



Correction to: Evaluation of optimum temperature for the early larval growth of Japanese eel in captivity

Mari Kuroki¹ · Akihiro Okamura² · Yoshiaki Yamada² · Shunsuke Hayasaka¹ · Katsumi Tsukamoto^{1,2}

Published online: 28 June 2019
© Japanese Society of Fisheries Science 2019, corrected publication 2019

Correction to: Fisheries Science
<https://doi.org/10.1007/s12562-019-01317-z>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In the original publication Table 2 is incorrectly published with inappropriate format. The correct version of Table 2 is presented in this correction.

The original article has been corrected.

Table 2 Otolith diameter (mean ± standard deviation) and growth of cultured larvae at different water temperatures, and estimated ambient temperature (bold) for wild larvae based on the relationship between otolith growth and temperature

Larvae	Target temperature (°C)	Mean temperature (°C)	Days post-hatching (dph)	Otolith diameter (μm)	Days post-hatching (dph)	Otolith diameter (μm)	Otolith growth (μm/day)
Cultured	19	20.1				29.53 ± 2.97	0.70
	21	21.7				32.08 ± 2.97	0.79
	23	23.4	0	10.73 ± 1.18	27	33.33 ± 3.27	0.84
	25	25.0				36.47 ± 3.42	0.95
	27	26.6				38.80 ± 2.96	1.04
Wild	–	23.2	0	10.49 ± 1.90	30	36.15 ± 4.88	0.86

The original article can be found online at <https://doi.org/10.1007/s12562-019-01317-z>.

✉ Mari Kuroki
mari.kuroki@aqua.fs.a.u-tokyo.ac.jp

¹ Department of Aquatic Bioscience, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1, Yayoi, Bunkyo, Tokyo 113-8657, Japan

² IRAGO Institute Co., Ltd., Tahara, Aichi 441-3605, Japan