



## Correction to: Low-Grade Inflammation Aggravates Rotenone Neurotoxicity and Disrupts Circadian Clock Gene Expression in Rats

Huan Li<sup>1,2</sup> · Sheng Song<sup>3</sup> · Yuan Wang<sup>1</sup> · Chun Huang<sup>1</sup> · Feng Zhang<sup>1</sup> · Jie Liu<sup>1,3</sup> · Jau-Shyong Hong<sup>3</sup>

Published online: 21 November 2018  
© Springer Science+Business Media, LLC, part of Springer Nature 2018

### Correction to: Neurotoxicity Research

<https://doi.org/10.1007/s12640-018-9968-1>

The original version of this article contained mistakes, and the authors would like to correct them. Some parts of the image in Figure 5 were missing. The correct Figure 5 is shown at the next page.

The original article has been corrected.

---

The online version of the original article can be found at <https://doi.org/10.1007/s12640-018-9968-1>


---

✉ Jie Liu  
Jie@liuonline.com

<sup>1</sup> Key Laboratory of Basic Pharmacology of Ministry of Education and Joint International Research Laboratory of Ethnomedicine of Ministry of Education, Zunyi Medical University, Zunyi, Guizhou, China

<sup>2</sup> Life Sciences and Technology Institute, China Pharmaceutical University, Nanjing 210023, China

<sup>3</sup> National Institute of Environmental Health Sciences, NIH, Research Triangle Park, NC 27709, USA



**Fig. 5** Effects of LPS and ROT on the expressions of circadian clock feedback gene Per1, Per2, Cry1, and Cry2. Rats were given LPS (5 mg/kg, i.p.  $\times 1$ ), and 7 months later 20 injections of rotenone (0.5 mg/kg, s.c., 5 times/week for 4 weeks). Two weeks after the last ROT treatment, rats

were sacrificed and cortex was collected for analysis. Values are mean  $\pm$  SEM ( $n = 6-9$ ). \*Significantly different from the control group, # Significantly different from the LPS group,  $p < 0.05$