



CORRECTION

Correction to: Gene activation in human cells using CRISPR/Cpf1-p300 and CRISPR/Cpf1-SunTag systems

Xin Zhang, Wei Wang, Lin Shan, Le Han, Shufeng Ma, Yan Zhang, Bingtao Hao, Ying Lin[⊠], Zhili Rong[⊠]

Cancer Research Institute, School of Basic Medical Sciences, Southern Medical University, Guangzhou 510515, China
Correspondence: linying0216@smu.edu.cn (Y. Lin), rongzhili@smu.edu.cn (Z. Rong)

CORRECTION TO: PROTEIN CELL 2018, 9(4):380–383 HTTPS://DOI.ORG/10.1007/S13238-017-0491-6

In the original publication the Supplementary Material and Fig. 2 are incorrect. The correct version of Supplementary Material and Fig. 2 are provided in this correction article. The text HBG2 appearing in the article should be read as HBG1.

The original article can be found online at https://doi.org/10.1007/s13238-017-0491-6.

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s13238-018-0585-9) contains supplementary material, which is available to authorized users.




Figure 2. Simultaneously transcriptional activation of multiple endogenous genes using either dLbCpf1-p300core or dLbCpf1-SunTag system with a single gRNA for each gene. (A) Relative mRNA expression of MYOD, IL1RN, and HBG1 revealed by quantitative real-time PCR, in HEK293Tcells co-transfected with dCpf1-p300core fusion proteins and four single gRNAs or pooled sets of all four single gRNAs targeting each promoter region of target genes. (B) Relative mRNA expression of HBG1 revealed by quantitative RT-PCR, in HEK293T cells co-transfected with dCpf1-p300core fusion proteins and four single gRNAs or pooled sets of all four single gRNAs targeting the enhancer region (HS2 region) of HBG1 gene. (C) Relative mRNA expression of MYOD, HBG1, and IL1RN revealed by quantitative RT-PCR, in HEK293Tcells co-transfected with dCpf1-p300core fusion proteins and three gRNAs targeting each promoter region of target genes. (D) Relative mRNA expression of MYOD, HBG1, and IL1RN revealed by quantitative RT-PCR, in HEK293T cells co-transfected with dLbCpf1 (M925)-SunTag and three gRNAs targeting each promoter region of target genes. For C and D, gRNA1, gRNA2 and gRNA1 were used for MYOD, HBG1 and IL1RN, respectively. For (A–D), mean value are presented with S.D. (n = 3). Tukey-test, P < 0.05 compared to cells transfected with dCpf1-p300core or dLbCpf1(M925)-SunTag only, n = 3 independent experiments.

OPEN ACCESS

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and

reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.