



Correction to: Three-dimensional bone formation including vascular networks derived from dental pulp stem cells in vitro

Miho Watanabe¹ · Akihiro Ohyama^{2,3} · Hiroshi Ishikawa^{2,3} · Akira Tanaka^{1,4}

Published online: 15 April 2019
© Japan Human Cell Society and Springer Japan KK, part of Springer Nature 2019

Correction to: Human Cell
<https://doi.org/10.1007/s13577-018-00228-y>

The authors would like to correct the error in the publication of the original article. The correction detail is given below:

The first sentence beneath the subheading “Preparation of endothelial cell beads (EC beads)” on page 3 should read as “To prepare endothelial cell beads (EC beads), 4 μ l atelocollagen solution containing 1×10^5 peripheral blood cells and 500 ng/ml vascular endothelial growth factor was dropped into mineral oil at 37 °C.”

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13577-018-00228-y>.

✉ Miho Watanabe
wmiho@ngt.ndu.ac.jp

- ¹ Oral and Maxillofacial Surgery, Field of Oral and Maxillofacial Surgery and Systemic Medicine, Course of Clinical Science, Nippon Dental University Graduate School of Life Dentistry at Niigata, 1-8 Hamaura-cho, Chuo-ku, Niigata 951-8580, Japan
- ² Department of NDU Life Sciences, Nippon Dental University School of Life Dentistry at Tokyo, 1-9-20 Fujimi, Chiyoda-ku, Tokyo 102-8159, Japan
- ³ Laboratory of Clinical Regenerative Medicine, Department of Neurosurgery, Faculty of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba-shi 305-8575, Ibaraki, Japan
- ⁴ Department of Oral and Maxillofacial Surgery, Nippon Dental University School of Life Dentistry at Niigata, 1-8 Hamaura-cho, Chuo-ku, Niigata 951-8580, Japan