

CORRECTION

## Correction to: The Expression of Immunomodulation-Related Cytokines and Genes of Adipose- and Bone Marrow-Derived Human Mesenchymal Stromal Cells from Early to Late Passages

Chin Hee Mun<sup>1,2,3</sup> · Mi-Il Kang<sup>4</sup> · Yong Dae Shin<sup>1,2</sup> · Yeseul Kim<sup>1</sup> · Yong-Beom Park<sup>1,2,3</sup>

Published online: 12 January 2019

© The Korean Tissue Engineering and Regenerative Medicine Society and Springer Nature B.V. 2019

**Correction to: Tissue Eng Regen Med (2018) 15(6):771–779**  
<https://doi.org/10.1007/s13770-018-0147-5>

Unfortunately, the online published article has error in Figure 3. The name of protein, “Gelectin-1” should be corrected to “Galectin-3”. The corrected Fig. 3 is placed in the following page.

---

The original article can be found online at  
<https://doi.org/10.1007/s13770-018-0147-5>.


✉ Yong-Beom Park  
yongbpark@yuhs.ac

<sup>1</sup> Division of Rheumatology, Department of Internal Medicine, and Department of Medical Sciences, Institute for Immunology and Immunological Disease, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea

<sup>2</sup> BK21 Plus Project, Department of Medical Sciences, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea

<sup>3</sup> Severance Biomedical Science Institute, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea

<sup>4</sup> Division of Rheumatology, Department of Internal Medicine, Dankook University College of Medicine, 201 Manghyang-ro, Dongnam-gu, Cheonan, Chung Nam, Republic of Korea



◀**Fig. 3** Gene expressions relevant to immune modulation of AD- and BM-MSCs through passages. **A–L** TSG-6 and HLA-G expressions were decreased, whereas galectin-1 and galectin-3 expressions were increased or maintained throughout passages in AD-MSCs and BM-MSCs. Data are shown as normalized gene expression (**A–B**, **D–E**, **G–H**, **J–K**) compared to the housekeeping gene, GAPDH, and as relative gene expression (**C**, **F**, **I**, **L**) compared to AD- and BM-MSC P4. Each experiment was conducted in triplicate. All values are the mean  $\pm$  SEM. \* $p < 0.05$  was considered statistically significant versus P4 of each origin, ns means not significant