

Spinning pulsating strings in $(AdS_5 \times S^5)_\kappa$

Sorna Prava Barik^a, Kamal L. Panigrahi^b, Manoranjan Samal^c

Department of Physics, Indian Institute of Technology Kharagpur, Kharagpur 721 302, India

Received: 1 March 2018 / Accepted: 19 March 2018 / Published online: 3 April 2018
© The Author(s) 2018

Abstract We study a general class of spinning pulsating strings in $(AdS_5 \times S^5)_\kappa$ background. For these family of solitons, we examine the scaling relation between the energy, spin or angular momentum. We verify that in $\kappa \rightarrow 0$ limit these relations reduce to the undeformed $AdS_5 \times S^5$ case. We further study an example of a string which is spinning in the κ -deformed AdS_5 and S^5 simultaneously and find out the scaling relation among various conserved charges.

1 Introduction

The AdS/CFT correspondence relates string states on AdS and gauge invariant operators in the gauge theory side. The most studied example of the AdS/CFT duality is the one between spectrum of closed superstrings (supergravity) in $AdS_5 \times S^5$ background and gauge invariant operators in four dimensional $\mathcal{N} = 4$ Supersymmetric Yang–Mills (SYM) theory based on the gauge group $SU(N)$ [1–3]. A remarkable feature of AdS/CFT duality is an underlying integrability [4] structure on both side, which provides an important tool for finding the spectrum on both sides and many significant properties have been revealed based on exact computations. To understand the structure of full string spectrum, one need to identify the classical solitonic solutions of $AdS_5 \times S^5$ sigma model carrying global charges. The Yang–Mills theory itself can be mapped to a integrable spin-chain system [5]. The basic idea of relating all string states to precise dual gauge theory operators is a tough job due to presence of infinite tower of string solutions on string theory side. One probable way out of this problem is that in the large angular momentum or large R-charge limit both sides of the duality become more tractable. One of the advantages of this limit is that the anomalous dimension of operators in the SYM theory

can be related to the dispersion relation between conserved charges of spinnings and pulsating strings in the large charge limit.

In this context, a large variety of rotating and spinning strings has been studied in $AdS_5 \times S^5$ precisely and also have been mapped to dual spin-chain excitations. These include the already well studied giant magnon, folded strings and spiky strings solutions and the gauge theory duals have been analyzed in great detail. In spinning string case the highly excited string states corresponds to the gauge theory operators with small anomalous dimension. This type of strings is the generalization of the folded [6,7] and spiky [8] strings with single spin in AdS_3 part of AdS_5 . The semiclassical multi spinning string states (strings spinning in AdS_5) have also been found to be dual to certain trace operators [9]. It has been shown that for these solutions, the string states are unstable for large charges [10]. On the other hand, the circular pulsating string solutions have been less explored. Pulsating strings were first introduced in [11] where they were expected to correspond to certain highly excited sigma model operators and later on were generalized to [12–14]. In [11] and [15], pulsating string solutions in AdS_5 and S^5 respectively have been worked out separately where as in [16], a string rotating and at the same time oscillating in AdS_5 have been derived. It has been shown that the addition of oscillation to spinning strings improve the stability of the string states [17]. An interesting class of solutions were proposed in [18] which generalized some of the earlier pulsating and spinning string solutions by looking at strings which are straight and spinning in one direction but circular and pulsating in another, and with a non-trivial coupling between the two in $AdS_5 \times S^5$ background. Even though the exact gauge theory operators corresponding to these class of string states are still unknown, they are interesting in their own right. In this paper, we wish to generalize such spinning pulsating strings in $(AdS_5 \times S^5)_\kappa$ background, the sigma model associated to which retains integrability of the original model.

^a e-mail: sorna@phy.iitkgp.ernet.in

^b e-mail: panigrahi@phy.iitkgp.ernet.in

^c e-mail: manoranjan@phy.iitkgp.ernet.in

To explore the integrability beyond the usual $AdS_5 \times S^5$, it is natural to consider the integrable deformation of the background to study the string motion as well as the underlying gauge theory, if any. Following the earlier proposals about Yang–Baxter deformation made in [19–21], a one-parameter deformed $AdS_5 \times S^5$ supercoset model was constructed in [22–24]. The background string metric and the NS–NS 2-form corresponding to deformed structure has been worked out [25]. The symmetry group $SO(2, 4) \times SO(6)$ of $AdS_5 \times S^5$ reduced to its Cartan subgroup $[U(1)]^6$ and it produced a nice laboratory to study string motion. Given the integrable nature of the deformed background, various rigidly rotating and pulsating strings have been investigated in detail [26–31]. Since the background suffers from the presence of a singularity surface in the AdS space, a new coordinate system to handle this has been developed [32]. The folded GKP like string solutions of [32] generalized to N-spike string solutions in [33]. In these cases, however, it was found out that in the long string limit, where the strings touch the singularity surface, the expression for cusp anomalous dimension does not reduce to the undeformed one in the $\kappa \rightarrow 0$ limit and a possible explanation was provided in [32]. Not only classical string solutions, various minimal surfaces and Wilson loops in this background have been found in [34]. In some recent works [35,36], conformal twist and non-commutative gauge theory have been studied in Yang–Baxter deformed $AdS_5 \times S^5$ background.

In the present paper, we are interested in a non-trivial string solitons in deformed $AdS_5 \times S^5$ background with κ being the deformed parameter. The strings, we consider here are straight and spinning in one sub-space while it is circular and pulsating in another. In each case, we obtain the exact form of string solutions in terms of elliptic functions and discuss their properties. We construct the scaling relations between energy and charges in various limits and compare the result with the undeformed results. We would like to mention in passing that the dual gauge theory of the κ -deformed background is not yet known and hence the operator dual to these class of solutions is not apriori clear. One of the main reasons is the lack of our understanding of incorporating the deformed parameter into the operators which would be present in the unknown gauge theory. Nevertheless studying the string states will surely enhance our understanding of the string theory in integrable deformed background. The rest of the paper is organized as follows. In Sect. 2, we outline the background that we are interested in for studying the string solutions. In Sect. 3, we study the strings which are pulsating in AdS_5 subspace and is spinning in S^5 simultaneously. We find out the scaling relation among the energy and the angular momentum. We study the behavior of the long and short strings in this case and have checked that they indeed reduce to the similar relations in the undeformed $AdS_5 \times S^5$ case. Section 4 is devoted to the study of strings which pulsate

in S^5 and at the same time spins along the deformed AdS_5 . We have looked at some interesting spinning string solution which spins along both deformed AdS_5 and S^5 subspace and write down the energy spin relationship in Sect. 5. In Sect. 6 we conclude with some discussions.

2 The κ deformed $(AdS_5 \times S^5)_\kappa$ geometry

The so called κ deformed $(AdS_5 \times S^5)_\kappa$ background is

$$ds^2_{AdS_5} = -h(\rho)dt^2 + f(\rho)d\rho^2 + \rho^2 (u(\rho, \psi)(d\psi^2 + \cos^2 \psi d\psi_1^2) + \sin^2 \psi d\psi_2^2), \tag{2.1}$$

$$ds^2_{S^5} = \tilde{h}(r)d\varphi^2 + \tilde{f}(r)dr^2 + r^2 (\tilde{u}(r, \phi)(d\phi^2 + \cos^2 \phi d\phi_1^2) + \sin^2 \phi d\phi_2^2), \tag{2.2}$$

where

$$h(\rho) = \frac{1 + \rho^2}{1 - \kappa^2 \rho^2}, \quad f(\rho) = \frac{1}{(1 + \rho^2)(1 - \kappa^2 \rho^2)},$$

$$u(\rho, \psi) = \frac{1}{1 + \kappa^2 \rho^4 \sin^2 \psi},$$

and

$$\tilde{h}(r) = \frac{1 - r^2}{1 + \kappa^2 r^2}, \quad \tilde{f}(r) = \frac{1}{(1 - r^2)(1 + \kappa^2 r^2)},$$

$$\tilde{u}(r, \phi) = \frac{1}{1 + \kappa^2 r^4 \sin^2 \phi},$$

supported by the following NS–NS B field

$$B_{\psi_1 \psi} = \frac{1}{2} \kappa^2 \rho^4 \sin 2\psi u(\rho, \psi),$$

$$B_{\phi_1 \phi} = -\frac{1}{2} \kappa^2 r^4 \sin 2\phi \tilde{u}(r, \phi), \tag{2.3}$$

where $\kappa \in [0, \infty)$ is the deforming parameter. It can be seen that for $\kappa = 0$, the above geometry reduces to the original $AdS_5 \times S^5$. It is convenient to write the above metric and the NS–NS B-field components in global coordinates which is obtained by the following co-ordinate transformations: $\rho \rightarrow \sinh \rho$ and $r \rightarrow \cos \theta$. The metric in the global coordinates looks like

$$ds^2_{AdS_5} = -\frac{\cosh^2 \rho}{1 - \kappa^2 \sinh^2 \rho} dt^2 + \frac{d\rho^2}{1 - \kappa^2 \sinh^2 \rho} + \frac{\sinh^2 \rho d\psi^2}{1 + \kappa^2 \sinh^4 \rho \sin^2 \psi} + \frac{\sinh^2 \rho \cos^2 \psi d\psi_1^2}{1 + \kappa^2 \sinh^4 \rho \sin^2 \psi} + \sinh^2 \rho \sin^2 \psi d\psi_2^2, \tag{2.4}$$

$$ds^2_{S^5} = \frac{\sin^2 \theta d\varphi^2}{1 + \kappa^2 \cos^2 \theta} + \frac{d\theta^2}{1 + \kappa^2 \cos^2 \theta}$$

$$\begin{aligned}
 & + \frac{\cos^2 \theta d\phi^2}{1 + \kappa^2 \cos^4 \theta \sin^2 \phi} + \frac{\cos^2 \theta \cos^2 \phi d\phi_1^2}{1 + \kappa^2 \cos^4 \theta \sin^2 \phi} \\
 & + \cos^2 \theta \sin^2 \phi d\phi_2^2. \tag{2.5}
 \end{aligned}$$

We would like to study a general class of spinning pulsating string in this background. The deformed bosonic string action in conformal gauge can be written as

$$S = \hat{T} \int d^2\sigma \left(\sqrt{-h} h^{\alpha\beta} G_{\mu\nu} \partial_\alpha X^\mu \partial_\beta X^\nu + \epsilon^{\alpha\beta} B_{\mu\nu} \partial_\alpha X^\mu \partial_\beta X^\nu \right), \tag{2.6}$$

where $\hat{T} = \frac{1}{4\pi\alpha'} \sqrt{1 + \kappa^2}$ is the effective string tension.

3 Pulsating in $(\text{AdS}_5)_\kappa$ and spinning in $(S^5)_\kappa$

In this section, we consider a string which is spinning in S^5 and pulsating in AdS_5 of the deformed background. We use the following ansatz

$$\begin{aligned}
 t &= t(\tau), \quad \rho = \rho(\tau), \quad \psi = \frac{\pi}{2}, \quad \psi_2 = \sigma, \\
 \varphi &= \varphi(\tau), \quad \theta = \theta(\sigma), \quad \phi = \frac{\pi}{2}, \quad \phi_2 = \text{const}. \tag{3.1}
 \end{aligned}$$

Taking the above ansatz the Polyakov action in conformal gauge takes the form,

$$\begin{aligned}
 S &= \hat{T} \int d^2\sigma \left(-\frac{\cosh^2 \rho \dot{t}^2}{1 - \kappa^2 \sinh^2 \rho} + \frac{\dot{\rho}^2}{1 - \kappa^2 \sinh^2 \rho} \right. \\
 &\quad \left. - \sinh^2 \rho + \frac{\sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} \dot{\varphi}^2 - \frac{\theta'^2}{1 + \kappa^2 \cos^2 \theta} \right). \tag{3.2}
 \end{aligned}$$

We notice that for such ansatz the B-fields do not contribute to the action. The equation of motion for t and φ can be found as

$$\dot{t} = \frac{c_0(1 - \kappa^2 \sinh^2 \rho)}{\cosh^2 \rho}, \quad \dot{\varphi} = v, \tag{3.3}$$

where c_0 and v are the integration constants. From the equation motion of the φ , we find that the string rotates in φ direction with angular velocity v . Now, the equations of motion for ρ and θ can be written as

$$\begin{aligned}
 & 2\ddot{\rho}(1 - \kappa^2 \sinh^2 \rho) + \kappa^2 \sinh^2 \rho \dot{\rho}^2 \\
 & + \sinh^2 2\rho \left((1 + \kappa^2) \dot{t}^2 + (1 - \kappa^2 \sinh^2 \rho)^2 \right) = 0, \tag{3.4}
 \end{aligned}$$

and

$$2\theta'' \left(1 + \kappa^2 \cos^2 \theta \right) + \kappa^2 \theta'^2 \sin 2\theta + v^2 \sin 2\theta (1 + \kappa^2) = 0. \tag{3.5}$$

These equations of motion are to be supported by the conformal gauge constraints

$$G_{mn} \left(\partial_\tau X^m \partial_\tau X^n + \partial_\sigma X^m \partial_\sigma X^n \right) = 0 \tag{3.6}$$

$$G_{mn} \partial_\tau X^m \partial_\sigma X^n = 0. \tag{3.7}$$

From (3.6), we get

$$\begin{aligned}
 & -\frac{c_0^2(1 - \kappa^2 \sinh^2 \rho)}{\cosh^2 \rho} + \frac{\dot{\rho}^2}{1 - \kappa^2 \sinh^2 \rho} + \sinh^2 \rho \\
 & + \frac{v^2 \sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} + \frac{\theta'^2}{1 + \kappa^2 \cos^2 \theta} = 0. \tag{3.8}
 \end{aligned}$$

The above equation is a coupled equation of $\rho(\tau)$ and $\theta(\sigma)$. However without loss of generality we can separate the above equation as

$$\begin{aligned}
 & -\frac{c_0^2(1 - \kappa^2 \sinh^2 \rho)}{\cosh^2 \rho} + \frac{\dot{\rho}^2}{1 - \kappa^2 \sinh^2 \rho} \\
 & + \sinh^2 \rho + c_1^2 = 0, \tag{3.9}
 \end{aligned}$$

$$\frac{v^2 \sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} + \frac{\theta'^2}{1 + \kappa^2 \cos^2 \theta} - c_1^2 = 0, \tag{3.10}$$

where c_1 is a constant. One can check that the above equations are consistent with the equations of motion ρ and θ . Now solving these equations we find the following solution of ρ and θ

$$\sinh \rho = \sqrt{\frac{R_- \text{sn}^2 \left((1 + \kappa^2 c_1^2) \sqrt{R_+} \tau \left| \frac{R_-}{R_+} \right. \right)}{1 + \kappa^2 \left[1 + R_- \text{sn}^2 \left((1 + \kappa^2 c_1^2) \sqrt{R_+} \tau \left| \frac{R_-}{R_+} \right. \right) \right]}}, \tag{3.11}$$

$$\sin \theta = \sqrt{\frac{c_1^2(1 + \kappa^2)}{v^2 + \kappa^2 c_1^2}} \text{sn} \left(\sqrt{v^2 + \kappa^2 c_1^2} \sigma \left| \frac{c_1^2(1 + \kappa^2)}{v^2 + \kappa^2 c_1^2} \right. \right), \tag{3.12}$$

with

$$R_\pm = \frac{-1 + \kappa^2 c_0^2 - c_1^2(1 + 2\kappa^2) \pm \sqrt{(c_1^2 - 1)^2 + \kappa^4 c_0^4 + 2c_0^2(2 + (1 + c_1^2)\kappa^2)}}{2 + 2\kappa^2 c_1^2}. \tag{3.13}$$

Now we take periodicity condition of θ which gives, the range $0 \leq \sigma \leq 2\pi$ is divided into four equal segments: in the first segment i.e. for $0 \leq \sigma \leq \frac{\pi}{2}$, the solution of $\theta(\sigma)$ increases from zero to its maximal value $\theta_{max} = \sin^{-1} \sqrt{\frac{c_1^2(1 + \kappa^2)}{v^2 + \kappa^2 c_1^2}}$, then it decreases to zero in the $\frac{\pi}{2} \leq \sigma \leq \pi$ segment and continues the similar pattern for $\pi \leq \sigma \leq 2\pi$ region.

$$2\pi = \int_0^{2\pi} d\sigma = 4 \int_0^{\theta_{max}} \frac{d\theta}{\theta'}$$

$$= 4 \int_0^{\theta_{max}} \frac{d\theta}{\sqrt{c_1^2(1 + \kappa^2) - \sin^2 \theta (v^2 + c_1^2 \kappa^2)}}. \tag{3.14}$$

Using the definition of complete elliptic integral of first kind (\mathbf{K}), we get

$$2\pi \sqrt{v^2 + \kappa^2 c_1^2} = 4\mathbf{K} \left[\frac{c_1^2(1 + \kappa^2)}{v^2 + \kappa^2 c_1^2} \right]. \tag{3.15}$$

The conserved charges can be evaluated as

$$E = \int_0^{2\pi} \frac{\sqrt{1 + \kappa^2} c_0}{2\pi \alpha'} d\sigma = \frac{c_0}{\alpha'} \sqrt{1 + \kappa^2}, \tag{3.16}$$

$$\mathcal{J} = \frac{v \sqrt{1 + \kappa^2}}{2\pi \alpha'} \int_0^{2\pi} \frac{\sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} d\sigma. \tag{3.17}$$

Let us define two parameters $A = \sinh \rho_{max} = \sqrt{R_-}$ and $B = \sin \theta_{max} = \sqrt{\frac{c_1^2(1 + \kappa^2)}{v^2 + \kappa^2 c_1^2}}$. Now we want to express the conserved charges in terms of these two parameters so that we can analyze both ‘‘short string’’ and ‘‘long string’’ limits. Let us first find out the other parameters in terms A and B . Using the periodicity condition (3.15), we get

$$c_1 = \frac{2B\mathbf{K}(B^2)}{\pi \sqrt{1 + \kappa^2}} \quad \text{and} \quad v = \frac{2}{\pi} \sqrt{\frac{1 + \kappa^2(1 - B^2)}{1 + \kappa^2}} \mathbf{K}(B^2). \tag{3.18}$$

Solving (3.13) for c_0 and then substituting the value of c_1 from the above equation, we get the conserved energy in the following form

$$E = \sqrt{\frac{(1 + A^2) [(1 + \kappa^2 + \kappa^2 A^2) 4B^2 \mathbf{K}^2(B^2) + \pi^2 A^2 (1 + \kappa^2)]}{\alpha'^2 \pi^2 (1 + \kappa^2 (1 + A^2))}}. \tag{3.19}$$

Similarly substituting the value of $\sin \theta$ in (3.17) and then using (3.18), we find the expression of angular momentum in terms B as

$$\mathcal{J} = \frac{2\sqrt{1 + \kappa^2 - \kappa^2 B^2}}{\alpha' \pi \kappa^2} \left[\mathbf{\Pi} \left(\frac{\kappa^2 B^2}{1 + \kappa^2}, B^2 \right) - \mathbf{K}(B^2) \right]. \tag{3.20}$$

Now we reach in a position where we can apply short string limit or long string limit to establish the relation between conserved charges. Let us discuss these two situations one by one.

Short string limit

Consider the short-string solution in S^5 which correspond to $\sin \theta_{max} \ll 1$ or $B \ll 1$. For which the leading order term in

energy and angular momentum turn out to be

$$E \approx \sqrt{\frac{(1 + A^2) [(1 + \kappa^2 + \kappa^2 A^2) B^2 + A^2 (1 + \kappa^2)]}{\alpha'^2 (1 + \kappa^2 (1 + A^2))}}, \tag{3.21}$$

$$\mathcal{J} \approx \frac{B^2}{2\alpha' \sqrt{1 + \kappa^2}}. \tag{3.22}$$

Therefore the scaling relation between E and \mathcal{J} becomes,

$$E \approx \sqrt{\frac{(1 + A^2) [2\alpha' (1 + \kappa^2 + \kappa^2 A^2) \sqrt{1 + \kappa^2} \mathcal{J} + A^2 (1 + \kappa^2)]}{\alpha'^2 (1 + \kappa^2 (1 + A^2))}}. \tag{3.23}$$

Now let us consider two sub cases depending upon the nature of the oscillation of string in AdS_5 part. For $A \ll 1$ which corresponds to small oscillation at center AdS, the relation between E and \mathcal{J} becomes,

$$\alpha' E^2 \approx 2J \sqrt{1 + \kappa^2}. \tag{3.24}$$

For $\kappa \rightarrow 0$, we get $\alpha' E^2 \approx 2\mathcal{J}$, which is a well known result for undeformed $AdS_5 \times S^5$ as found in [18,37]. Similarly, for $A \gg 1$, we write the scaling relation by expanding it for small κ ,

$$E = \left(\frac{A^2}{\alpha'} + \frac{1}{2\alpha'} + \mathcal{J} \right) + \left(-\frac{A^4}{2\alpha'} + \left(\mathcal{J} - \frac{1}{2\alpha'} \right) \frac{A^2}{2} + \left(\frac{3\mathcal{J}}{4} + \frac{1}{16\alpha'} \right) \right) \kappa^2 + \mathcal{O}(\kappa^4). \tag{3.25}$$

We can notice, as expected, the zeroth order κ term matches with its corresponding undeformed case as found in [18].

Long string limit

For long string in S^5 , the string extends to the equator so $B \approx 1$ for which E and \mathcal{J} become

$$E \approx \sqrt{\frac{(1 + A^2) \left((1 + \kappa^2 + \kappa^2 A^2) \log^2 \frac{16}{1 - B^2} + \pi^2 (1 + \kappa^2) A^2 \right)}{\alpha'^2 \pi^2 (1 + \kappa^2 (1 + A^2))}}, \tag{3.26}$$

$$\mathcal{J} \approx \frac{1}{\alpha' \pi} \log \frac{16}{1 - B^2} - \frac{6 + 2\kappa^2}{3\pi \alpha'}. \tag{3.27}$$

Therefore the relation between E and \mathcal{J} turns out to be

$$E \approx \sqrt{\frac{(1 + A^2) \left[(1 + \kappa^2 + \kappa^2 A^2) (\alpha' \pi \mathcal{J} + 2 + \frac{2}{3} \kappa^2)^2 + \pi^2 (1 + \kappa^2) A^2 \right]}{\alpha'^2 \pi^2 [1 + \kappa^2 (1 + A^2)]}}. \tag{3.28}$$

For small oscillation in AdS i.e $A \ll 1$, we get

$$E - \mathcal{J} \approx \frac{2}{\pi\alpha'} \left(1 + \frac{\kappa^2}{3}\right) + \frac{\mathcal{J}A^2}{2}. \tag{3.29}$$

On the other hand, for $A \gg 1$ we have to distinguish two different scenarios,

(i) for $\alpha'J \ll A$, we get the scaling relation as

$$E = \left(\frac{A^2}{\alpha'} + \frac{\alpha' \mathcal{J}^2}{2}\right) + \left(-\frac{A^4}{2\alpha'} + \left(\frac{\mathcal{J}}{\alpha'} + \frac{\alpha' \mathcal{J}^2}{4}\right) \frac{A^2}{2}\right) \kappa^2 + \mathcal{O}(\kappa^4). \tag{3.30}$$

(ii) for $\alpha'J \gg A$, we get

$$E = A\mathcal{J} + \frac{A^3}{2\alpha'^2 \mathcal{J}} + \left(\frac{2A^2}{3\pi\alpha'} - \frac{A^3}{3\pi\alpha'^3 \mathcal{J}^2} + \left(-\frac{1}{2\alpha'^2 \mathcal{J}} + \frac{1}{\alpha'^3 \pi \mathcal{J}^2}\right) A^5\right) \kappa^2 + \mathcal{O}(\kappa^4). \tag{3.31}$$

We can notice that for $\kappa \rightarrow 0$ the above results agree with their undeformed case as found in [6, 18, 37].

4 Spinning in $(AdS_5)_\kappa$ and pulsating in $(S^5)_\kappa$

In this section, we consider a string which is spinning in AdS_5 and pulsating in S^5 of the deformed geometry. The corresponding string ansatz:

$$t = k\tau, \quad \rho = \rho(\sigma), \quad \psi = \frac{\pi}{2}, \quad \psi_2 = \omega\tau, \quad \theta = \theta(\tau), \quad \varphi = \sigma. \tag{4.1}$$

The Polyakov action for the above string ansatz can be written as

$$S_p = \hat{T} \int d^2\sigma \left(-\frac{k^2 \cosh^2 \rho}{1 - \kappa^2 \sinh^2 \rho} - \frac{\rho'^2}{1 - \kappa^2 \sinh^2 \rho} + \omega^2 \sinh^2 \rho + \frac{\dot{\theta}^2}{1 + \kappa^2 \cos^2 \theta} - \frac{\sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} \right). \tag{4.2}$$

The equations of motion for ρ and θ can be found as

$$2\rho''(1 - \kappa^2 \sinh^2 \rho) + \kappa^2 \rho'^2 \sinh 2\rho + \sinh 2\rho \left(\omega^2 (1 - \kappa^2 \sinh^2 \rho)^2 - k^2 (1 + \kappa^2) \right) = 0, \tag{4.3}$$

$$2\ddot{\theta}(1 + \kappa^2 \cos^2 \theta) + \dot{\theta}^2 \kappa^2 \sin 2\theta + (1 + \kappa^2) \sin 2\theta = 0. \tag{4.4}$$

The above equations of motion are supplemented with the following conformal gauge constraints:

$$-\frac{\cosh^2 \rho k^2}{1 - \kappa^2 \sinh^2 \rho} + \frac{\rho'^2}{1 - \kappa^2 \sinh^2 \rho} + \omega^2 \sinh^2 \rho + \frac{\dot{\theta}^2}{1 + \kappa^2 \cos^2 \theta} + \frac{\sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} = 0. \tag{4.5}$$

We can decouple the equations of ρ and θ as done in the previous section.

$$\frac{-\cosh^2 \rho k^2}{1 - \kappa^2 \sinh^2 \rho} + \frac{\rho'^2}{1 - \kappa^2 \sinh^2 \rho} + \omega^2 \sinh^2 \rho + c_1^2 = 0, \tag{4.6}$$

$$\frac{\dot{\theta}^2}{1 + \kappa^2 \cos^2 \theta} + \frac{\sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} - c_1^2 = 0, \tag{4.7}$$

where c_1 is a constant. We can also check that the above equations are consistent with equations of motion of ρ (4.3) and θ (4.4). Now it is easy to obtain the solution of ρ and θ from these equations. From (4.6) we get

$$\rho'^2 = \kappa^2 \omega^2 (\sinh^2 \rho - \sinh^2 \rho_-)(\sinh^2 \rho - \sinh^2 \rho_+), \tag{4.8}$$

where

$$z_{\pm} = \sinh^2 \rho_{\pm} = \frac{\omega^2 - k^2 - c_1^2 \kappa^2 \pm \sqrt{(\omega^2 - k^2 - c_1^2 \kappa^2)^2 - 4\kappa^2 \omega^2 (k^2 - c_1^2)}}{2\kappa^2 \omega^2}. \tag{4.9}$$

Now integrating (4.8), we get the following solution of ρ in terms of elliptic function

$$\coth^2 \rho = \frac{z_+ + 1}{z_+} \frac{\mathbf{dc}^2 \left(\kappa \omega \sqrt{z_+(1+z_-)} \sigma, \mathbf{k}^2 \right)}{\mathbf{k}^2}, \tag{4.10}$$

where \mathbf{dc} is Jacobi elliptic \mathbf{dc} function and $\mathbf{k}^2 = \frac{z_-(1+z_+)}{z_+(1+z_-)}$. Similarly the solution of θ can be found from (4.7) as

$$\sin \theta = \begin{cases} \mathbf{sn} \left(c_1 \sqrt{1 + \kappa^2} \tau, \frac{1}{B^2} \right) & \text{for } B > 1 \\ B \mathbf{sn} \left(\sqrt{1 + c_1^2 \kappa^2} \tau, B^2 \right) & \text{for } B < 1, \end{cases} \tag{4.11}$$

where $B = \frac{c_1 \sqrt{1 + \kappa^2}}{\sqrt{1 + c_1^2 \kappa^2}}$. The above solution of θ implies the pulsating nature of string along θ direction.

The positiveness of the left hand side of (4.8) put constraint on the values of ρ , which gives the allowed regions of ρ are

$$(i) \quad 0 \leq \rho \leq \rho_- \quad \text{with } \rho_- \leq \rho^* < \rho_s \tag{4.12}$$

$$(ii) \quad \rho_+ \leq \rho < \infty \quad \text{with } \rho^* \leq \rho_+ < \rho_s \tag{4.13}$$

where ρ^* is the value of ρ where $\rho_- = \rho_+$ and $\rho_s = \sinh^{-1} \left(\frac{1}{\kappa} \right)$ is the singularity in $(AdS_5)_\kappa$. In our case i.e

spinning string solution we study the string motion in the first region because it contains the center of AdS. Again since $\sinh^2 \rho_{\pm} \in \mathbb{R}_+$ it gives the following inequality,

$$\omega \geq \left(\frac{\sqrt{1 + \kappa^2} + \kappa \sqrt{1 - \frac{c_1^2}{\omega^2}}}{1 + \frac{c_1^2}{\omega^2} \kappa^2} \right) k. \tag{4.14}$$

Using this approximation and the periodicity condition (4.15), we get

$$\frac{k}{\omega} = \sqrt{z_- + \frac{B^2 \pi^2 (1 + z_-)}{4(1 + \kappa^2 - B^2 \kappa^2) \mathbf{K}^2 \left(\frac{z_-(1 + \kappa^2)}{z_- + 1} \right)}}. \tag{4.20}$$

Substituting these above result in (4.17) and (4.18) and then expanding for small z_- , the leading order terms in energy and spin expressions for sort string turn out to be

$$\begin{aligned} E &\approx \mathcal{A} \left(1 + \frac{(1 + \kappa^2 + B^2) z_-}{2B^2} - \frac{2(1 + \kappa^2)^2 - 12B^2 \kappa^2 (1 + \kappa^2) + B^4 (1 - 6\kappa^2 + 3\kappa^4)}{16B^4} z_-^2 \right), \\ S &\approx \frac{\sqrt{1 + \kappa^2}}{2\alpha'} \left[z_- + \frac{3(-1 + \kappa^2)}{8} z_-^2 \right], \end{aligned} \tag{4.21}$$

The periodicity condition can be written as,

$$2\pi = \int_0^{2\pi} d\sigma = 4 \int_0^{\rho_-} \frac{d\rho}{\rho'},$$

which gives the following relation between the parameters,

$$\pi \kappa \omega \sqrt{z_+(z_- + 1)} = 2\mathbf{K} \left(\frac{z_-(z_+ + 1)}{z_+(z_- + 1)} \right). \tag{4.15}$$

The conserved energy and spin can be evaluated as,

$$\begin{aligned} E &= \frac{k\sqrt{1 + \kappa^2}}{2\pi\alpha'} \int_0^{2\pi} \frac{\cosh^2 \rho}{1 - \kappa^2 \sinh^2 \rho} d\sigma \\ &= \frac{k\sqrt{1 + \kappa^2}}{2\pi\alpha'} \int_0^{2\pi} \frac{\cosh^2 \rho}{1 - \kappa^2 \sinh^2 \rho} \frac{d\rho}{\rho'}, \\ S &= \frac{\omega\sqrt{1 + \kappa^2}}{2\pi\alpha'} \int_0^{2\pi} \sinh^2 \rho d\sigma \\ &= \frac{\omega\sqrt{1 + \kappa^2}}{2\pi\alpha'} \int_0^{2\pi} \sinh^2 \rho \frac{d\rho}{\rho'}. \end{aligned} \tag{4.16}$$

Substituting the value of ρ' from (4.8), we get,

$$E = \frac{2k\sqrt{1 + \kappa^2}}{\pi\alpha'\kappa\omega\sqrt{z_+(z_- + 1)}} \mathbf{\Pi} \left(\frac{z_-(1 + \kappa^2)}{z_- + 1}, \frac{z_-(z_+ + 1)}{z_+(z_- + 1)} \right), \tag{4.17}$$

$$\begin{aligned} S &= \frac{2\sqrt{1 + \kappa^2}}{\pi\alpha'\kappa\sqrt{z_+(z_- + 1)}} \\ &\times \left(\mathbf{\Pi} \left(\frac{z_-}{z_- + 1}, \frac{z_-(z_+ + 1)}{z_+(z_- + 1)} \right) - \mathbf{K} \left(\frac{z_-(z_+ + 1)}{z_+(z_- + 1)} \right) \right), \end{aligned} \tag{4.18}$$

Short string limit

As the maximum value of ρ is ρ_- , for short string limit $\rho_- \rightarrow 0$ or $z_- \rightarrow 0$. For which

$$z_- \approx \frac{k^2 - c_1^2}{\omega^2} \text{ and } z_+ \approx \frac{1}{\kappa^2} \text{ with } \omega \gg k. \tag{4.19}$$

with $\mathcal{A} = \frac{B\sqrt{1 + \kappa^2}}{\alpha'\sqrt{1 + \kappa^2 - B^2 \kappa^2}}$. Therefore the expression for energy in terms of spin can be written as

$$E^2 \approx \frac{(1 + \kappa^2)}{\alpha'^2(1 + \kappa^2 - B^2 \kappa^2)} \left(B^2 + \frac{2\alpha'(1 + B^2 + \kappa^2)}{\sqrt{1 + \kappa^2}} S \right). \tag{4.22}$$

For small oscillation i.e. $B \rightarrow 0$

$$E^2 \approx 2 \frac{\sqrt{1 + \kappa^2}}{\alpha'} S. \tag{4.23}$$

For $\kappa \rightarrow 0$ the above results reduces to result obtained in undeformed case [18,38].

Long string limit

Now let us consider the long string solution, for which the length of string becomes maximum i.e. $\rho_- \approx \rho^*$. Now using (4.8), (4.12) and (4.14), we get

$$z_- = \sinh^2 \rho_- = \sqrt{\left(1 + \frac{1}{\kappa^2} \right) \left(1 - \frac{c_1^2}{\omega^2} \right)} - 1, \tag{4.24}$$

$$\frac{k}{\omega} = \sqrt{1 + \kappa^2} - \kappa \sqrt{1 - \frac{c_1^2}{\omega^2}}. \tag{4.25}$$

We can also notice that for long string solution ρ_- becomes very nearer to ρ_+ . Therefore taking $\frac{z_-}{z_+} = 1 - \varepsilon$ where $\varepsilon \ll 1$, the expressions for energy and spin for long string limit become

$$\begin{aligned} E &= \frac{2k\sqrt{1 + \kappa^2}}{\pi\alpha'\kappa\omega\sqrt{z_-(z_- + 1)}} \sqrt{1 - \varepsilon} \mathbf{\Pi} \\ &\times \left(\frac{z_-(1 + \kappa^2)}{z_- + 1}, 1 - \frac{\varepsilon}{1 + z_-} \right), \\ S &= \frac{2\sqrt{1 + \kappa^2}}{\pi\alpha'\kappa\sqrt{z_-(z_- + 1)}} \sqrt{1 - \varepsilon} \end{aligned} \tag{4.26}$$

$$\times \left(\mathbf{\Pi} \left(\frac{z_-}{z_- + 1}, 1 - \frac{\varepsilon}{1 + z_-} \right) - \mathbf{K} \left(1 - \frac{\varepsilon}{1 + z_-} \right) \right) \tag{4.27}$$

Now using the expansions of elliptic integrals for $\varepsilon \ll 1$

$$\begin{aligned} & \sqrt{1 - \varepsilon} \mathbf{\Pi} \left(m, 1 - \frac{\varepsilon}{1 + z_-} \right) \\ &= \frac{4 \log 2 - \sqrt{m} \log \left[\frac{1 + \sqrt{m}}{1 - \sqrt{m}} \right] - \log \frac{\varepsilon}{1 + z_-}}{2(1 - m)} + \mathcal{O}(\varepsilon), \\ & \sqrt{1 - \varepsilon} \left[\mathbf{\Pi} \left(n, 1 - \frac{\varepsilon}{1 + z_-} \right) - \mathbf{K} \left(1 - \frac{\varepsilon}{1 + z_-} \right) \right] \\ &= \frac{n \left(4 \log 2 - \frac{1}{\sqrt{n}} \log \left[\frac{1 + \sqrt{n}}{1 - \sqrt{n}} \right] - \log \frac{\varepsilon}{1 + z_-} \right)}{2(1 - n)} + \mathcal{O}(\varepsilon) \end{aligned}$$

and identifying $m = \frac{z_-(1 + \kappa^2)}{z_- + 1}$ and $n = \frac{z_-}{z_- + 1}$, we get

$$E = \frac{\sqrt{1 + \kappa^2}}{\pi \alpha' \kappa \sqrt{m}} \left(4 \log 2 - \sqrt{m} \log \left[\frac{1 + \sqrt{m}}{1 - \sqrt{m}} \right] - \log \frac{\varepsilon}{1 + z_-} \right), \tag{4.28}$$

$$S = \frac{\sqrt{1 + \kappa^2}}{\pi \alpha' \kappa} \sqrt{n} \left(4 \log 2 - \frac{1}{\sqrt{n}} \log \left[\frac{1 + \sqrt{n}}{1 - \sqrt{n}} \right] - \log \frac{\varepsilon}{1 + z_-} \right), \tag{4.29}$$

where we have used the relations

$$\begin{aligned} \frac{k}{\omega \sqrt{z_-(z_- + 1)(1 - m)}} &= \frac{1}{\sqrt{m}} \quad \text{and} \\ \frac{n}{\sqrt{z_-(z_- + 1)(1 - n)}} &= \sqrt{n}. \end{aligned} \tag{4.30}$$

Now the relation between energy and spin turns out to be

$$E - \frac{1}{\sqrt{mn}} S = \frac{\sqrt{1 + \kappa^2}}{\pi \alpha' \kappa} \left(\log \left[\frac{1 - \sqrt{m}}{1 + \sqrt{m}} \right] \times + \frac{1}{\sqrt{mn}} \log \left[\frac{1 + \sqrt{n}}{1 - \sqrt{n}} \right] \right). \tag{4.31}$$

finally substituting $\frac{1}{\sqrt{m}} = \omega_0$ and $\sqrt{n} = k_0$, we get

$$E - \frac{\omega_0}{k_0} S = \frac{\sqrt{1 + \kappa^2}}{\pi \alpha' \kappa} \times \left(\log \left[\frac{\omega_0 - 1}{\omega_0 + 1} \right] + \frac{\omega_0}{k_0} \log \left[\frac{1 + k_0}{1 - k_0} \right] \right). \tag{4.32}$$

The above relation agrees with result found in [32].

5 Spinning both in $(\text{AdS}_5)_\kappa$ and $(\text{S}^5)_\kappa$

In this section we consider a string which spins both in $(\text{AdS}_5)_\kappa$ and $(\text{S}^5)_\kappa$. The ansatz for such string can be written as

$$\begin{aligned} t &= t(\tau), \quad \rho = \rho(\sigma), \quad \psi = \frac{\pi}{2}, \quad \psi_2 = \omega \tau, \\ \theta &= \theta(\sigma), \quad \varphi = \nu \tau. \end{aligned} \tag{5.1}$$

Since the embeddings of AdS_5 part remains unchanged from the previous section, the equation of motion of ρ and so as its solution and periodicity condition are remain same. While the equation of motion θ and Virasoro constraint equation are given by

$$\begin{aligned} 2\theta'' \left(1 + \kappa^2 \cos^2 \theta \right) + \kappa^2 \theta'^2 \sin 2\theta \\ + \nu^2 \sin 2\theta (1 + \kappa^2) &= 0, \end{aligned} \tag{5.2}$$

$$\begin{aligned} -\frac{\cosh^2 \rho k^2}{1 - \kappa^2 \sinh^2 \rho} + \frac{\rho^2}{1 - \kappa^2 \sinh^2 \rho} + \omega^2 \sinh^2 \rho \\ + \frac{\theta'^2}{1 + \kappa^2 \cos^2 \theta} + \frac{\nu^2 \sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} &= 0. \end{aligned} \tag{5.3}$$

The solution to the equation motion of θ can be obtained in terms of elliptic function,

$$\sin \theta = \begin{cases} \mathbf{sn} \left(c_1 \sqrt{1 + \kappa^2} \sigma, \frac{1}{B^2} \right) & \text{for } B > 1 \\ B \mathbf{sn} \left(\sqrt{\nu^2 + c_1^2 \kappa^2} \sigma, B^2 \right) & \text{for } B < 1, \end{cases} \tag{5.4}$$

where $B = \sqrt{\frac{c_1^2(1 + \kappa^2)}{\nu^2 + c_1^2 \kappa^2}}$ and c_1 is an integration constant.

There is another periodicity condition comes from the string configuration along θ angle, which gives,

$$2\pi \sqrt{\nu^2 + \kappa^2 c_1^2} = 4\mathbf{K} \left[\frac{c_1^2(1 + \kappa^2)}{\nu^2 + \kappa^2 c_1^2} \right].$$

The general expressions of conserved charges E and S will be same as (4.17) and (4.18), while the extra angular momentum can be evaluated as,

$$\begin{aligned} \mathcal{J} &= \frac{\nu \sqrt{1 + \kappa^2}}{2\pi \alpha'} \int_0^{2\pi} d\sigma \frac{\sin^2 \theta}{1 + \kappa^2 \cos^2 \theta} \\ &= \frac{2\nu \sqrt{1 + \kappa^2}}{\alpha' \pi} \int_0^{\theta_{max}} \frac{d\theta}{\theta'} \frac{\sin^2 \theta}{1 + \kappa^2 \cos^2 \theta}. \end{aligned}$$

So $\mathcal{J} = \frac{2\nu \sqrt{1 + \kappa^2}}{\alpha' \pi \kappa^2 \sqrt{\nu^2 + c_1^2 \kappa^2}} \times \left[\mathbf{\Pi} \left(\frac{\kappa^2 B^2}{1 + \kappa^2}, B^2 \right) - \mathbf{K}(B^2) \right]. \tag{5.5}$

Now using the periodicity condition of θ and ρ we express all the parameters in terms of B and z_-

$$c_1 = \frac{2BK[B^2]}{\pi\sqrt{1+\kappa^2}}, \quad v = \frac{2}{\pi}\sqrt{\frac{1+\kappa^2(1-B^2)}{1+\kappa^2}}\mathbf{K}[B^2], \tag{5.6}$$

$$\omega = \frac{2}{\pi\sqrt{1+\kappa^2}}\mathbf{K}\left(\frac{(1+\kappa^2)z_-}{1+z_-}\right),$$

$$k^2 = \frac{4}{\pi^2}\left[\frac{z_-}{(1+z_-)}\mathbf{K}^2\left(\frac{(1+\kappa^2)z_-}{1+z_-}\right) + \frac{B^2}{(1+\kappa^2)}\mathbf{K}^2(B^2)\right]. \tag{5.7}$$

Now substituting these relation in conserves charges we get

$$E = \frac{2\sqrt{(1+\kappa^2)(1+z_-)}}{\alpha'\pi(1-z_- \kappa^2)} \frac{\sqrt{1+z_-}}{\mathbf{K}\left(\frac{z_-(1+\kappa^2)}{z_-+1}\right)}$$

$$\times \sqrt{\frac{B^2\mathbf{K}^2(B^2)}{1+\kappa^2} + \frac{z_- \mathbf{K}^2\left(\frac{z_-(1+\kappa^2)}{z_-+1}\right)}{1+z_-}} \mathbf{E}\left(\frac{z_-(1+\kappa^2)}{z_-+1}\right),$$

$$S = \frac{2\sqrt{1+\kappa^2}}{\pi\alpha'\sqrt{(z_-+1)}}$$

$$\times \left[\mathbf{\Pi}\left(\frac{z_-}{z_-+1}, \frac{z_-(1+\kappa^2)}{z_-+1}\right) - \mathbf{K}\left(\frac{z_-(1+\kappa^2)}{z_-+1}\right) \right],$$

$$J = \frac{2\sqrt{1+\kappa^2 - \kappa^2 B^2}}{\alpha'\pi\kappa^2} \left[\mathbf{\Pi}\left(\frac{\kappa^2 B^2}{1+\kappa^2}, B^2\right) - \mathbf{K}(B^2) \right]. \tag{5.8}$$

Using the short string limit $z_- \rightarrow 0$ and $B \ll 1$, the leading terms in E , S and \mathcal{J} turn out to be

$$E^2 \approx \frac{1}{\alpha'^2} \left(B^2 + (1+\kappa^2)z_- \right),$$

$$S \approx \frac{\sqrt{1+\kappa^2}}{2\alpha'} z_-, \quad \text{and} \quad \mathcal{J} \approx \frac{B^2}{2\alpha'\sqrt{1+\kappa^2}}. \tag{5.9}$$

Finally the relation between E , S and \mathcal{J} becomes

$$E^2 \approx 2 \frac{\sqrt{1+\kappa^2}}{\alpha'} (J + S). \tag{5.10}$$

For $\kappa \rightarrow 0$, the above result reduces to result found in [37]. This scaling relation is of the form of usual Regge-type spectrum. In the case of $\kappa \rightarrow 0$ the dual gauge theory operator has been identified in [37]. In the present case of nonzero κ , the proper identification is still missing even for the simplest class of folded and spinning string solutions.

6 Summary and conclusion

In this paper, we have studied a generic class of semiclassical string solitons in κ deformed $AdS_5 \times S^5$ background. We have restricted ourselves to the bosonic part of the string

action in conformal gauge. We have found the exact solutions for the string configuration which are pulsating in $(AdS_5)_\kappa$ and at the same time spinning in the S^5_κ and vice versa and computed the general expressions of conserved charges in terms Jacobi elliptic functions and elliptic integrals. We have obtained the scaling relation between the energy and angular momentum for such configurations in short and long string limits. As expected these relations reduce to their corresponding undeformed results when $\kappa \rightarrow 0$. We have further studied an example of string soliton which spins along both subspaces of $(AdS_5 \times S^5)_\kappa$ and write down the scaling relation among the charges in the leading orders.

There are some open questions which could be addressed later. First, it would be interesting to study the corresponding gauge theory operator, if any, dual to these string solitons. Recall that the novelty of the solutions presented in [18] was to find out a scaling relation between various charges through an additional constant, whose gauge theory interpretation was highly nontrivial. It was argued that suitably adjusting this nontrivial constant one was able to see some direct relationship with the underlying gauge theory. Ours is a generalization of those results and we have been able to successfully incorporate all nontrivial factors which have been introduced by the deformation parameter κ . These are a generic class of solutions for the string trajectories in the gravity side. We hope this will give further insight in the study of string solutions in κ -deformed $AdS_5 \times S^5$. The recent studies suggest that the underlying gauge theory might be some nonlocal theory which at best remains obscure. Therefore, studying classical rotating and pulsating string solutions are expected to give us hints along these directions. Secondly, as the deformed geometry is question is not well understood as a generic string background it would be interesting to see if we can take clue from the deformed supergravity equations proposed in [39] and look for possible ‘‘modified’’ intersecting D-branes whose near horizon geometry gives rise to these deformed geometries. Thirdly, finding the multi spin string solution in this deformed background is worth attempting, though in this case the string rotating with equal spins along the two subspace will not be possible because of loss of symmetry due to the deformation. Finally, one can do a one loop calculation to find out the leading order corrections to the conserved charges along the lines of [15]. We will report on some of these issues in near future.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

References

1. J.M. Maldacena, The Large N limit of superconformal field theories and supergravity. *Int. J. Theor. Phys.* **38**, 1113–1133 (1999). [arXiv:hep-th/9711200](#)
2. E. Witten, Anti-de Sitter space and holography. *Adv. Theor. Math. Phys.* **2**, 253–291 (1998). [arXiv:hep-th/9802150](#)
3. O. Aharony, S.S. Gubser, J.M. Maldacena, H. Ooguri, Y. Oz, Large N field theories, string theory and gravity. *Phys. Rept.* **323**, 183–386 (2000). [arXiv:hep-th/9905111](#)
4. I. Bena, J. Polchinski, R. Roiban, Hidden symmetries of the AdS(5) x S**5 superstring. *Phys. Rev. D* **69**, 046002 (2004). [arXiv:hep-th/0305116](#)
5. J.A. Minahan, K. Zarembo, The Bethe ansatz for N=4 superYang–Mills. *JHEP* **03**, 013 (2003). [arXiv:hep-th/0212208](#)
6. S.S. Gubser, I.R. Klebanov, A.M. Polyakov, A semiclassical limit of the gauge/string correspondence. *Nucl. Phys. B* **636**, 99–114 (2002). [arXiv:hep-th/0204051](#)
7. H.J. de Vega, I.L. Egusquiza, Planetoid string solutions in (3+1) axisymmetric space-times. *Phys. Rev. D* **54**, 7513–7519 (1996). [arXiv:hep-th/9607056](#)
8. M. Kruczenski, Spiky strings and single trace operators in gauge theories. *JHEP* **08**, 014 (2005). [arXiv:hep-th/0410226](#)
9. A. Tirziu, A.A. Tseytlin, Semiclassical rigid strings with two spins in AdS(5). *Phys. Rev. D* **81**, 026006 (2010). [arXiv:0911.2417](#)
10. S. Frolov, A.A. Tseytlin, Multispin string solutions in AdS(5) x S**5. *Nucl. Phys. B* **668**, 77–110 (2003). [arXiv:hep-th/0304255](#)
11. J.A. Minahan, Circular semiclassical string solutions on AdS(5) x S(5). *Nucl. Phys. B* **648**, 203–214 (2003). [arXiv:hep-th/0209047](#)
12. J. Engquist, J.A. Minahan, K. Zarembo, Yang–Mills duals for semiclassical strings on AdS(5) x S(5). *JHEP* **11**, 063 (2003). [arXiv:hep-th/0310188](#)
13. H. Dimov, R.C. Rashkov, Generalized pulsating strings. *JHEP* **05**, 068 (2004). [arXiv:hep-th/0404012](#)
14. M. Smedback, Pulsating strings on AdS(5) x S**5. *JHEP* **07**, 004 (2004). [arXiv:hep-th/0405102](#)
15. M. Beccaria, G.V. Dunne, G. Macorini, A. Tirziu, A.A. Tseytlin, Exact computation of one-loop correction to energy of pulsating strings in AdS₅ x S⁵. *J. Phys. A* **44**, 015404 (2011). [arXiv:1009.2318](#)
16. I.Y. Park, A. Tirziu, A.A. Tseytlin, Semiclassical circular strings in AdS(5) and 'long' gauge field strength operators. *Phys. Rev. D* **71**, 126008 (2005). [arXiv:hep-th/0505130](#)
17. A. Khan, A.L. Larsen, Improved stability for pulsating multi-spin string solitons. *Int. J. Mod. Phys. A* **21**, 133–150 (2006). [arXiv:hep-th/0502063](#)
18. A. Khan, A.L. Larsen, Spinning pulsating string solitons in AdS(5) x S**5. *Phys. Rev. D* **69**, 026001 (2004). [arXiv:hep-th/0310019](#)
19. C. Klimcik, Yang–Baxter sigma models and dS/AdS T duality. *JHEP* **12**, 051 (2002). [arXiv:hep-th/0210095](#)
20. C. Klimcik, On integrability of the Yang–Baxter sigma-model. *J. Math. Phys.* **50**, 043508 (2009). [arXiv:0802.3518](#)
21. C. Klimcik, Integrability of the bi-Yang–Baxter sigma-model. *Lett. Math. Phys.* **104**, 1095–1106 (2014). [arXiv:1402.2105](#)
22. F. Delduc, M. Magro, B. Vicedo, An integrable deformation of the AdS₅ x S⁵ superstring action. *Phys. Rev. Lett.* **112**, 051601 (2014). [arXiv:1309.5850](#)
23. F. Delduc, M. Magro, B. Vicedo, Derivation of the action and symmetries of the q-deformed AdS₅ x S⁵ superstring. *JHEP* **10**, 132 (2014). [arXiv:1406.6286](#)
24. B. Hoare, R. Roiban, A.A. Tseytlin, On deformations of AdS_n x Sⁿ supercosets. *JHEP* **06**, 002 (2014). [arXiv:1403.5517](#)
25. G. Arutyunov, R. Borsato, S. Frolov, S-matrix for strings on η-deformed AdS₅ x S⁵. *JHEP* **04**, 002 (2014). [arXiv:1312.3542](#)
26. M. Khouchen, J. Kluson, Giant magnon on deformed AdS(3)xS(3). *Phys. Rev. D* **90**, 066001 (2014). [arXiv:1405.5017](#)
27. A. Banerjee, K.L. Panigrahi, On the rotating and oscillating strings in (AdS₃ x S³)_κ. *JHEP* **09**, 048 (2014). [arXiv:1406.3642](#)
28. K.L. Panigrahi, P.M. Pradhan, M. Samal, Pulsating strings on (AdS₃S³). *JHEP* **03**, 010 (2015). [arXiv:1412.6936](#)
29. G. Arutyunov, M. de Leeuw, S.J. van Tongeren, The exact spectrum and mirror duality of the (AdS₅ x S⁵)_η superstring. *Theor. Math. Phys.* **182**, 23–51 (2015). [arXiv:1403.6104](#)
30. R. Hernandez, J.M. Nieto, Spinning strings in the η-deformed Neumann-Rosochatius system. *Phys. Rev. D* **96**, 086010 (2017). [arXiv:1707.08032](#)
31. A. Banerjee, K.L. Panigrahi, On circular strings in (AdS₃ x S³)_κ. *JHEP* **09**, 061 (2016). [arXiv:1607.04208](#)
32. T. Kameyama, K. Yoshida, A new coordinate system for q-deformed AdS₅ x S⁵ and classical string solutions. *J. Phys. A* **48**, 075401 (2015). [arXiv:1408.2189](#)
33. A. Banerjee, S. Bhattacharya, K.L. Panigrahi, Spiky strings in κ-deformed AdS. *JHEP* **06**, 057 (2015). [arXiv:1503.07447](#)
34. T. Kameyama, K. Yoshida, Minimal surfaces in q-deformed AdS₅³ with Poincare coordinates. *J. Phys. A* **48**, 245401 (2015). [arXiv:1410.5544](#)
35. T. Araujo, I. Bakhmatov, E. Colgin, J.-i. Sakamoto, M.M. Sheikh-Jabbari, K. Yoshida, Conformal twists, Yang–Baxter σ-models & Holographic Noncommutativity. [arxiv:1705.02063](#)
36. T. Araujo, I. Bakhmatov, E. Colgin, J. Sakamoto, M.M. Sheikh-Jabbari, K. Yoshida, Yang–Baxter σ-models, conformal twists, and noncommutative Yang–Mills theory. *Phys. Rev. D* **95**, 105006 (2017). [arXiv:1702.02861](#)
37. J.G. Russo, Anomalous dimensions in gauge theories from rotating strings in AdS(5) x S**5. *JHEP* **06**, 038 (2002). [arXiv:hep-th/0205244](#)
38. S. Frolov, A.A. Tseytlin, Semiclassical quantization of rotating superstring in AdS(5) x S**5. *JHEP* **06**, 007 (2002). [arXiv:hep-th/0204226](#)
39. G. Arutyunov, S. Frolov, B. Hoare, R. Roiban, A.A. Tseytlin, Scale invariance of the η-deformed AdS₅ x S⁵ superstring, T-duality and modified type II equations. *Nucl. Phys. B* **903**, 262–303 (2016). [arXiv:1511.05795](#)