

youngstock in a given period is slightly better correlated with coat type at the end of the period than with coat type at the beginning of a period. Differences between breed groups do not exist. Partial correlation of coat type and youngstock weight do not differ significantly from zero.

## VI. — Séance ouverte et boîte aux suggestions. II

### MAXIMAL UTILIZATION OF HETEROSIS IN MILK AND BEEF PRODUCTION

A. HORN, S. BOZO and A. DUNAY

*Dep. of Animal Husbandry University of Veterinary Science, Budapest 7, P.O.B. 2, Hungary*

Authors worked out a breeding scheme in order to utilize the yearly genetic progress in milk and beef production as well as "hybrid vigor" and "type heterosis". Latter term is proposed to be introduced into the terminology of animal production. This scheme which integrates milk and beef production, keeps space with the international genetic progress and attains a heterosis effect over 60 per cent in milk production and at the same time assures a heterosis effect (hybrid vigor as well as type heterosis) of 100 per cent in the single suckler cow stock as well as in the population for direct fattening. In this scheme pure breeding is practiced only in herds meant for producing sires of high genetic quality whereas in milk and beef production only crossbred populations are involved. The scheme is already in practice within a large scale experiment of 6 000 cows in criss-cross breeding by using the best genetic material of *USA-Canadian* and *Danish Jersey* breeds and of three types of beef breeds. In this conception breeds are no more the objects of production but the building bricks of the production systems.

### HITCH-HIKING EFFECT AND LINKAGE DISEQUILIBRIUM: THE EXAMPLE OF TWO CLOSELY LINKED LOCI IN THE PIG, HALOTHANE SENSITIVITY (*Hal*) AND PHOSPHOHEXOSE ISOMERASE (*PHI*)

G. GUÉRIN, L. OLLIVIER and P. SELLIER

*Département de Génétique animale, I.N.R.A., 78350 Jouy-en-Josas, France*

The gene for halothane sensitivity (*Hal*<sup>s</sup>) is favoured in the selection for muscle development which has occurred in some modern breeds of pig (e.g. *Pietrain*). Such a selection tends to increase the frequency of another gene (*PHI*<sup>B</sup>) at the closely linked locus for *PHI*, by virtue of a so-called "hitch-hiking effect", the *PHI* locus itself being selectively neutral. Another effect of selection at the *Hal* locus is to generate a linkage disequilibrium between the two loci, *Hal* and *PHI*, the evolution of this disequilibrium depending on the type of selection applied at the *Hal* locus. This point is demonstrated by the observation of two lines of pigs selected differently for 4 generations, and thus confirms a recent theoretical work (THOMSON, 1977) showing that a possible source of linkage disequilibrium may be the hitch-hiking effect of a selected locus on another closely linked neutral locus.

### GRENZEN DER MAST- UND SCHLACHTLEISTUNG BEIM SCHWEIN UND SELEKTION ENTLANG VON LEISTUNGSGRENZEN

F. ZIMMERMANN

*Av. des Peupliers, 1012 Pully (Schweiz)*

Anhand von integral leistungsgeprüften Herden werden Papierselektionen durchgeführt. Die daraus resultierenden Merkmalsmuster zeigen, dass Niveaugrenzen auftreten können, welche das Zusammenwirken der Teilleistungen in ungünstiger Weise verändern. In einer Arbeitshypothese wird gezeigt, wie die Merkmalsmuster von Tieren, deren Leistungen Niveaugrenzen