

RESEARCH ARTICLE

Open Access



A *Glycine max* sodium/hydrogen exchanger enhances salt tolerance through maintaining higher Na^+ efflux rate and K^+/Na^+ ratio in *Arabidopsis*

Tian-Jie Sun^{1†}, Long Fan^{1†}, Jun Yang¹, Ren-Zhi Cao², Chun-Yan Yang³, Jie Zhang^{1*} and Dong-Mei Wang^{1*} 

Abstract

Background: Soybean (*Glycine max* (L.)) is one the most important oil-yielding cash crops. However, the soybean production has been seriously restricted by salinization. It is therefore crucial to identify salt tolerance-related genes and reveal molecular mechanisms underlying salt tolerance in soybean crops. A better understanding of how plants resist salt stress provides insights in improving existing soybean varieties as well as cultivating novel salt tolerant varieties. In this study, the biological function of *GmNHX1*, a NHX-like gene, and the molecular basis underlying *GmNHX1*-mediated salt stress resistance have been revealed.

Results: We found that the transcription level of *GmNHX1* was up-regulated under salt stress condition in soybean, reaching its peak at 24 h after salt treatment. By employing the virus-induced gene silencing technique (VIGS), we also found that soybean plants became more susceptible to salt stress after silencing *GmNHX1* than wild-type and more silenced plants wilted than wild-type under salt treatment. Furthermore, *Arabidopsis thaliana* expressing *GmNHX1* grew taller and generated more rosette leaves under salt stress condition compared to wild-type. Exogenous expression of *GmNHX1* resulted in an increase of Na^+ transportation to leaves along with a reduction of Na^+ absorption in roots, and the consequent maintenance of a high K^+/Na^+ ratio under salt stress condition. *GmNHX1*-GFP-transformed onion bulb endothelium cells showed fluorescent pattern in which GFP fluorescence signals enriched in vacuolar membranes. Using the non-invasive micro-test technique (NMT), we found that the Na^+ efflux rate of both wild-type and transformed plants after salt treatment were significantly higher than that of before salt treatment. Additionally, the Na^+ efflux rate of transformed plants after salt treatment were significantly higher than that of wild-type. Meanwhile, the transcription levels of three osmotic stress-related genes, *SKOR*, *SOS1* and *AKT1* were all up-regulated in *GmNHX1*-expressing plants under salt stress condition.

Conclusion: Vacuolar membrane-localized *GmNHX1* enhances plant salt tolerance through maintaining a high K^+/Na^+ ratio along with inducing the expression of *SKOR*, *SOS1* and *AKT1*. Our findings provide molecular insights on the roles of *GmNHX1* and similar sodium/hydrogen exchangers in regulating salt tolerance.

Keywords: Soybean, *GmNHX1*, Salt stress, VIGS, K^+/Na^+ ratio

* Correspondence: zhangjiezhao1h@163.com; dongmeiwang63@126.com

[†]Tian-Jie Sun and Long Fan contributed equally to this work.

¹State Key Laboratory of North China Crop Improvement and Regulation, Hebei Key Laboratory of Plant Physiology and Molecular Pathology, Hebei Agricultural University, Baoding 071000, Hebei, China

Full list of author information is available at the end of the article



Background

Plants are subjected to various biotic as well as abiotic stresses during their growth. Due to increasingly exacerbated salinization worldwide [1], compounded with the abiotic stresses such as cold and drought, the damage caused by salt stress has been significantly worse. Plants reserve many ways to tolerate salt stress, including efflux of salt and intracellular partitioning [2]. Salt tolerance, similar to many other metabolic processes, requires the proper control of cellular pH [3]. Na^+/H^+ exchangers (NHXs) are integral membrane transporters that catalyze the electroneutral exchange of K^+ or Na^+ for H^+ and are implicated in cell expansion [4], development [5], ion homeostasis [6] and salt tolerance [7]. The *Arabidopsis* genome contains eight NHX homolog-encoding genes which are grouped based on their sequence similarity and localization into three distinct classes, those enriched in plasma membrane (NHX7/SOS1 and NHX8), endosomal/vesicular (NHX5, NHX6), and vacuolar membrane (NHX1, NHX2, NHX3, NHX4), respectively [2]. In *Arabidopsis*, NHXs that localize in vacuolar and plasma membrane are generally considered critical for maintaining Na^+/K^+ homeostasis [8]. NHX5 and NHX6, however, function as pH regulators of Golgi, trans-Golgi network, and pre-vacuolar compartments, regulating the sorting of newly synthesized peptides and the direction of Golgi-cargo movement [9].

Despite that *Glycine soja* (wild soybean) is often unsusceptible to salt stress [10], its close relative *Glycine max* (soybean) is typically osmotic sensitive [11]. Thus, the identification and characterization of endogenous genes that are involved in salt tolerance regulation would substantially benefit genetic breeding of soybeans. Overexpression of *GmNAC15*, a member of the NAC transcription factor family in soybean, enhances salt tolerance in soybean hairy roots [12]. Besides, overexpression of *GmSKI*, one of the multi-subunit E3 ligases, enhances tolerance to high salinity and drought stress when transformed into tobacco (*Nicotiana tabacum*) plants [13]. Overexpression of *GmBIN2*, a serine/threonine kinase related to brassinosteroid sensitivity, increases cellular Ca^{2+} content and reduces Na^+ content, together enhances salt tolerance in transgenic *Arabidopsis* plants [14]. Previous studies in our lab have demonstrated that the overexpression of *GmNHX1* (Gene ID: LOC100816746) is able to complement the defect of the *Saccharomyces cerevisiae* *ena1-4*, *nhx1* and *nha1* mutant and reduce the hindering effect of salt stress on cell growth [15]. In this study, we extend our understanding of the function of *GmNHX1* gene and its roles in regulating salt tolerance. We uncover its subcellular localization as well as how its transcription responds to salt stress in soybeans. We further use *Arabidopsis* ecotype Col-0, a model plant allele that has been used in

numerous physiology studies, as a host to evaluate the function and mechanism of *GmNHX1* under salt stress condition. We uncover the role of *GmNHX1* by utilizing this model organism, which is to maintain a higher Na^+ efflux rate and K^+/Na^+ ratio under salt stress. Our findings provide important implications for understanding the molecular basis underlying salt tolerance in plants.

Results

GmNHX1 gene is related to salt stress resistance in plants

In order to examine the relation between *GmNHX1* and the response to salt stress in soybeans, we examined the transcription level of *GmNHX1* in soybean variety “Jidou-7”. We observed an increase of *GmNHX1* gene expression after salt treatment and the peak of its expression was at 24 h post treatment (hpt), reaching approximately 10-fold of that of before treatment (0 hpt). The expression of *GmNHX1* then began to decrease to approximately 5-fold of that of 0 hpt at 48 hpt (Fig. 1a). The upregulation of *GmNHX1* expression under salt stress implies its relevance to salt stress response in soybeans and its potential role in regulating salt tolerance in soybeans.

We then picked a specific fragment in the *GmNHX1* coding region, to construct the TRV-VIGS vector, and the expression of *GmNHX1* was examined using RT-qPCR after gene silencing. The result showed a satisfactory silencing efficiency after a routine period of TRV-VIGS in soybean plants. Compared to EV (unsilenced plants), TRV vector carrying *GmNHX1* specific fragment reduced the expression level of *GmNHX1* in infected soybeans by nearly 50% (Fig. 1b). As a result of the silencing of *GmNHX1*, the top of the plant drooped and the leaves wilted after 200 mM NaCl solution treatment compared to EV plant (Fig. 1c), suggesting that *GmNHX1* is a critical gene that is involved in the process of plant adaptation to salt stress.

GmNHX1 enhances salt tolerance in *Arabidopsis*

Since the silence of *GmNHX1* reduces salt stress resistance in soybean, we wonder if exogenous expression of this gene could cause the opposite. We constructed the coding sequence of *GmNHX1* gene into a T-DNA within which the *GmNHX1* expression is driven by a *CaMV* 35S promoter, and the construct was then transformed into *Arabidopsis* Col-0. The expression of *GmNHX1* in transformed plants was verified by RT-PCR. We observed significant expression of the gene in transformed plants but undetectable level of the expression in untransformed col-0 (Fig. 2). In order to investigate whether *GmNHX1* overexpression is able to enhance salt tolerance, 21-day-old *Arabidopsis* plants expressing *GmNHX1* were irrigated with 170 mM NaCl solution for 20 days, and plants that were irrigated with water were set as control. Wilting and chlorosis phenotypes were observed in wild-type plants after salt stress treatment,




Fig. 1 *GmNHX1* positively regulates plant salt tolerance. Transcription Level of *GmNHX1* under salt stress in Soybean was detected by RT-qPCR, *ACTIN* gene was used as a reference gene. The data shows the mean \pm S.E. of triplicate experiments. Columns with different letters indicate significant differences at $P < 0.05$ (a). The silencing efficiency of *GmNHX1* in silenced plants using VIGS were detected by RT-qPCR, using *ACTIN* as reference gene (b). Phenotype of TRV- VIGS plants under 200 mM NaCl solution treatment for 24 h (c)

whereas only slight chlorosis was observed in all three lines that express *GmNHX1* exogenously (Fig. 3a, b). Parameters such as stem length, number of rosette leaves, fresh weight, and dry weight were also measured. Plants expressing *GmNHX1* showed significantly higher values in the measurements of stem length (Fig. 3c), rosette leaves (Fig. 3d), fresh weight (Fig. 3e) and dry weight (Fig. 3f), compared to wild-type under salt stress

treatment. Taken together, we concluded that *GmNHX1* enhanced plant resistance to salt stress condition.

***GmNHX1* enhances salt tolerance through maintaining K^+/Na^+ ratio in root**

After revealing the positive role of *GmNHX1* in salt resistance, we wondered the molecular mechanism underlying the *GmNHX1*-mediated salt resistance. *GmNHX1* has a typical NHX domain, which encodes a functional unit that pumps Na^+ against its concentration pressure in exchange for a proton. We therefore measured K^+ and Na^+ content in *GmNHX1* expressing plants before and after salt stress treatment. K^+ content in all three *GmNHX1* expressing lines were significantly higher in roots and leaves both before and after salt stress treatment (Fig. 4a, b). As for Na^+ content, all the three lines expressing *GmNHX1* had no significant difference compared to wild-type plants in leaves before salt stress treatment (Fig. 4c), however all the three lines showed higher Na^+ level in leaves after salt stress treatment than wild-type (Fig. 4c). In the meantime, Na^+ content was significantly lower after salt stress treatment in roots of two of the three lines which express *GmNHX1* (OE 1-1 and OE 1-4) compared to wild-type





Fig. 2 Expression analysis of *GmNHX1* in transformed *Arabidopsis*. Using *ACTIN* as reference gene, RT-PCR was performed to detect the expression of *GmNHX1*




plants (Fig. 4d). These results suggested that exogenous expression of *GmNHX1* might elevate Na^+ transportation to leaves and reduce Na^+ content in roots.

The maintenance of a high K^+/Na^+ ratio within plant cells is one of the key factors that mediate salt tolerance in plants [16], especially for the root tissue. Compared to K^+/Na^+ ratio in leaves (Fig. 4e), K^+/Na^+ ratio in roots is significantly higher in all three *GmNHX1* expressing

lines compared to wild-type after salt stress treatment (Fig. 4f). *GmNHX1* therefore contributes to the high K^+/Na^+ ratio in roots, in accordance with our results above.

Vacuolar membrane-localized *GmNHX1* regulates K^+ and Na^+ efflux

Plants overcome salt stress by means of regulating Na^+ efflux and the partitioning of Na^+ into vacuolar [2]. Given



the observations above, we wondered whether GmNHX1 regulated salt tolerance via adjusting the efflux of these two ion molecules, or modifying the cellular ion storage. To address this question, subcellular localization of GmNHX1 was examined. An eGFP was fused to the C-terminal of GmNHX1 and the resulted fusion protein was constitutively expressed under the control of a *CaMV 35S* promoter. Transformed onion bulb endothelium cells showed a pattern of GFP fluorescence in which signals were enriched in the vacuolar membranes (Fig. 5). To explore the absorption law of Na⁺ and K⁺ under salt stress in *Arabidopsis* alleles expressing *GmNHX1*, we used NMT to detect the flow changes of Na⁺ and K⁺ after 25 h of 100 mM

NaCl treatment. The result showed that the efflux of K⁺ in transformed plants were significantly fewer than that of wild-type, with no significant difference for Na⁺ efflux between those before and after salt treatment. The K⁺ efflux of both transformed and wild-type *Arabidopsis* after salt treatment were fewer than that of before salt treatment, but the rate of Na⁺ efflux of transformed plants was greater than that of wild-type. The rate of Na⁺ efflux of both transformed and wild-type *Arabidopsis* after salt treatment was significantly greater than that of before salt treatment, and the rate of Na⁺ efflux of transformed plants after salt treatment was significantly higher than that of wild-type (Fig. 6). This result suggested that *Arabidopsis* plants expressing




Fig. 5 Subcellular localization of GmNHX1. Onion bulb endothelium cells expressing eGFP (a-c) and *GmNHX1*-eGFP (d-f) was analyzed under fluorescent microscopy, images were acquired using the 488 nm excitation (a, d) and light (b, e). Superimposed images were generated in c and f, respectively. Arrows indicate the nucleus regions. Bar = 100 μ m

GmNHX1, a vacuolar membrane-localized protein, maintain K^+/Na^+ ratio via elevating Na^+ efflux rate in roots, along with reducing Na^+ accumulation, which thereby avoiding the toxic effects of excessive salt in cells.



GmNHX1 regulates stress responsive genes

The seemingly contradictory observations above, in which GmNHX1 localized in vacuolar membranes whereas it was able to mediate Na^+ efflux, prompted us to wonder if there were other mechanisms involved. The process of salt stress defense in plants relies on regulation of a series of stress responsive genes [17]. To reveal the relationship between GmNHX1 and stress responsive genes, we employed RT-qPCR to quantify the expression of *AKT1*, *HKT1*, *SOS1* and *SKOR*, which are closely related to salt stress response [2]. Before salt stress treatment, the expression of *SKOR* and *HKT1* in transformed plants were significantly lower than those of wild-type, whereas *SOS1*, *AKT1* showed no significant

difference between the two (Fig. 7). Salt stress treatment elevated the expression of *SOS1* and *AKT1*, and reduced the expression of *SKOR* and *HKT1*. After salt treatment, GmNHX1 upregulated the expression of *SKOR*, *SOS1* and *AKT1*, but not *HKT1*.

Discussion

Na^+/H^+ exchangers (NHX) function as regulators of intracellular ion homeostasis, mainly by increasing Na^+ excretion of cells (such as *SOS1*) [18] or compartmenting Na^+ into vacuolar, such as *AtNHX1*, to improve plant salt tolerance [19]. Ape et al. suggested that *AtNHX1* is critical to the resistance to salt stress in plants [20]. Since *GmNHX1* possesses classic sodium/hydrogen exchanger (NHX) features in its sequence, we therefore wonder if *GmNHX1* is also related to salt stress resistance. In this work, we use VIGS to investigate the function of GmNHX1 in soybeans under salt stress condition. VIGS is a fast, simple and reliable



approach that has been used in many functional biology studies, yet is still limited due to the difficulty in finding compatible plant virus stains [21]. TRV (*Tobacco rattle virus*)-mediated VIGS has been widely used in many functional biology studies [22], and is applicable in soybeans according to Liu et al. [23]. We found that the salt resistance has been reduced in *GmNHX1*-silenced plants, suggesting that this gene is closely related to plant salt stress resistance. Previous studies suggest that heterologous expression of chrysanthemum *DgNHX1* is able to improve salt tolerance in tobacco, causing an increase of Na^+ and K^+ accumulation in tobacco leaves [24]. *ZxNHX1* and *ZxVP1-1* could increase the salt and drought resistance of roots, as well as increase the accumulation of Na^+ , K^+ and Ca^{2+} in leaves [25]. Similarly, Heterologous expression of salt-tolerant plant hippocampus *SbNHX1* can improve salt tolerance of *Jatropha curcas*, increase Na^+ content and decrease K^+ content in leaves when exposed to 200 mM NaCl [26]. *Arachis hypogaea* plants expressing *AtNHX1* are resistant to drought, and the content of Na^+ and K^+ in leaves are increased [27]. Stress-inducible expression of *TaNHX2* significantly improves growth performance as well as Na^+ and K^+ content from the leaf and root tissue of T_2 transgenic eggplants (*Solanum melongena* L.) under salt stress, compared to non-transformed plants [28]. *NHX* in sweet sorghum is mainly involved in the transportation of Na^+ , facilitating Na^+ homeostasis in response to the increase of salt concentration [29]. Our work showed that the contents of Na^+ and K^+ in roots have been significantly increased and the K^+/Na^+ ratio also increases significantly in plants expressing *GmNHX1* under salt stress, in accordance with these reports.

Previous studies have shown that *SOS1*, membrane Na^+/H^+ exchanger protein, mediates the efflux of Na^+ in roots [30], previous research shown that the high efficiency K^+ channel protein *HKT1* is located in the membrane, which plays an important role in maintaining the homeostasis of K^+ and Na^+ on the aboveground part of the plant [31–33]. *SOS1* and *HKT1* are located in the membrane, which play key role in regulating K^+ absorbance from soil to the root cells, and the K^+ and Na^+ homeostasis [34]. *AKT1* encodes an internal rectifier K^+ channel protein, mainly regulates internal K^+ flow into the root cells [35]. *SKOR* as an external rectifier K^+ channel protein mainly involved in K^+ loading from the column cell to the xylem [36]. Yuan et al. [37] prove that the *ZxNHX1* regulates the whole plant K^+/Na^+ homeostasis, and the expression of ion transport protein genes such as *SKOR*, *SOS1*, *AKT1* and *HKT1* were significantly down-regulated. In the NMT result, the net flux rate of K^+ in transformed plants under normal conditions is significantly lower than that in the WT plants, consistent with the RT-qPCR results that *GmNHX1* down-regulated the

expression of *SKOR* and *HKT1* genes in *Arabidopsis* under normal conditions, but had little effect on *SOS1* and *AKT1*. Salt stress induced the expression of *SOS1* and *AKT1*, and decreased the expression of *SKOR* and *HKT1*. Under salt stress, compared with WT, *SKOR*, *SOS1* and *AKT1* in *GmNHX1* transformed plants increased significantly, but not *HKT1*, which may explain why efflux of Na^+ in the transformed plants after salt treatment is significantly greater than that in the WT plants.

We show that *GmNHX1* overexpression enhances salt tolerance of *Arabidopsis*. We speculate that, on one hand, by increasing the efflux of Na^+ in root cells, Na^+ content is consequently reduced in roots and the K^+/Na^+ ratio increases; on the other hand, the Na^+ is transported to leaves through the xylem, accumulating in the vacuolar in leaves, and K^+/Na^+ ratio is maintained stable. When under salt stress, the efflux rate of Na^+ in roots is greatly increased, and the possible reason is that the expression of intermembrane Na^+/H^+ exchanger *SOS1* is induced by *GmNHX1*. Our results further show that the salt tolerance of the transgenic *Arabidopsis* is mainly achieved by regulating the Na^+ distribution in plants.

Conclusions

In summary, soybean Na^+/H^+ exchanger *GmNHX1* responds to and regulates plant tolerance to salt stress. In transformed *Arabidopsis* which expresses *GmNHX1*, *GmNHX1* changes the flow rate of K^+ and Na^+ in root cells by altering the expression of *SKOR* and *SOS1*, in order to regulate the accumulation of K^+ and Na^+ in roots and leaves, as well as the maintenance of a high K^+/Na^+ ratio in roots, together improve the tolerance to salt stress in plants.

Methods

Cultivation and salt treatment of plant materials

Soybean cv. Jidou-7 was obtained from the Institute of Grain and Oil Crops, Hebei Academy of Agricultural and Forestry Sciences, and was cultivated in a greenhouse with a 14 h light/10 h dark cycle at a constant temperature of 25 °C and 700 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. The 10-day-old soybean seedlings were transferred to Hoagland nutrient solution for 24 h, then transferred to Hoagland nutrient solution containing 170 mM NaCl, and sampled at 0 h, 6 h, 12 h, 24 h and 48 h, before RT-qPCR analysis. *Arabidopsis* ecotype Col-0 was obtained from the Arabidopsis Biological Resource Center (ABRC; <http://abrc.osu.edu>). The wild-type and transformed *Arabidopsis* seeds were surface sterilized and planted on MS medium, then transferred to vermiculite for 20 days and watered in Hoagland nutrient solution. For salt treatment, 21-day-old *Arabidopsis* plants were irrigated with 170 mM NaCl solution, biomass measurement including plant height, rosette number, fresh weight and dry weight,

measurement of K^+ and Na^+ content and expression quantitation of salt stress related genes were performed 20 days after salt stress treatment initiation.

Virus induced gene silence

A specific fragment of *GmNHX1* was amplified using primer pair *GmNHX1*-F and *GmNHX1*-R (Table 1), with PrimeSTAR HS DNA Polymerase (TaKaRa). The virus vector that is used to silence *GmNHX1*, pTRV2-*GmNHX1*, was constructed by inserting the amplified fragment of *GmNHX1* into pTRV2 vector between *Bam*H I and *Kpn* I recognition sites. TRV-VIGS was performed according to the previous report [22]. After infection, soybean seedlings were treated with 170 mM NaCl solution for 24 h, then *GmNHX1* silencing efficiency was determined by RT-qPCR.

RT-PCR and RT-qPCR

Total RNA was isolated from plant material using UNIQ-10 Column Trizol Total RNA Isolation Kit (Sangon), and reverse transcript with PrimeScript RT reagent Kit with gDNA Eraser (TaKaRa). RT-PCR and RT-qPCR was performed using Ex Taq DNA polymerase (TaKaRa) and SYBR Premix DimerEraser (TaKaRa) according to user manual, respectively, using *ACTIN* as reference

Table 1 Synthetic DNA oligo used in this research

Oligo name	Sequence (5' - 3')	Application
GmNHX1-F	acgttgacacggggtcccccttc atgccatgggaca	Construction of TRV induced <i>GmNHX1</i> silencing vector.
GmNHX1-R	ctagctaggggtacctccaga ggaccaacatccaac	
RT GmNHX1 F	actgcaagcaatgcaatca	Detection of transcriptional level of <i>GmNHX1</i> and using RT-PCR.
RT GmNHX1 R	ggccattacgttcagttgggtg	
RT ACTIN F	atggctgatggtgaagacattc	
RT ACTIN R	tccatgctcaatagggtacttg	
OE GmNHX1 F	ggtaccatggttttgaatcagttc	Construction of binary vector pCAMBIA1300-GmNHX1.
OE GmNHX1 R	tctagatcaacgcattgatggcca	
GFP GmNHX1 F	tgccatgggacaaaatggttttgaatc	Construction of GFP fused vector pCAMBIA1300-GmNHX1-GFP.
GFP GmNHX1 R	cgccccgggacgccattgatgg	
qRT AtSKOR F	accgaaacaaactcggtaggaa	Detection of transcriptional level of salt stress related genes using RT-qPCR.
qRT AtSKOR R	ttagcacgtagatagacaggaatg	
qRT AtSOS1 F	gtgaagcaatcaagcggaaa	
qRT AtSOS1 R	tgcaagaagcgtagaaca	
qRT AtHKT1 F	gattgtcccacgaatgaga	
qRT AtHKT1 R	caaaaccaagaagcaaggaac	
qRT AtAKT1 F	aaaggtctcactcatcaacaacga	
qRT AtAKT1 R	tcggcaaaagagcgaataag	
qRT ACTIN F	gcaccgcagagagaaaatac	
qRT ACTIN R	caccaccgaaccagataaga	

gene. Primers used in RT-PCR and RT-qPCR experiments have been listed in Table 1.

Arabidopsis transformation

Full length CDS of *GmNHX1* was PCR amplified using primer pair OE GmNHX1 F/ R (Table 1), and constructed into binary vector pCAMBIA1300 between the restriction enzyme recognize site of *Kpn* I and *Xba* I, under control of *CaMV* 35S promoter.

In vivo measurement of K^+ and Na^+

Dried plant materials were ground into fine powder. Concentrated sulfuric acid was added to the ground powder, and the mixture was boiled at 170 °C for 20 min. A few drops of 30% hydrogen peroxide were added till a large amount of white smoke appeared, followed by digestion at 220 °C for 40 min, then 330 °C for 2 h. The content of Na^+ and K^+ ions were measured using a flame spectrophotometer (Sherwood M410).

Subcellular localization of GmNHX1

Full-length CDS of GmNHX1 was PCR amplified using primer pair GFP GmNHX1 F/ R (Table 1), and constructed into pCAMBIA1300-GFP, between the restriction enzyme recognize sites *Nco* I and *Sma* I, and was fused to the N⁺ terminal of GFP, resulting a fusion protein that is expressed under the control of *CaMV* 35S promoter. Purified pCAMBIA1300-*GmNHX1*-GFP plasmid was bombarded with a particle gun (BioRad PDS- 1000/He). Transformed onion bulb endothelium cells were cultivated in 1/2 MS medium for 24 h, before analyzed under the fluorescence microscopy (Olympus BX53).

Measurement of Na^+ and K^+ flow rate

Fluxes of Na^+ and K^+ ion was measured using NMT. 7-day-old *Arabidopsis* seedlings were transferred to MS medium containing 100 mM NaCl, NMT test was performed by Xuyue (Beijing) Sci.& tech. co., ltd., in accordance with previous report [38].

Abbreviations

hpt: Hours post treatment; NHX: Na^+/H^+ exchanger; NMT: Non-invasive micro-test technique; PCR: Polymerase chain reaction; TRV: *Tobacco rattle virus*; VIGS: Virus-induced gene silencing

Acknowledgements

We thank Clara Elizabeth of the Pacific Lutheran University, Tacoma, and Zhou-Liang YU of the University of California, Berkeley, for their help in polishing the language of this article.

Authors' contributions

TJS wrote this manuscript, TJS, LF and JY performed the experiment, RZC and CYY provided technical support, JZ and DMW conceived and designed the experiments. All authors have read and approved the final manuscript.

Funding

The sources of funding for this study are provided by China Ministry of Agriculture and Hebei Provincial Department of Science and Technology through the following projects:

(1) Major scientific and technological projects for breeding new varieties of genetically modified organisms (2009ZX08004-001B), Dong-Mei Wang; (2) Major scientific and technological projects for breeding new varieties of genetically modified organisms (2014ZX0800402B-001), Dong-Mei Wang; (3) Introducing Foreign Student Funding Program (CN201706), Jie Zhang. The funding bodies had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹State Key Laboratory of North China Crop Improvement and Regulation, Hebei Key Laboratory of Plant Physiology and Molecular Pathology, Hebei Agricultural University, Baoding 071000, Hebei, China. ²Department of Computer Science, Pacific Lutheran University, Tacoma, WA 98447, USA. ³Hebei Food and Oil Crops Institute, Shijiazhuang 050031, Hebei, China.

Received: 29 January 2019 Accepted: 17 October 2019

References

- Xu Y. Envirotyping for deciphering environmental impacts on crop plants. *Theor Appl Genet.* 2016;129:653–73 Springer.
- Deinlein U, Stephan AB, Horie T, Luo W, Xu G, Schroeder JJ. Plant salt-tolerance mechanisms. *Trends Plant Sci.* 2014;19:371–9.
- Chen X, Bao H, Guo J, Jia W, Tai F, Nie L, et al. Na⁺/H⁺ exchanger 1 participates in tobacco disease defence against *Phytophthora parasitica* var. *nicotianae* by affecting vacuolar pH and priming the antioxidative system. *J Exp Bot.* 2014;65:6107–22.
- Walker DJ, Leigh RA, Miller AJ. Potassium homeostasis in vacuolate plant cells (cytosolic K⁺/cytosolic pH/plant vacuole). *Plant Biol.* 1996;93:10510–4.
- Bassil E, Ohto M, Esumi T, Tajima H, Zhu Z, Cagnac O, et al. The *Arabidopsis* intracellular Na⁺/H⁺ antiporters NHX5 and NHX6 are endosome associated and necessary for plant growth and development. *Plant Cell.* 2011;23:224–39.
- Almeida DM, Margarida Oliveira M, Saibo NJM. Regulation of Na⁺ and K⁺ homeostasis in plants: towards improved salt stress tolerance in crop plants. *Genet Mol Biol.* 2017;40:326–45.
- Gao J, Sun J, Cao P, Ren L, Liu C, Chen S, et al. Variation in tissue Na⁺ content and the activity of *SOS1* genes among two species and two related genera of *Chrysanthemum*. *BMC Plant Biol.* 2016;16:1–15.
- Bassil E, Tajima H, Liang Y-C, Ohto M-A, Ushijima K, Nakano R, et al. The *Arabidopsis* Na⁺/H⁺ antiporters NHX1 and NHX2 control vacuolar pH and K⁺ homeostasis to regulate growth, flower development, and reproduction. *Plant Cell.* 2011;23:3482–97.
- Reguera M, Bassil E, Tajima H, Wimmer M, Chanoca A, Otegui MS, et al. pH regulation by NHX-type Antiporters is required for receptor-mediated protein trafficking to the vacuole in *Arabidopsis*. *Plant Cell.* 2015;27:1200–17.
- Ji W, Li Y, Li J, Dai CH, Wang X, Bai X, et al. Generation and analysis of expressed sequence tags from NaCl treated *Glycine soja*. *BMC Plant Biol.* 2006;6:4.
- Li M, Guo R, Jiao Y, Jin X, Zhang H, Shi L. Comparison of salt tolerance in *Soja* based on metabolomics of seedling roots. *Front Plant Sci.* 2017;8:1101.
- Ming LI, Zheng HU, Jiang QY, et al. *GmNAC15* overexpression in hairy roots enhances salt tolerance in soybean. *J Integr Agric.* 2018;17(3):530–8.
- Chen Y, Chi Y, Meng Q, et al. *GmSK1*, an *SKP1* homologue in soybean, is involved in the tolerance to salt and drought. *Plant Physiol Biochem.* 2018; 127:25–31.
- Ling-Shuang W, Qing-Shan C, Da-Wei X, et al. Overexpression of *GmBIN2*, a soybean glycogen synthase kinase 3 gene, enhances tolerance to salt and drought in transgenic *Arabidopsis* and soybean hairy roots. *Sci Agric Sin.* 2018;17(9):1959–71.
- Fan L, Sun T, Yang J, Zhang J, Wang D. Cloning and functional characterization of a vacuolar Na⁺/H⁺ antiporter gene (*GmNHX1*) from soybean. *J Agric Univ Hebei.* 2015;38:7–12,24.
- Li J, Bao S, Zhang Y, Ma X, Mishra-Knyrim M, Sun J, et al. *Paxillus involutus* strains MAJ and NAU mediate K⁽⁺⁾/Na⁽⁺⁾ homeostasis in *ectomycorrhizal Populus x canescens* under sodium chloride stress. *Plant Physiol.* 2012;159:1771–86.
- Villarino GH, Bombarely A, Giovannoni JJ, Scanlon MJ, Mattson NS. Transcriptomic analysis of *Petunia hybrida* in response to salt stress using high throughput RNA sequencing. *PLoS One.* 2014;9:1–13.
- Qiu Q-S, Guo Y, Dietrich MA, Schumaker KS, Zhu J-K. Regulation of *SOS1*, a plasma membrane Na⁺/H⁺ exchanger in *Arabidopsis thaliana*, by *SOS2* and *SOS3*. *Proc Natl Acad Sci U S A.* 2002;99:8436–41.
- Apse MP. Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ Antiport in *Arabidopsis*. *Science.* 1999;285:1256–8.
- Yamaguchi T, Apse MP, Shi H, Blumwald E. Topological analysis of a plant vacuolar Na⁺/H⁺ antiporter reveals a luminal C terminus that regulates antiporter cation selectivity. *Proc Natl Acad Sci U S A.* 2003;100:12510–5.
- Ramegowda V, Mysore KS, Senthil-Kumar M. Virus-induced gene silencing is a versatile tool for unraveling the functional relevance of multiple abiotic-stress-responsive genes in crop plants. *Front Plant Sci.* 2014;5:1–12.
- Senthil-Kumar M, Mysore KS. Tobacco rattle virus-based virus-induced gene silencing in *Nicotiana benthamiana*. *Nat Protoc.* 2014;9:1549–62.
- Liu X, Liu N, Li F, Wu L, Zhang J, Wang D. Establishment of TRV-mediated transient gene-silencing system in soybean. *Sci Agric Sin.* 2015;48:2479–86.
- Liu QL, Xu KD, Zhong M, Pan YZ, Jiang BB, Liu GL, et al. Cloning and characterization of a novel vacuolar Na⁺/H⁺ antiporter gene (*DgNHX1*) from chrysanthemum. *PLoS One.* 2013;8:1–7.
- Wu G-Q, Feng R-J, Wang S-M, Wang C-M, Bao A-K, Wei L, et al. Co-expression of xerophyte *Zygophyllum xanthoxylum* ZxNHX and ZxVP1–1 confers enhanced salinity tolerance in chimeric sugar beet (*Beta vulgaris* L.). *Front Plant Sci.* 2015;6:1–11.
- Jha B, Mishra A, Jha A, Joshi M. Developing transgenic *Jatropha* using the *SbNHX1* gene from an extreme halophyte for cultivation in saline wasteland. *PLoS One.* 2013;8(8):e71136.
- Banjara M, Zhu L, Shen G. Expression of an *Arabidopsis* sodium/ proton antiporter gene (*AtNHX1*) in peanut to improve salt tolerance. *Plant Biotechnol Rep.* 2012;6(1):59–67.
- Yarra R, Kirti PB. Expressing class I wheat NHX (*TaNHX2*) gene in eggplant (*Solanum melongena* L.) improves plant performance under saline condition. *Funct Integr Genomics.* 2019;19:541–54.
- Gu WT, Zhou LB, Liu RY, Jin WJ, Qu Y, et al. Synergistic responses of NHX, AKT1, and SOS1 in the control of Na⁺ homeostasis in sweet sorghum mutants induced by ¹²C⁶⁺-ion irradiation. *Nucl Sci Tech.* 2018;29:10.
- Yamaguchi T, Hamamoto S, Uozumi N. Sodium transport system in plant cells. *Front Plant Sci.* 2013;4:410.
- Davenport RJ, Muñoz-Mayor A, Jha D, Essah PA, Rus A, Tester M. The Na⁺ transporter AtHKT1;1 controls retrieval of Na⁺ from the xylem in *Arabidopsis*. *Plant Cell Environ.* 2007;30:497–507.
- Berthomieu P, Conéjéro G, Nublat A, Brackenburg WJ, Lambert C, Savio C, et al. Functional analysis of AtHKT1 in *Arabidopsis* shows that Na⁺ recirculation by the phloem is crucial for salt tolerance. *EMBO J.* 2003;22:2004–14.
- Mäser P, Eckelman B, Vaidyanathan R, Horie T, Fairbairn DJ, Kubo M, et al. Altered shoot/root Na⁺ distribution and bifurcating salt sensitivity in *Arabidopsis* by genetic disruption of the Na⁺ transporter AtHKT1. *FEBS Lett.* 2002;531:157–61.
- Assaha DVM, Ueda A, Saneoka H, Al-Yahyai R, Yaish MW. The role of Na⁺ and K⁺ transporters in salt stress adaptation in glycolphytes. *Front Physiol.* 2017;8:509.
- Xu J, Li HD, Chen LQ, Wang Y, Liu LL, He L, et al. A protein kinase, interacting with two Calcineurin B-like proteins, regulates K⁺ transporter AKT1 in *Arabidopsis*. *Cell.* 2006;125:1347–60.
- Liu K, Li L, Luan S. Intracellular K⁺ sensing of SKOR, a shaker-type K⁺ channel from *Arabidopsis*. *Plant J.* 2006;46:260–8.
- Yuan HJ, Ma Q, Wu GQ, Wang P, Hu J, Wang SM. ZxNHX controls Na⁺ and K⁺ homeostasis at the whole-plant level in *Zygophyllum xanthoxylum* through feedback regulation of the expression of genes involved in their transport. *Ann Bot.* 2015;115(3):495–507.
- Sun J, Chen S, Dai S, Wang R, Li N, Shen X, et al. NaCl-induced alternations of cellular and tissue ion fluxes in roots of salt-resistant and salt-sensitive poplar species. *Plant Physiol.* 2008;149:1141–53.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.