

RESEARCH

Open Access



A generalized Lyapunov-type inequality in the frame of conformable derivatives

Thabet Abdeljawad^{1*}, Jehad Alzabut¹ and Fahd Jarad²

*Correspondence: tabdeljawad@psu.edu.sa
¹Department of Mathematics and General Sciences, Prince Sultan University, P.O. Box 66833, Riyadh, 11586, Saudi Arabia
Full list of author information is available at the end of the article

Abstract

We prove a generalized Lyapunov-type inequality for a conformable boundary value problem (BVP) of order $\alpha \in (1, 2]$. Indeed, it is shown that if the boundary value problem

$$(\mathbf{T}_\alpha^c x)(t) + r(t)x(t) = 0, \quad t \in (c, d), x(c) = x(d) = 0$$

has a nontrivial solution, where r is a real-valued continuous function on $[c, d]$, then

$$\int_c^d |r(t)| dt > \frac{\alpha^\alpha}{(\alpha - 1)^{\alpha-1} (d - c)^{\alpha-1}}. \tag{1}$$

Moreover, a Lyapunov type inequality of the form

$$\int_c^d |r(t)| dt > \frac{3\alpha - 1}{(d - c)^{2\alpha-1}} \left(\frac{3\alpha - 1}{2\alpha - 1} \right)^{\frac{2\alpha-1}{\alpha}}, \quad \frac{1}{2} < \alpha \leq 1, \tag{2}$$

is obtained for a sequential conformable BVP. Some examples are given and an application to conformable Sturm-Liouville eigenvalue problem is analyzed.

MSC: 34A08; 26D15

Keywords: Lyapunov inequality; conformable derivative; Green's function; boundary value problem; Sturm-Liouville eigenvalue problem

1 Background

In [1], it was proved that if the boundary value problem (BVP)

$$x''(t) + r(t)x(t) = 0, \quad t \in (c, d), x(c) = x(d) = 0,$$

has a nontrivial solution, where r is a real-valued continuous function, then

$$\int_c^d |r(s)| ds > \frac{4}{d - c}. \tag{3}$$

Undoubtedly, the Lyapunov inequality (3) proves cooperative and supportive in differential equations. Indeed, it is frequently used to dominate certain quantities for the sake of

proving qualitative properties of solutions for differential equations. The Lyapunov inequality has been extensively under consideration for integer-order differential equations. As fractional differential equations become of high interest due to their demonstrated applications [2], nevertheless, the last few years have witnessed the appearance of many papers which systematically studied the fractional analog of this inequality and other types of inequalities; we refer the reader to [3–11] for Lyapunov type and to [12] for Gronwall type. The newly defined conformable fractional calculus was initiated in [13] and studied later on by the current author in [14, 15], where many properties of conformable operators were introduced. However, the progress in this direction is still at its earliest stage [16–18]. In this paper, we prove a Lyapunov-type inequality for a conformable fractional BVP. Moreover, a Lyapunov type inequality is obtained for a sequential conformable BVP. Some examples are given and an application to conformable Sturm-Liouville eigenvalue problem is analyzed. The new obtained inequalities generalize the existing ones.

2 Preliminaries on conformable derivatives

This section is devoted to the presentation of some preliminaries about higher order fractional conformable derivatives developed in [14].

Definition 1 ([13, 14]) The (left) conformable fractional derivative starting from c of a function $f : [c, \infty) \rightarrow \mathbb{R}$ of order $0 < \alpha \leq 1$ is defined by

$$(T_\alpha^c g)(t) = \lim_{\epsilon \rightarrow 0} \frac{g(t + \epsilon(t - c)^{1-\alpha}) - g(t)}{\epsilon}. \tag{4}$$

In the case when $c = 0$, we write T_α . If $(T_\alpha^c g)(t)$ exists on (c, d) then $(T_\alpha^c g)(c) = \lim_{t \rightarrow c^+} (T_\alpha^c g)(t)$.

Note that if g is differentiable then

$$(T_\alpha^c g)(t) = (t - c)^{1-\alpha} g'(t). \tag{5}$$

Moreover, the conformable fractional integral of order $0 < \alpha \leq 1$ starting at $c \geq 0$ is defined by $(I_\alpha^c g)(t) = \int_c^t g(x)(x - c)^{\alpha-1} dx$ and following [13] by $(I_\alpha^c g)(t) = \int_c^t g(x)x^{\alpha-1} dx$.

In the case of higher order, the following definition becomes true.

Definition 2 ([14]) Let $n < \alpha \leq n + 1$ and set $\gamma = \alpha - n$. Then the conformable fractional derivative starting from c of a function $g : [c, \infty) \rightarrow \mathbb{R}$ of order α , where $g^{(n)}(t)$ exists, is defined by

$$(T_\alpha^c g)(t) = (T_\gamma^c g^{(n)})(t). \tag{6}$$

In the case $c = 0$, we write T_α .

Note that if $\alpha = n + 1$ then $\gamma = 1$ and the fractional derivative of g becomes $g^{(n+1)}(t)$. Also when $n = 0$ (or $\alpha \in (0, 1)$) then $\gamma = \alpha$ and the definition coincides with that in Definition 1. From (6), it is immediate that if $n < \alpha \leq n + 1$ then $\gamma = \alpha - n$ and if moreover, the $(n + 1)$ th derivative (or the derivative of $g^{(n)}$) exists then we have

$$(T_\alpha^c g)(t) = (T_\gamma^c g^{(n)})(t) = (t - c)^{1-\gamma} g^{(n+1)}(t) = (t - c)^{1-\alpha+n} g^{(n+1)}(t). \tag{7}$$

Lemma 1 ([13]) *Assume that $g : [c, \infty) \rightarrow \mathbb{R}$ is continuous and $0 < \alpha \leq 1$. Then, for all $t > c$, we have*

$$T_\alpha^c I_\alpha^c g(t) = g(t).$$

In the case of higher order, the following definition is valid.

Definition 3 ([14]) *Let $\alpha \in (n, n + 1]$ and set $\gamma = \alpha - n$. Then the left conformable fractional integral starting at c of order α is defined by*

$$(I_\alpha^c g)(t) = \mathbf{I}_{n+1}^c((t - c)^{\gamma-1} g) = \frac{1}{n!} \int_c^t (t - x)^n (x - c)^{\gamma-1} g(x) dx. \tag{8}$$

Notice that if $\alpha = n + 1$ then $\gamma = 1$ and hence $(I_\alpha^c g)(t) = (\mathbf{I}_{n+1}^c g)(t) = \frac{1}{n!} \int_c^t (t - x)^n g(x) dx$, which is the iterative integral of g , $n + 1$ times over $(c, t]$.

Recall that the left Riemann-Liouville fractional integral of order $\alpha > 0$ starting from c is defined by

$$({}_c \mathbf{I}^\alpha g)(t) = \frac{1}{\Gamma(\alpha)} \int_c^t (t - s)^{\alpha-1} g(s) ds. \tag{9}$$

We see that $(I_\alpha^c g)(t) = ({}_c \mathbf{I}^\alpha g)(t)$ for $\alpha = n + 1, n = 0, 1, 2, \dots$

Example 1 In virtue of [14], we recall that $({}_c \mathbf{I}^\alpha (t - c)^{\mu-1})(x) = \frac{\Gamma(\mu)}{\Gamma(\mu + \alpha)} (x - c)^{\alpha + \mu - 1}, \alpha, \mu > 0$. Indeed, if $\mu \in \mathbb{R}$ such that $\alpha + \mu - n > 0$ then the conformable fractional integral of $(t - c)^\mu$ of order $\alpha \in (n, n + 1]$ is

$$(I_\alpha^c (t - c)^\mu)(x) = (\mathbf{I}_{n+1}^c (t - c)^{\mu + \alpha - n - 1})(x) = \frac{\Gamma(\alpha + \mu - n)}{\Gamma(\alpha + \mu + 1)} (x - c)^{\alpha + \mu}. \tag{10}$$

The following is a generalization of Lemma 1.

Lemma 2 ([14]) *Assume that $f : [c, \infty) \rightarrow \mathbb{R}$ such that $g^{(n)}(t)$ is continuous and $n < \alpha \leq n + 1$. Then, for all $t > c$, we have*

$$T_\alpha^c I_\alpha^c g(t) = g(t).$$

Theorem 1 ([14]) *Let $\alpha \in (n, n + 1]$ and $g : [c, \infty) \rightarrow \mathbb{R}$ be $(n + 1)$ times differentiable for $t > c$. Then, for all $t > c$, we have*

$$I_\alpha^c T_\alpha^c (g)(t) = g(t) - \sum_{k=0}^n \frac{g^{(k)}(c)(t - c)^k}{k!}. \tag{11}$$

Example 2 From [14], we recall that the solution of the following conformable fractional initial value problem:

$$(T_\alpha^c x)(t) = \lambda x(t), \quad 0 < \alpha \leq 1, x(c) = x_0, t > c, \tag{12}$$

is $x(t) = x_0 e^{\lambda \frac{(t-c)^\alpha}{\alpha}}$.

The following example shows why it is useful to work in conformable differential systems.

Example 3 Consider the following second order ordinary differential equation:

$$t^{2-2\alpha}y'' + t^{1-\alpha}((1-\alpha)t^{-\alpha} - 2)y' - 3y = 0, \quad 0 < \alpha \leq 1, \tag{13}$$

where $y(t)$ is assumed to have second order continuous derivative. In view of (13), one can figure out that it is not easy to solve it using standard methods. However, equation (13) is rewritten in the form of the sequential (local) conformable differential equation as

$$y^{(2\alpha)} - 2y^{(\alpha)} - 3y = 0, \tag{14}$$

where $y^{(2\alpha)} = t^{1-\alpha}(t^{1-\alpha}y')' = t^{1-\alpha}[t^{1-\alpha}y'' + (1-\alpha)t^{-\alpha}y']$ such that $y^{(2\alpha)}$ stands for $(T_\alpha^0 T_\alpha^0 y)(t)$ and T_α^0 is defined in (5). By using the operator method, we may write (14) in the form

$$(D^\alpha - 3)(D^\alpha + 1)y = 0.$$

The solution of the above equation has the form

$$y(t) = c_1 e^{3\frac{t^\alpha}{\alpha}} + c_2 e^{-\frac{t^\alpha}{\alpha}}.$$

For $\alpha = \frac{1}{2}$, (13) reduces to $ty'' + (\frac{1}{2} - 2\sqrt{t})y' - 3y = 0$, which by the above arguments has a solution $y(t) = c_1 e^{6\sqrt{t}} + c_2 e^{-2\sqrt{t}}$. It is worth mentioning that, for $\alpha = 1$, the solution $y(t) = c_1 e^{3t} + c_2 e^{-t}$ of the ordinary differential equation $y'' - 2y' - 3y = 0$ is verified.

3 A Lyapunov-type inequality for a conformable BVP

Consider the following (local) conformable BVP:

$$(T_\alpha^c x)(t) + r(t)x(t) = 0, \quad 1 < \alpha \leq 2, c < t < d, x(c) = x(d) = 0. \tag{15}$$

Lemma 3 $x(t)$ is a solution of the BVP (15) if and only if it satisfies the integral equation

$$x(t) = \int_c^d H(t,s)r(s)x(s) ds, \tag{16}$$

where H is the Green function for (15) defined by

$$H(t,s) = \begin{cases} \frac{(t-c)(d-s)}{d-c} \cdot (s-c)^{\alpha-2}, & c \leq t \leq s \leq d, \\ (\frac{(t-c)(d-s)}{d-c} - (t-s)) \cdot (s-c)^{\alpha-2}, & c \leq s \leq t \leq d. \end{cases}$$

Proof Applying the integral I_α^c to (15) and making use of Definition 3 and Theorem 1 with $n = 1$ and $\gamma = \alpha - 1$, we obtain

$$x(t) = c_1 + c_2(t-c) - \int_c^t (t-s)r(s)x(s)(s-c)^{\alpha-2} ds.$$

The condition $x(c) = 0$ implies that $c_1 = 0$ and the condition $x(d) = 0$ implies that $c_2 = \frac{1}{d-c} \int_c^d (d-s)r(s)x(s)(s-c)^{\alpha-2} ds$ and hence

$$x(t) = \frac{t-c}{d-c} \int_c^d (d-s)r(s)x(s)(s-c)^{\alpha-2} ds - \int_c^t (t-s)r(s)x(s)(s-c)^{\alpha-2} ds.$$

Then, using

$$\begin{aligned} & \int_c^d (d-s)r(s)x(s)(s-c)^{\alpha-2} ds \\ &= \int_c^t (d-s)r(s)x(s)(s-c)^{\alpha-2} ds + \int_t^d (d-s)r(s)x(s)(s-c)^{\alpha-2} ds, \end{aligned}$$

the proof is concluded. □

Lemma 4 *The Green function H defined above has the following properties:*

- (i) $H(t, s) \geq 0$ for all $c \leq t, s \leq d$.
- (ii) $\max_{t \in [c, d]} H(t, s) = H(s, s)$ for $s \in [c, d]$.
- (iii) $H(s, s)$ has a unique maximum, given by

$$\max_{s \in [c, d]} H(s, s) = H\left(\frac{c + (\alpha - 1)d}{\alpha}, \frac{c + (\alpha - 1)d}{\alpha}\right) = \frac{(d - c)^{\alpha-1}(\alpha - 1)^{\alpha-1}}{\alpha^\alpha}.$$

Proof Define the functions $h_1(t, s) = \frac{(t-c)(d-s)}{d-c} \cdot (s - c)^{\alpha-2}$ and $h_2(t, s) = (\frac{(t-c)(d-s)}{d-c} - (t - s)) \cdot (s - c)^{\alpha-2}$.

- (i) It is clear that $h_1 \geq 0$. To determine the sign of h_2 , we observe that $(t - s) = \frac{t-c}{d-c}(d - (c + \frac{(s-c)(d-c)}{(t-c)}))$ and that $c + \frac{(s-c)(d-c)}{(t-c)} \geq s$ if and only if $s \geq c$. Together with $(s - c)^{\alpha-2} \geq 0$ we conclude that $h_2 \geq 0$ as well. Hence, the proof of the first part is complete.
- (ii) It is clear that $h_1(t, s)$ is an increasing function in t . Differentiating h_2 with respect to t for every fixed s and following similar analysis as in first part we conclude that h_2 is a decreasing function.
- (iii) Let $g(s) = H(s, s) = \frac{(s-c)^{\alpha-1}(d-s)}{d-c}$. Then it is sufficient to show that $g'(s) = 0$ if $s = \frac{c+(\alpha-1)d}{\alpha}$ and hence the proof is completed. □

Theorem 2 *If the BVP (15) has a nontrivial solution, where r is a real-valued continuous function on $[c, d]$, then*

$$\int_c^d |r(t)| dt > \frac{\alpha^\alpha}{(\alpha - 1)^{\alpha-1}(d - c)^{\alpha-1}}. \tag{17}$$

Proof Let $x \in Y = C[c, d]$ be a nontrivial solution of the BVP (15), where $\|x\| = \sup_{t \in [c, d]} |x(t)|$. By Lemma 3, x must satisfy

$$x(t) = \int_c^d H(t, s)r(s)x(s) ds.$$

Therefore

$$\|x\| \leq \max_{t \in [c, d]} \int_c^d |H(t, s)r(s)| ds \|x\|$$

or

$$\max_{t \in [c, d]} \int_c^d |H(t, s)r(s)| ds \geq 1.$$

By using the properties of the Green function H proved in Lemma 4, we end up with

$$\frac{(d - c)^{\alpha-1}(\alpha - 1)^{\alpha-1}}{\alpha^\alpha} \int_c^d |r(s)| ds > 1. \tag{18}$$

Inequality (17) is an immediate conclusion of (18). □

Remark 1 If $\alpha = 2$, then (17) reduces to the classical Lyapunov inequality (3). We also invite the reader to compare the obtained generalized Lyapunov inequality in Theorem 2 and the one obtained recently and independently in [19]. The approach in [19] is different and the authors there proved the existence of solution in the space $AC^2[c, d] = \{u \in C^1[c, d] : u' \in AC[c, d]\}$. Further, we see that our obtained inequality provides, for example when applied to the Sturm-Liouville eigenvalue problem, sharper lower estimate for the eigenvalues. Indeed, in Section 5 we can see that the lower estimate $\frac{\alpha^\alpha}{(\alpha-1)^{\alpha-1}}$ is bigger than $4(\alpha - 1)$ for $1 < \alpha \leq 2$. This is due to the observation that $(\frac{\alpha}{\alpha-1})^\alpha \geq 4$, for $1 < \alpha \leq 2$. Further, for convenience, in the next section we prove a sequential type Lyapunov inequality version as well.

4 A Lyapunov-type inequality for a sequential conformable BVP

Consider the following sequential conformable BVP:

$$x^{(2\alpha)}(t) + r(t)x(t) = 0, \quad c < t < d, \frac{1}{2} < \alpha \leq 1, x(c) = x(d) = 0. \tag{19}$$

We shall prove a Lyapunov inequality for (19).

Lemma 5 $x(t)$ is a solution of the BVP (19) if and only if it satisfies the integral equation

$$x(t) = \int_c^d G(t, s)r(s)x(s) ds, \tag{20}$$

where G is the Green function of (19) defined by

$$G(t, s) = \begin{cases} g_1(t, s), & c \leq s \leq t \leq d, \\ g_2(t, s), & c \leq t \leq s \leq d, \end{cases}$$

such that

$$g_1(t, s) = (s - c)^{\alpha-1} \left[\frac{(s - c)^\alpha}{\alpha} - \frac{(t - c)^\alpha (s - c)^\alpha}{\alpha (d - c)^\alpha} \right]$$

and

$$g_2(t, s) = (s - c)^{\alpha-1} \left[\frac{(t - c)^\alpha}{\alpha} - \frac{(t - c)^\alpha (s - c)^\alpha}{\alpha (d - c)^\alpha} \right].$$

Proof Applying the integral I_α^c twice to (19) and using Definition 3, we get

$$y(t) = c_1 + c_2 \frac{(t - c)^\alpha}{\alpha} - \int_c^d r(s)x(s)(s - c)^{\alpha-1} \left[\frac{(t - c)^\alpha}{\alpha} - \frac{(s - c)^\alpha}{\alpha} \right] ds.$$

The condition $x(c) = 0$ implies that $c_1 = 0$ and the condition $x(d) = 0$ implies that $c_2 = \frac{\alpha}{(d - c)^\alpha} \int_c^d r(s)x(s)(s - c)^{\alpha-1} \left[\frac{(d - c)^\alpha}{\alpha} - \frac{(s - c)^\alpha}{\alpha} \right] ds$ and hence

$$x(t) = \frac{(t - c)^\alpha}{(d - c)^\alpha} \int_c^d r(s)x(s)(s - c)^{\alpha-1} \left[\frac{(d - c)^\alpha}{\alpha} - \frac{(s - c)^\alpha}{\alpha} \right] ds \tag{21}$$

$$- \int_c^t r(s)x(s)(s - c)^{\alpha-1} \left[\frac{(t - c)^\alpha}{\alpha} - \frac{(s - c)^\alpha}{\alpha} \right] ds. \tag{22}$$

Finally, the proof is concluded by splitting the first integral in (21) to

$$\begin{aligned} & \int_c^t r(s)x(s)(s - c)^{\alpha-1} \left[\frac{(d - c)^\alpha}{\alpha} - \frac{(s - c)^\alpha}{\alpha} \right] ds \\ & + \int_t^d r(s)x(s)(s - c)^{\alpha-1} \left[\frac{(d - c)^\alpha}{\alpha} - \frac{(s - c)^\alpha}{\alpha} \right] ds. \end{aligned} \quad \square$$

Lemma 6 *The Green function G defined above has the following properties:*

- (i) $G(t, s) \geq 0$ for all $c \leq t, s \leq d$.
- (ii) $\max_{t \in [c, d]} G(t, s) = G(s, s)$ for $s \in [c, d]$.
- (iii) $f(s) = G(s, s)$ has a unique maximum, given by

$$\begin{aligned} \max_{s \in [c, d]} G(s, s) &= G(\Lambda(c, d, \alpha), \Lambda(c, d, \alpha)) \\ &= \frac{(d - c)^{2\alpha-1}}{3\alpha - 1} \left(\frac{2\alpha - 1}{3\alpha - 1} \right)^{\frac{2\alpha-1}{\alpha}}, \end{aligned}$$

where

$$\Lambda(c, d, \alpha) = c + (d - c) \left[\frac{2\alpha - 1}{3\alpha - 1} \right]^{\frac{1}{\alpha}}.$$

Proof Define the two functions g_1 and g_2 as in Lemma 5.

- (i) The proof follows by noting that the function $g_1 \geq 0$ since $g_1(t, s)$ is decreasing in t for any s and $g_1(d, s) = 0$ for any s . Also, $g_2 \geq 0$ since $g_2(t, s)$ is increasing in t for any s and $g_2(c, s) = 0$ for any s .
- (ii) The proof of this part follows by noting that the function $g_1(t, s)$ is decreasing in t for any s and that $g_2(t, s)$ is increasing in t for any s by realizing that $(1 - \frac{(s - c)^\alpha}{(d - c)^\alpha}) \geq 0$ for all s .

(iii) Let $f(s) = G(s, s) = (s - c)^{\alpha-1} [\frac{(s-c)^\alpha}{\alpha} - \frac{(t-c)^\alpha (s-c)^\alpha}{\alpha(d-c)^\alpha}]$. Then one can show that $f'(s) = 0$ if $s = \Lambda(c, d, \alpha)$ and hence the proof is concluded. \square

Theorem 3 *If the BVP (19) has a nontrivial solution, where r is a real-valued continuous function on $[c, d]$, then*

$$\int_c^d |r(t)| dt > \frac{1}{G(\Lambda(c, d, \alpha), \Lambda(c, d, \alpha))} = \frac{3\alpha - 1}{(d - c)^{2\alpha-1}} \left(\frac{3\alpha - 1}{2\alpha - 1}\right)^{\frac{2\alpha-1}{\alpha}}. \tag{23}$$

Proof Let $x \in X = C[c, d]$ be a nontrivial solution of the BVP (19) where $\|x\| = \sup_{t \in [c, d]} |x(t)|$. By Lemma 5, x must satisfy

$$y(t) = \int_c^d G(t, s)r(s)x(s) ds.$$

Taking the norm leads to

$$\|x\| \leq \max_{t \in [c, d]} \int_c^d |G(t, s)r(s)| ds \|x\|$$

or equivalently

$$\max_{t \in [c, d]} \int_c^d |G(t, s)r(s)| ds \geq 1.$$

By using the properties of the Green function $G(t, s)$ given in Lemma 6, we come to the conclusion that

$$G(\Lambda(c, d, \alpha), \Lambda(c, d, \alpha)) \int_c^d |r(s)| ds > 1,$$

from which (23) follows. \square

Remark 2 Since $G(\Lambda(c, d, \alpha), \Lambda(c, d, \alpha))$ tends to $\frac{d-c}{4}$ as $\alpha \rightarrow 1$ then the classical Lyapunov inequality (3) is obtained again: $\alpha \rightarrow 1$. In this case, one may also deduce that $x^{(2\alpha)}(t) \rightarrow x''(t)$ as $\alpha \rightarrow 1$.

5 Application

Consider the following conformable Sturm-Liouville eigenvalue problem of order $\alpha \in (1, 2]$.

$$(\mathbf{T}_0^\alpha x)(t) + \lambda x(t) = 0, \quad 0 < t < 1, x(0) = x(1) = 0. \tag{24}$$

If λ is an eigenvalue of (24), then by Theorem 2, we must have $|\lambda| > \frac{\alpha^\alpha}{(\alpha-1)^{\alpha-1}}$. If x is twice differentiable, then by means of (5), equation (24) is equivalent to

$$t^{2-\alpha} x''(t) + \lambda x(t) = 0, \quad t \in (0, 1), x(0) = x(1) = 0, \tag{25}$$

which can be considered as a type of generalized Sturm-Liouville eigenvalue problem. It is not easy to find the eigenvalues and eigenfunctions of this problem. However, if we consider the sequential conformable Sturm-Liouville eigenvalue problem

$$x^{(2\alpha)}(t) + \lambda^2 x(t) = 0, \quad 0 < t < 1, x(0) = x(1) = 0, \quad \frac{1}{2} < \alpha \leq 1,$$

then by using the operator method its solution is given by

$$x(t) = c_1 \cos \lambda \frac{t^\alpha}{\alpha} + c_2 \sin \lambda \frac{t^\alpha}{\alpha}.$$

The boundary conditions imply that the eigenvalues are $\lambda = n\alpha\pi$, $n \in \mathbb{Z}$ and the correspondent eigenfunctions are $\sin \lambda \frac{t^\alpha}{\alpha}$. Notice that, for any $n \in \mathbb{Z}$, we have

$$\lambda^2 = n^2 \alpha^2 \pi^2 > G(\Lambda(0, 1, \alpha), \Lambda(0, 1, \alpha))^{-1} = (3\alpha - 1) \left(\frac{3\alpha - 1}{2\alpha - 1} \right)^{\frac{2\alpha - 1}{\alpha}}.$$

This is a direct verification of Theorem 3.

Acknowledgements

The first and the second author would like to thank Prince Sultan University for funding this work through research group Nonlinear Analysis Methods in Applied Mathematics (NAMAM) group number RG-DES-2017-01-17.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors read and approved the final manuscript.

Author details

¹Department of Mathematics and General Sciences, Prince Sultan University, P.O. Box 66833, Riyadh, 11586, Saudi Arabia.

²Department of Mathematics and Computer Science, Çankaya University, Ankara, 06790, Turkey.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 15 June 2017 Accepted: 1 October 2017 Published online: 11 October 2017

References

- Lyapunov, AM: Problème général de la stabilité du mouvement. *Ann. Fac. Sci. Univ. Toulouse* (2) **9**, 203-469 (1907). Reprinted in *Ann. Math. Stud.*, vol. 17, Princeton University Press, Princeton (1947)
- Podlubny, I: *Fractional Differential Equations*. Academic Press, San Diego (1999)
- Abdeljawad, T: A Lyapunov type inequality for fractional operators with nonsingular Mittag-Leffler kernel. *J. Inequal. Appl.* **2017**, Article ID 130 (2017). doi:10.1186/s13660-017-1400-5
- Agarwal, RP, Özbekler, A: Lyapunov inequalities for even order differential equations with mixed nonlinearities. *J. Inequal. Appl.* **2015**, Article ID 142 (2015)
- Hashizume, M: Minimization problem related to a Lyapunov inequality. *J. Math. Anal. Appl.* **432**(1), 517-530 (2015)
- Jleli, M, Ragoub, L, Samet, B: A Lyapunov-type inequality for a fractional differential equation under a Robin boundary condition. *J. Funct. Spaces* **2015**, Article ID 468536 (2015)
- Jleli, M, Samet, B: Lyapunov-type inequalities for a fractional differential equation with mixed boundary conditions. *Math. Inequal. Appl.* **18**(2), 443-451 (2015)
- Jleli, M, Samet, B: Lyapunov-type inequalities for fractional boundary value problems. *Electron. J. Differ. Equ.* **2015**, Article ID 88 (2015)
- O'Regan, D, Samet, B: Lyapunov-type inequalities for a class of fractional differential equations. *J. Inequal. Appl.* **2015**, Article ID 247 (2015)
- Rong, J, Bai, C: Lyapunov-type inequality for a fractional differential equation with fractional boundary conditions. *Adv. Differ. Equ.* **2015**, Article ID 82 (2015)
- Chdouh, A, Torres, DFM: A generalized Lyapunov's inequality for a fractional boundary value problem. *J. Comput. Appl. Math.* **312**, 192-197 (2017)
- Fečkan, M, Pospíšil, M: Note on fractional difference Gronwall inequalities. *Electron. J. Qual. Theory Differ. Equ.* **2014**, Article ID 44 (2014)

13. Khalil, R, Al Horani, M, Yousef, A, Sababheh, M: A new definition of fractional derivative. *J. Comput. Appl. Math.* **264**, 65-70 (2014)
14. Abdeljawad, T: On conformable fractional calculus. *J. Comput. Appl. Math.* **279**(1), 57-66 (2015)
15. Abdeljawad, T, Al Horani, M, Khalil, R: Conformable fractional semigroup operators. *J. Semigroup Theory Appl.* **2015**, Article ID 7 (2015)
16. Abu Hammad, M, Khalil, R: Abel's formula and Wronskian for conformable fractional differential equations. *Int. J. Differ. Equ. Appl.* **13**(3), 177-183 (2014)
17. Anderson, DR, Ulness, DJ: Newly defined conformable derivatives. *Adv. Dyn. Syst. Appl.* **10**(2), 109-137 (2015)
18. Pospíšil, M, Škripkova, LP: Sturm's theorems for conformable fractional differential equations. *Math. Commun.* **21**, 273-281 (2016)
19. Khaldi, R, Guezane-Lakoud, A: Lyapunov inequality for a boundary value problem involving conformable derivative. *Prog. Fract. Differ. Appl.* **3**(4), 323-329 (2017). doi:10.18576/pfda

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
