

RESEARCH ARTICLE

Open Access

DX5⁺NKT cells display phenotypical and functional differences between spleen and liver as well as NK1.1⁻Balb/c and NK1.1⁺ C57Bl/6 mice

Jens M Werner, Elisabeth Busl, Stefan A Farkas, Hans J Schlitt, Edward K Geissler and Matthias Hornung*

Abstract

Background: Natural killer T cells represent a linkage between innate and adaptive immunity. They are a heterogeneous population of specialized T lymphocytes composed of different subsets. DX5⁺NKT cells are characterized by expression of the NK cell marker DX5 in the context of CD3. However, little is known about the phenotype and functional capacity of this unique cell population. Therefore, we investigated the expression of several T cell and NK cell markers, as well as functional parameters in spleen and liver subsets of DX5⁺NKT cells in NK1.1⁻ Balb/c mice and compared our findings to NK1.1⁺ C57Bl/6 mice.

Results: In the spleen 34% of DX5⁺NKT cells expressed CD62L and they up-regulated the functional receptors CD154 as well as CD178 upon activation. In contrast, only a few liver DX5⁺NKT cells expressed CD62L, and they did not up-regulate CD154 upon activation. A further difference between spleen and liver subsets was observed in cytokine production. Spleen DX5⁺NKT cells produced more Th1 cytokines including IL-2, IFN- γ and TNF- α , while liver DX5⁺NKT cells secreted more Th2 cytokines (e.g. IL-4) and even the Th17 cytokine, IL-17a. Furthermore, we found inter-strain differences. In NK1.1⁺ C57Bl/6 mice DX5⁺NKT cells represented a distinct T cell population expressing less CD4 and more CD8. Accordingly, these cells showed a CD178 and Th2-type functional capacity upon activation.

Conclusion: These results show that DX5⁺NKT cells are a heterogeneous population, depending on the dedicated organ and mouse strain, that has diverse functional capacity.

Background

Natural killer T (NKT) cells represent a small but important subset of T lymphocytes with characteristics of both T and NK cells. They have potent immunoregulatory function that reportedly can promote cell-mediated immunity to tumors and infectious organisms and, paradoxically, suppress cell-mediated immunity associated with autoimmune disease and allograft rejection [1]. In mice, these cells express NK cell markers such as NK1.1 and CD94, as well as T-cell receptors (TCR) α/β with a restricted repertoire [2,3]. The invariant T cell receptor α chain V α 14-J α 18 with a conserved CDR3 region is associated with V β 8.2, V β 7 or V β 2 gene segments [3,4].

In contrast to conventional T-lymphocytes, the TCR of NKT cells does not interact with antigens presented by classical major histocompatibility complex (MHC)-encoded class I or II molecules. Instead, their TCR recognizes glycolipids presented by CD1d, which is a MHC class-I-like glycoprotein that belongs to a group of CD1 molecules associated with β 2-microglobulin [5-7]. CD1d is known to present lipids including glycosylceramides and glycosylphosphatidylinositol [8,9]. Activation via CD1d initiates the production of both Th1 (IFN) and Th2 cytokines (IL-4, IL-5, IL-13) [10], and increases the cytolytic activity of NKT cells [11].

NKT cells do not represent a homogeneous population. So far, three types of NKT cells have been described. First, there is an invariant V α 14-NKT cell (iV α 14-NKT) also called a type I NKT cell or iNKT cell. This group can be further differentiated into CD4⁺ single positive and CD4⁻CD8⁻ double-negative variants.

* Correspondence: matthias.hornung@klinik.uni-regensburg.de
Department of Surgery, University Hospital Regensburg, Franz-Josef-Strauss-Allee 11, 93053 Regensburg, Germany

Second, a population of CD1d-reactive NKT cells expressing diverse non-V α 14TCRs, referred to as type II NKT cells, has also been characterized. A third category has been termed NKT-like cells, which are CD1d-independent and express diverse TCRs [2].

Despite many years of NKT cell research, controversy remains about defining these cells. The expression of several surrogate markers, such as NK1.1 in C57Bl/6 mice and CD161 in humans, co-expressed with the TCR $_{\alpha/\beta}$, have been frequently used for NKT cell identification [2]. There are also NK1.1⁺ T cells which do not express the semiinvariant V α 14-J α 18 T cell receptor and are not CD1d-dependent, excluding their consideration as NKT cells.

A common marker for NKT cells in NK1.1⁻ mice strains is the antibody DX5, which recognizes the α_2 -integrin CD49b [12]. DX5 was initially characterized as a marker for NK cells [13] and more recently DX5 co-expressing CD3⁺ lymphocytes have been described [14]. Several studies, including some from our group, revealed evidence for immunoregulatory properties of these cells [15-18]. DX5⁺NKT cells produce Th1 and Th2 cytokines after stimulation like other NKT cells and seem to play a central role in anti-tumor immunity [13,19]. Recently published studies suggest an immune modulatory function of DX5⁺NKT cells after bone marrow and solid organ transplantation [20,21]. Although all these studies describe typical characteristics of NKT cells, Pellecci et al. showed that DX5⁺ T cell numbers were normal in CD1d^{-/-} and even in TCR J α 18^{-/-} mice [22]. Therefore, it is more likely that DX5⁺NKT cells belong to the third group of NKT-like cells [22,23].

In the present study we have further characterized DX5⁺CD3⁺ T cells referred to as DX5⁺NKT cells in NK1.1⁻ (Balb/c) and NK1.1⁺ (C57Bl/6) mice. For this purpose DX5⁺NKT cells were isolated from spleen and liver and several T and NK cell surface markers as well as maturation and activation markers were studied by flow cytometry. Distinct differences could be found between both subsets and between mouse strains.

Results

T cell marker expression is different between spleen and liver DX5⁺NKT cells

Lymphocytes were isolated from spleen and liver of Balb/c mice by MACS and FACS sorting (Figure 1A). First, expression of different T cell markers were analyzed to evaluate which subtype of NKT cells the DX5⁺NKT subset belong to. In the spleen of Balb/c mice, 93 \pm 1.4% of freshly isolated DX5⁺NKT cells expressed TCR $_{\alpha/\beta}$, with 18 \pm 1.2% of cells expressing V β 8.1/8.2 (Figure 1B). Surprisingly, most DX5⁺NKT cells expressed CD4 (82 \pm 0.85%), with 11 \pm 2% expressing CD8a and CD25. Interestingly, freshly isolated liver DX5

⁺NKT cells displayed a different receptor pattern. These cells expressed more V β 8.1/8.2 (36 \pm 3.4%, $P = 0.0159$) and less CD4 (67 \pm 2.6%, $P = 0.0079$).

Thus far, surface receptor expression was only examined in freshly isolated cells. Since DX5⁺NKT cells display their immunoregulating function especially upon activation [22], expression patterns after stimulation were analyzed. For this purpose, splenic and hepatic mononuclear cells were stimulated for 4 hours with anti-CD3 and anti-CD28 antibodies. Activation of spleen DX5⁺NKT cells from Balb/c mice (Figure 1C) resulted in a down-regulation of TCR $_{\alpha/\beta}$ (90 \pm 2% vs. 51 \pm 2%, $P = 0.004$) and an up-regulation of CD25 (14 \pm 2.6% vs. 27 \pm 3%, $P = 0.0411$). However, stimulation had little effect on liver derived cells of Balb/c mice. DX5⁺NKT cells revealed an expression pattern similar to that of freshly isolated cells: there was no down-regulation of TCR $_{\alpha/\beta}$ in liver derived DX5⁺NKT cells upon stimulation.

NK cell and activatory markers are expressed differently in spleen and liver DX5⁺NKT cells

To further characterize the differences between spleen and liver DX5⁺NKT cells, specific antibodies were used to detect different NK cell types, as well as maturation and activation markers by flow cytometry. In the spleen of Balb/c mice (Figure 2A) the NK cell marker CD94 was expressed on 8 \pm 1.8% of DX5⁺NKT cells. Again, DX5⁺NKT cells in the liver displayed differences. CD94 was expressed on significantly more cells than in the spleen (22 \pm 3.2%, $P = 0.0095$). After 4 h stimulation, no significant change in the NK cell receptor pattern was observed in splenic or hepatic DX5⁺NKT cells. As expected, the NK cell marker NK1.1 was expressed only on very few splenic (0.5 \pm 0.3%) and hepatic (0.8 \pm 0.6%) DX5⁺NKT cells (data not shown).

We also analyzed if the distinct phenotype of spleen and liver DX5⁺NKT cells was based on differences in the maturation status. Therefore, freshly isolated cells were stained with anti-CD103 and anti-CD62L antibodies (Figure 2B). Spleen DX5⁺NKT cells were positive for CD103 in 15 \pm 2.4% of cells. In contrast, freshly isolated liver DX5⁺NKT cells expressed this maturation marker less frequently (4.5 \pm 1.7%, $P = 0.014$). Freshly isolated spleen DX5⁺NKT cells were even more frequently positive for the maturation marker CD62L (34 \pm 5.5%) compared to liver DX5⁺NKT cells (3 \pm 2.2%, $P = 0.0079$). Upon 4 h stimulation, liver DX5⁺NKT cells displayed an increase in CD62L expression (14 \pm 1.7%, $P = 0.0317$), but still significantly less than the spleen subset (36 \pm 4%, $P = 0.0028$).

Next, differences in activation markers were evaluated. In freshly isolated DX5⁺NKT cells the activation marker CD38 was expressed on approximately 80% of splenic




Figure 1 FACS analysis of T cell marker from spleen and liver CD3⁺/DX5⁺NKT cells in Balb/c mice. DX5⁺NKT cells were isolated from spleen and liver of Balb/c mice using MACS and FACS-Sorting. Representative dot plot (A). Expression of different T cell markers revealed distinct subsets in Balb/c mice. In the spleen, more DX5⁺NKT cells appeared CD4⁺, in the liver, they expressed more V β 8.1/8.2 (B). Upon 4 h activation with anti-CD3 and anti-CD28, spleen DX5⁺NKT cells displayed a TCR α/β down-regulation, whereas liver DX5⁺NKT cells did not (C). Results are given as mean + SEM. Experiments were repeated at least three times (* p < 0.05).

and hepatic cells and displayed no significant change upon stimulation (data not shown). In contrast, upon 4 h stimulation, there was a strong up-regulation of the activating marker CD154 (3 ± 2.2% vs. 39 ± 5.8%, P = 0.0238) in spleen DX5⁺NKT cells (Figure 2C). However, liver DX5⁺NKT cells did not display a similar up-regulation of CD154 (4 ± 1%, P = 0.0498). Another activating marker, CD178, was also not significantly expressed on freshly isolated DX5⁺NKT cells, but upon 4 h stimulation, there was an up-regulation of CD178 in the spleen subset (2 ± 1.3% vs. 19 ± 1.4%, P = 0.0087).

Spleen and liver DX5⁺NKT cells display a distinct cytokine secretion pattern in Balb/c mice

Results thus far suggest distinct phenotypic differences between spleen and liver DX5⁺NKT cells. Next, the production of Th1 and Th2 cytokines were compared to

further assess functional differences between these subsets. Lymphocytes were isolated from spleen and liver of Balb/c mice and cultured for 4, 24 or 48 h in the presence of anti-CD3 and anti-CD28 antibodies. The cells were additionally incubated with PMA, ionomycin and GolgiPlug for intracellular cytokine staining.

As shown in Figure 3, we first stained for Th1 cytokines such as IL-2, IFN- γ and TNF α . After 4h stimulation, 33 ± 8% of spleen DX5⁺NKT cells produced IL-2. After 24 h, the number of IL-2-producing cells decreased (0.3 ± 0.1%, P = 0.0238). As expected, liver DX5⁺NKT cells showed a distinct Th1 cytokine profile. After 4 h, lower cell numbers produced IL-2 (11 ± 4.4%, P = 0.049) compared to spleen cells. Spleen DX5⁺NKT cells were also positive for IFN- γ after 4 h (46 ± 3.1%), 24 h (40 ± 4.6%) and 48 h (38 ± 6.6%). In contrast, liver DX5⁺NKT cells produced less frequently IFN- γ after 4 h




Figure 2 Activating markers are expressed differently in spleen and liver DX5⁺NKT cells. Detection of NK cell markers confirmed differences between spleen and liver subsets of DX5⁺NKT cells in NK1.1⁺Balb/c mice. CD94 expression was higher in liver DX5⁺NKT cells (A). Furthermore, the liver subset revealed less expression of maturation marker CD103 and CD62L compared to spleen (B). The activating markers CD154 and CD178 were in both freshly-isolated cell subsets only little expressed (C). Upon stimulation spleen DX5⁺NKT cells displayed an up-regulation of CD154, whereas the liver subset did not. Results are given as mean + SEM. Experiments were repeated at least three times (* p < 0.05).

(24 ± 2.4%, P = 0.0079) and 24 h (14.5 ± 2.2%, P = 0.0095). After 4 h splenic DX5⁺NKT cells produced also TNFα (49 ± 0.6%). However, the number of TNFα-producing cells decreased after 24 h (22 ± 5.1%, P = 0.0358). Liver DX5⁺NKT cells again displayed differences; after 4 h they produced slightly less TNFα compared to the spleen subset (39 ± 4.1%) and, after 24 h, the number of TNFα-producing cells decreased even more (4 ± 1.2%, P = 0.0159).

Next, Th2 cytokines such as IL-4, IL-6, IL-10 and IL-13 were analyzed. Only very few spleen DX5⁺NKT cells produced IL-4 after 4, 24 and 48 h. However, the number of IL-4 positive liver DX5⁺NKT cells increased after 48 h (11 ± 1.2%, P = 0.0286). Neither spleen nor liver DX5⁺NKT cells were significantly positive for IL-6 at any time point

(data not shown). Spleen DX5⁺NKT cells showed a significant increase in the frequency of IL-10-producing cells after 24 h (13 ± 2.1%) compared to 4 h (0.4 ± 0.2%, P = 0.0187), and compared to liver DX5⁺NKT cells after 24 h stimulation (1 ± 0.6%, P = 0.0357). In the spleen the number of IL-13-producing DX5⁺NKT cells significantly increased after 48 h compared to 4 h (20 ± 3% vs. 10 ± 1.2%, P = 0.0061), whereas in liver the number of IL-13 positive DX5⁺NKT cells peaked after 4 h compared to 24 h (16 ± 2.1% vs. 3 ± 1.2%, P = 0.0286). Furthermore, we tested for IL-17a as a Th17-type cytokine. Liver DX5⁺NKT cells after 4 h more frequently produced IL-17a (7 ± 1%) compared to the splenic-derived cells (1 ± 0.7%, P = 0.0061); after 24 h, the frequency of liver-derived IL-17a-producing DX5⁺NKT cells decreased (0.3 ± 0.3%, P = 0.286).




Figure 3 Spleen and liver DX5⁺NKT cells display a distinct cytokine secretion pattern in Balb/c mice. Isolated DX5⁺NKT cells were stimulated with antibodies against CD3 and CD28 in the presence of IL-2 for 4, 24 and 48 h, respectively. Control cells were cultured simultaneously without stimulation. Values were subtracted as background. FACS analysis revealed again distinct differences between spleen and liver subsets of DX5⁺NKT cells. Results are given as mean + SEM. Experiments were repeated at least three times (* p < 0.05).

NK1.1⁺C57Bl/6 mice reveal distinct surface marker expression on DX5⁺NKT cells

Data thus far were obtained using a NK1.1⁻ mouse strain. These findings were compared to a second mouse strain known to express NK1.1, C57Bl/6 mice. Several T cell markers were analyzed. A down-regulation of TCR α/β (65 \pm 4.8% vs. 89 \pm 2.1%, P = 0.01) was also observed upon activation of spleen DX5⁺NKT cells from C57Bl/6 mice (Figure 4A). Compared to Balb/c, fewer freshly isolated spleen DX5⁺NKT cells expressed CD4 (82 \pm 1% vs. 47 \pm 1.1%, P = 0.0043), but more expressed CD8a (11 \pm 1.6% vs. 32 \pm 0.8%, P = 0.0043). NK cell markers showed significant differences between both mouse strains (Figure 4B). In C57Bl/6 mice 33 \pm 2.8% of freshly isolated spleen DX5⁺NKT cells were stained for CD94 (P = 0.0095) and 39 \pm 1% for NK1.1 (P = 0.0022). Compared to Balb/c mice (Figure 4C), fewer spleen DX5⁺NKT cells displayed the maturation marker CD103 (15 \pm 1.9% vs. 6 \pm 1.6%, P = 0.0030), but more expressed CD62L (33 \pm 5.5% vs. 47 \pm 1%, P = 0.0047). In terms of activating markers, both mouse strains showed little expression of CD154 or CD178 in freshly isolated DX5⁺NKT cells (Figure 4D). However, upon activation, the up-regulation of CD154 (39 \pm 5.8% vs. 13 \pm 1.2%, P = 0.0012) was less distinctive in C57Bl/

6 mice, whereas CD178 expression was similar to Balb/c mice (19 \pm 1.4% vs. 18 \pm 1.2%).

Cytokine production of spleen DX5⁺NKT cells is different in NK1.1⁻Balb/c and NK1.1⁺ C57Bl/6 mice

Results have already shown distinct phenotypic differences in DX5⁺NKT cells between NK1.1⁻ and NK1.1⁺ mouse strains. Next, cytokine production was compared to further assess potential functional differences between these subsets. Lymphocytes were isolated from spleens of Balb/c and C57Bl/6 mice and cultured for 4, 24 or 48 h. As shown in Figure 5 spleen DX5⁺NKT cells from the NK1.1⁺ mouse strain displayed a distinct Th1 cytokine profile. Compared to NK1.1⁻Balb/c mice, DX5⁺NKT cells less often produced IL-2 (33 \pm 8.6% vs. 1 \pm 0.5%, P = 0.0357) and TNF- α (49 \pm 0.6% vs. 14 \pm 8.8%, P = 0.0167) after 4h; however, production of IFN- γ was similar. In terms of Th2 cytokines, there was also no significant IL-4 production in spleen DX5⁺NKT cells from the NK1.1⁺ mouse strain. There were some differences in Th2 cytokine production between NK1.1⁻ and NK1.1⁺ mice. After 24 h, spleen DX5⁺NKT cells in NK1.1⁺ C57Bl/6 mice less frequently produced IL-10 (13 \pm 2.1% vs. 1 \pm 0.3%, P = 0.0357) and IL-13 (14 \pm 2.9% vs. 4 \pm 0.9%, P = 0.0381). The number of splenic-





Figure 4 FACS-Analysis of spleen DX5⁺NKT cells revealed interstrain differences in several phenotypical markers between NK1.1⁻Balb/c and NK1.1⁺C57Bl/6 mice. In the NK1.1⁺ mice strain spleen DX5⁺NKT cells expressed less CD4 but more CD8 (A). Upon activation the TCR α/β down-regulation was confirmed. In addition to NK1.1 more spleen DX5⁺NKT cells in C57Bl/6 mice were positive for CD94 (B). Maturation marker CD62L was slightly higher expressed in freshly-isolated spleen DX5⁺NKT cells in C57Bl/6 mice compared to Balb/c, whereas CD103 was less (C). Functional marker CD154 and CD178 were only little expressed in freshly-isolated spleen DX5⁺NKT cells in both mice strains (D). Upon stimulation spleen DX5⁺NKT cells in NK1.1⁺ C57Bl/6 mice displayed less up-regulation of CD154 compared to Balb/c. Results are given as mean + SEM. Experiments were repeated at least three times (* p < 0.05).



derived IL-17a-producing DX5⁺NKT cells was similar in both mouse strains. Only a few spleen DX5⁺NKT cells produced IL17a after 4 h and 24 h.

Discussion

All freshly isolated DX5⁺NKT cells from NK1.1 deficient Balb/c mice were positive for TCR_{α/β}, as expected for type III NKT cells [2,3,24]. Most of them additionally expressed CD4, and a few expressed CD8a and CD25. As mentioned in other reports for spleen and thymus [25], a difference between spleen and liver subsets was observed. Liver DX5⁺NKT cells expressed less CD4 and CD62L. Since expression of CD62L is associated with naivety or homing to peripheral lymph nodes in iNKT-cells [26], and CD4 becomes down-regulated after repeated activation [27], this suggests a higher maturational status of liver DX5⁺NKT cells in Balb/c mice. This is supported by the finding that Vβ8.2-containing

Vα14i TCRs in NKT cells have a higher antigen affinity than those containing Vβ7 or Vβ2 [28,29]. Consistent with the idea of a negative selection, liver DX5⁺NKT cells displayed a higher expression of Vβ8.1/8.2.

Upon activation, a down-regulation of TCR_{α/β} was observed in spleen DX5⁺NKT cells in Balb/c mice. A decrease of TCR_{α/β}-expression was also reported for NK1.1⁺NKT cells [30]. Due to a lack of detection, a rapid apoptotic death of this cell was initially claimed upon stimulation [30]. More recently TCR_{α/β}-down-regulation in spleen and liver NK1.1⁺NKT cells of C57Bl/6 mice was discovered to be the reason for this assumed cell disappearance [31,32]. However, our study does not confirm this finding for liver DX5⁺NKT cells in Balb/c mice, supporting evidence that there are distinct subsets of DX5⁺NKT cells in the spleen and liver. CD154 expression was also increased upon stimulation. As the ligand of CD40, CD154 represents a co-stimulatory

molecule and is expressed by activated and regulatory T cells [33,34]. Furthermore, CD178, the ligand of Fas responsible for the extrinsic induction of apoptosis (e.g. expressed on CD8⁺ cytotoxic T cells) was up-regulated [35]. This up-regulation was more evident in spleen than in liver DX5⁺NKT cells and is also reported for NK1.1⁺NKT cells [36]. Taken together, liver DX5⁺NKT cells are more mature and display a different activating phenotype upon stimulation compared to the spleen subset.

Since NKT cells are supposed to link innate and adaptive immunity by rapid secretion of cytokines, including IFN γ , TNF α , IL-4, IL-10, IL-13, GM-CSF and IL-2 [24,37,38], we further assessed differences between spleen and liver DX5⁺NKT cells on a functional basis. Fewer liver DX5⁺NKT cells in Balb/c mice produced Th1 cytokines such as IL-2, IFN- γ , and TNF- α . In terms of Th2 cytokines, more liver DX5⁺NKT cells produced IL-4, but fewer produced IL-10 compared to the spleen subset. Charbonnier et al. described an induction of IL-10 production in liver CD49b⁺CD4⁺ T cells by immature dendritic cell vaccination [18]. However, we did not confirm this finding for liver DX5⁺NKT cells in Balb/c mice after anti-CD3/anti-CD28 stimulation. Collectively, the limited Th1 cytokine secretion upon stimulation confirmed the observation that a different activating phenotype of DX5⁺NKT cells exists in the liver of Balb/c mice.

Recently, type I NKT cells have been shown to be capable of producing IL-17, potentially implicating this cell type in inflammatory conditions [39,40]. We showed that a subset of DX5⁺NKT cells in the liver of Balb/c mice has to be taken into account as an IL-17 producing cell. These results are consistent with our result regarding a more mature phenotype of liver-derived DX5⁺NKT cells, with a different phenotypical profile upon stimulation compared to the spleen subset.

In NK1.1⁺ C57Bl/6 mice fewer spleen DX5⁺NKT cells revealed expression of CD4, but more of CD8a. These findings are consistent with data from Pellici et al., who observed similar differences in C57Bl/6 mice [22]. In accordance with Conzalez et al., 40-50% of DX5⁺NKT cells express NK1.1 [15]. Taking these results together with the higher CD4 expression in NK1.1-deficient Balb/c mice, we suggest that these mice compensate for a reduced number of NK1.1⁺NKT cells with a higher prevalence of CD4⁺DX5⁺NKT cells. In contrast to results from Gonzalez et al. [15], about one-third of these cells in the spleen of C57Bl/6 mice, and 11% in Balb/c mice, were CD8⁺.

Upon activation a down-regulation of TCR α/β was observed in spleen DX5⁺NKT cells in Balb/c and in C57Bl/6 mice, as it has been reported for NK1.1⁺NKT cells [30]. Furthermore, upon a 4 h stimulation there

was a notable up-regulation of CD25 in Balb/c versus C57Bl/6 mice. CD25, the IL-2 receptor α -chain, is proposed to be a phenotypical marker of regulatory T cells [41-43]. The up-regulation we observed especially for spleen DX5⁺NKT cells in Balb/c mice could be either due to activation and increased receptor expression based on the regulatory background of DX5⁺NKT cells or due to a natural selection because of a higher apoptotic resistance in CD4⁺CD25⁺ T cells [44]. Another marker for activation on regulatory T cells is CD154 [34]. CD154 expression was increased upon stimulation in spleen DX5⁺NKT cells in Balb/c mice, but was significantly less prominent in C57Bl/6 mice. Upon 4 h stimulation, CD178 (ligand of Fas) was similarly up-regulated [35] on splenic DX5⁺NKT cells in both mice strains. However, after 24 h stimulation CD178 expression was decreased in Balb/c versus C57Bl/6 mice, and CD154 was further up-regulated in Balb/c, but down-regulated in C57Bl/6 mice (data not shown). These changes suggest a more regulatory CD4⁺CD25⁺ phenotype in NK1.1⁻ Balb/c and a more CD8⁺ and CD178⁺ driven cytotoxic component in the NK1.1⁺ C57Bl/6 mouse strain [45].

High IL-2 production by DX5⁺NKT cells in the early phase of activation was observed only in Balb/c mice, further supporting our findings of interstrain differences. Moreover, splenic DX5⁺NKT cells in C57Bl/6 mice displayed a more one-sided Th1 cytokine profile, with mostly IFN γ and TNF α producing cells, which is also described for CD8⁺ T cells [46]. DX5⁺NKT cells in Balb/c mice displayed a more polyfunctional cytokine profile. In the late phase of stimulation, IFN γ and TNF α production decreased and IL-10 and IL-13 secretion increased. In contrast to liver CD49b⁺CD4⁺ cells [18], and supporting our data regarding splenic DX5⁺NKT cells, Kassiotis et al. revealed no IL-10 production for splenic CD49b⁺CD4⁺ T cells in C57Bl/6 mice [47]. Furthermore, Pellicci et al. could not detect any significant IL-10 production in DX5⁺TCR α/β ⁺CD1d-Tet⁺ cells in the same mouse strain [22]. Confirming our finding and giving again evidence for a difference between both mouse strains, we showed a significant increase in IL-10 production after 24h stimulation for splenic DX5⁺NKT cells in Balb/c mice. Taken together, these results suggesting a more one-sided Th1 cytokine production from DX5⁺NKT cells in C57Bl/6 mice confirm their more CD8 and CD178 expressing phenotype.

Conclusions

In conclusion, our data show that distinct subsets of DX5⁺NKT cells exist within NK1.1⁻ Balb/c mice. In the spleen these cells appear more naive with a higher CD62L expression and an up-regulation of CD154 and CD178 upon activation. In contrast, in the liver DX5

*NKT cells are more mature and display a different regulatory potency upon stimulation. Furthermore, there are remarkable inter-strain differences. In NK1.1⁺ C57Bl/6 mice, fewer DX5⁺NKT cells are CD4⁺ and more are CD8⁺, confirming their CD178 and Th1-sided functional behavior upon activation.

Methods

Cell harvesting and isolation

All experiments were approved by the institutional animal care committee at the University Hospital Regensburg. Different lymphocyte subsets were purified from splenic or hepatic mononuclear cells isolated from Balb/c mice or C57Bl/6NCRl mice (Charles River Laboratories, Wilmington, MA, USA). If necessary, further isolation was performed by magnetic activated cell sorting (MACS; Miltenyi Biotec, Bergisch Gladbach, Germany) or by FACS (FACSaria I, BD Bioscience, San Jose, CA, USA). Briefly, cell suspension of the spleen was prepared by cutting the tissue into small pieces and gently pressing through a 100 µm wire mesh. For preparation of the liver the portal vein was flushed with 10 ml sodium chloride until the organ became pale. Then the liver was cut into small parts and passed through a 100 µm wire mesh. Hepatic lymphocytes were further isolated by density gradient centrifugation using 80% and 40% Percoll (Biochrom, Berlin, Germany). For further isolation, DX5⁺ cells were purified using anti-mouse-DX5⁺ MicroBeads (Miltenyi Biotec). Cells were passed through a MACS-column (type LS) attached to a Midi-MACS-magnet (Miltenyi Biotec). DX5⁺ cells were labeled with FITC-conjugated anti-mouse CD3 (clone: 17A2, rat IgG2b) and PE-conjugated anti-mouse CD49b (clone: DX5, rat IgM) (all from BD Biosciences) for further CD3⁺DX5⁺ enrichment by FACS, and are referred to as DX5⁺NKT cells throughout the study.

Antibodies and flow cytometry

The following reagents were used for cell surface labeling in multiparameter flow cytometric analysis (FACS Calibur, BD Bioscience): FITC or Alexa Fluor 647-conjugated anti-mouse CD3 (clone: 17A2, rat IgG2b), APC-conjugated anti-mouse-CD4 (clone: GK1.5, rat IgG2b), APC-conjugated anti-mouse-CD8a (clone: 53-6.7, rat IgG2a), PE-conjugated anti-mouse-CD38 (clone: 90, rat IgG2a), PE-conjugated anti-mouse-CD49b (clone: DX5, rat IgM), APC-conjugated anti-mouse-CD62L (clone: MEL-14, rat IgG2a), PE-conjugated anti-mouse-CD103 (clone: M290, rat IgG2a), PE-conjugated anti-mouse-CD178 (Fas-Ligand) (clone: MFL3, hamster IgG1), PE-conjugated anti-mouse- β 8.1, 8.2 TCR (clone: MR5-2, rat IgG2a) all BD Biosciences. APC-conjugated anti-mouse-CD49b (clone: DX5, rat IgM), FITC-conjugated anti-mouse-CD49b (clone: DX5, rat IgM) all obtained from Miltenyi Biotec. FITC-

conjugated anti-mouse-CD94 (clone: 18d3, rat IgG2a), APC-conjugated anti-mouse-CD154 (CD40L) (clone: MR1, hamster IgG), APC-conjugated anti-mouse-NK1.1 (clone: PK136, mouse IgG2a) all obtained from eBioscience (San Diego, CA, USA). APC-conjugated anti-mouse-CD25 (clone: PC61 5.3, rat IgG1), FITC-conjugated anti-mouse-TCR α/β (clone: HM 3601, hamster IgG) from Caltag (Towcester, UK).

Activation of DX5⁺NKT cells

For lymphocyte stimulation, 96 well cell culture plates (Corning costar, Sigma Aldrich) were coated with anti-mouse-CD3e (Clone: 145-2C11, BD Biosciences) at 10 µg/ml and stored overnight at 4°C. Wells were washed twice with PBS. And up to 1×10^6 isolated lymphocytes from spleen and liver were plated in 200 µl RPMI culture medium (Biochrom). For activation, 5 µg/ml of anti-mouse-CD28 (clone: 37.51, BD Biosciences) and 2000 IU/ml IL-2 (Peprotech, Rocky Hill, NJ, USA) were added. Plates were incubated for the indicated time at 37°C and 5% CO₂.

Intracellular cytokine staining

4×10^5 DX5⁺-NKT cells were incubated in 200 µl culture medium in a 96 well plate as mentioned above. Additionally, 50 ng/ml PMA (InvivoGen, San Diego, CA, USA) was added at the beginning, with 750 ng/ml ionomycin (Sigma-Aldrich, St. Louis, USA) being added for the last 4h; 1 µg/ml GolgiPlug (BD Bioscience) was added 2 h before cell harvesting. Culture supernatants were harvested and stored at -20°C for IFN γ ELISA. Cells were fixed in 1ml Fix/Perm (eBioscience) for 60 min at 4°C. After incubation with permeabilization buffer (eBioscience) cells were stained with PE-conjugated anti-mouse-cytokine Abs (IL-2, clone: JES6-5H4/IL-4, clone: BVD4-1D11/IL-6, clone: MP5-20F3/IL-10, clone: JES5-16E3/IFN γ , clone: XMG1.2/TNF α , clone: MP6-XT22) from BD Bioscience and with PE-conjugated anti-mouse-Abs (IL-13, clone: eBio13A/IL-17a, clone: eBio17B7) and FITC-conjugated anti-mouse-IFN γ (clone: XMG1.2) (all eBioscience).

IFN γ analysis by Enzyme-Linked Immunosorbent Assay

Stimulation of DX5⁺NKT cells was confirmed by IFN γ ELISA. IFN γ concentration from harvested supernatants was analyzed using a commercially available sandwich ELISA kit (BD Bioscience). Tetramethylbenzidine dihydrochloride was used for detection. ELISA readings were determined by OD scanning at 450 nm using an Emax precision microplate reader from Molecular Devices (Downingtown, PA, USA).

Statistics

All in vitro experiments were repeated at least 3 times and data are presented as the mean value \pm SEM.

Statistical analyses were performed using either a student's t-test or the Mann-Whitney-U-test. Differences were considered significant at $P < 0.05$.

Acknowledgements

We wish to thank Joachim Schweimer for his excellent technical assistance. This work was supported by funds from the "Regensburger Forschungsförderung Medizin (ReForM)".

Authors' contributions

JMW carried out FACS analysis, participated in the design of the study, data analysis and the preparation of the manuscript. EB carried out isolation of cells and FACS analysis. SAF participated in the design and coordination of the study and helped to draft the manuscript. HJS participated in the design of the study and revised the manuscript and gave important intellectual content. EKG has made substantial contributions to the conception and the design of the study, to the analysis and interpretation of the data and revised the manuscript. MH conceived of the study, and participated in its design and coordination, analyzed the data and drafted the manuscript. All authors read and approved the final manuscript.

Received: 9 January 2011 Accepted: 29 April 2011

Published: 29 April 2011

References

- Godfrey DI, Stankovic S, Baxter AG: **Raising the NKT cell family.** *Nat Immunol* 2004, **5**:1197-1206.
- Godfrey DI, MacDonald HR, Kronenberg M, Smyth MJ, Van Kaer L: **NKT cells: what's in a name?** *Nat Rev Immunol* 2004, **4**:231-7.
- Godfrey DI, Hammond KJ, Poulton LD, Smyth MJ, Baxter AG: **NKT cells: facts, functions and fallacies.** *Immunol Today* 2000, **21**:573-83.
- Lantz O, Bendelac A: **An invariant T cell receptor alpha chain is used by a unique subset of major histocompatibility complex class I-specific CD4+ and CD4-8- T cells in mice and humans.** *J Exp Med* 1994, **180**:1097-106.
- Brigl M, Brenner MB: **CD1: antigen presentation and T cell function.** *Annu Rev Immunol* 2004, **22**:817-90.
- Burdin N, Brossay L, Degano M, Iijima H, Gui M, Wilson IA, Kronenberg M: **Structural requirements for antigen presentation by mouse CD1.** *Proc Natl Acad Sci USA* 2000, **97**:10156-61.
- Burdin N, Kronenberg M: **CD1-mediated immune responses to glycolipids.** *Curr Opin Immunol* 1999, **11**:326-31.
- Joyce S, Woods AS, Yewdell JW, Bennink JR, De Silva AD, Boesteanu A, Balk SP, Cotter RJ, Brutkiewicz RR: **Natural ligand of mouse CD1d1: cellular glycosylphosphatidylinositol.** *Science* 1998, **279**:1541-4.
- Kawano T, Tanaka Y, Shimizu E, Kaneko Y, Kamata N, Sato H, Osada H, Sekiya S, Nakayama T, Taniguchi M: **A novel recognition motif of human NKT antigen receptor for a glycolipid ligand.** *Int Immunol* 1999, **11**:881-7.
- Benlagha K, Weiss A, Beavis A, Teyton L, Bendelac A: **In vivo identification of glycolipid antigen-specific T cells using fluorescent CD1d tetramers.** *J Exp Med* 2000, **191**:1895-903.
- Cui J, Shin T, Kawano T, Sato H, Kondo E, Taura I, Kaneko Y, Koseki H, Kanno M, Taniguchi M: **Requirement for Valpha14 NKT cells in IL-12-mediated rejection of tumors.** *Science* 1997, **278**:1623-6.
- Arase H, Saito T, Phillips JH, Lanier LL: **Cutting edge: the mouse NK cell-associated antigen recognized by DX5 monoclonal antibody is CD49b (alpha 2 integrin, very late antigen-2).** *J Immunol* 2001, **167**:1141-4.
- Moodycliffe AM, Maiti S, Ullrich SE: **Splenic NK1.1-negative, TCR alpha beta intermediate CD4+ T cells exist in naive NK1.1 allelic positive and negative mice, with the capacity to rapidly secrete large amounts of IL-4 and IFN-gamma upon primary TCR stimulation.** *J Immunol* 1999, **162**:5156-63.
- Ortaldo JR, Winkler-Pickett R, Mason AT, Mason LH: **The Ly-49 family: regulation of cytotoxicity and cytokine production in murine CD3+ cells.** *J Immunol* 1998, **160**:1158-65.
- Gonzalez A, Andre-Schmutz I, Carnaud C, Mathis D, Benoist C: **Damage control, rather than unresponsiveness, effected by protective DX5+ T cells in autoimmune diabetes.** *Nat Immunol* 2001, **2**:1117-25.
- Adachi K, Tsutsui H, Seki E, Nakano H, Takeda K, Okumura K, Van Kaer L, Nakanishi K: **Contribution of CD1d-unrestricted hepatic DX5+ NKT cells to liver injury in Plasmodium berghei-parasitized erythrocyte-injected mice.** *Int Immunol* 2004, **16**:787-98.
- Hornung M, Farkas SA, Sattler C, Schlitt HJ, Geissler EK: **DX5+ NKT cells induce the death of colitis-associated cells: involvement of programmed death ligand-1.** *Eur J Immunol* 2006, **36**:1210-21.
- Charbonnier LM, van Duivenvoorde LM, Apparailly F, Cantos C, Han WG, Noel D, Duperray C, Huizinga TW, Toes RE, Jorgensen C, et al: **Immature dendritic cells suppress collagen-induced arthritis by in vivo expansion of CD49b+ regulatory T cells.** *J Immunol* 2006, **177**:3806-13.
- Osada T, Nagawa H, Shibata Y: **Tumor-infiltrating effector cells of alpha-galactosylceramide-induced antitumor immunity in metastatic liver tumor.** *J Immune Based Ther Vaccines* 2004, **2**:7.
- Lan F, Zeng D, Higuchi M, Higgins JP, Strober S: **Host conditioning with total lymphoid irradiation and antithymocyte globulin prevents graft-versus-host disease: the role of CD1-reactive natural killer T cells.** *Biol Blood Marrow Transplant* 2003, **9**:355-63.
- Margalit M, Ilan Y, Ohana M, Safadi R, Alper R, Sherman Y, Doviner V, Rabbani E, Engelhardt D, Nagler A: **Adoptive transfer of small numbers of DX5+ cells alleviates graft-versus-host disease in a murine model of semiallogeneic bone marrow transplantation: a potential role for NKT lymphocytes.** *Bone Marrow Transplant* 2005, **35**:191-7.
- Pellicci DG, Hammond KJ, Coquet J, Kyprisoudis K, Brooks AG, Kedzierska K, Keating R, Turner S, Berzins S, Smyth MJ, et al: **DX5/CD49b-positive T cells are not synonymous with CD1d-dependent NKT cells.** *J Immunol* 2005, **175**:4416-25.
- Wingender G, Kronenberg M: **Role of NKT cells in the digestive system. IV. The role of canonical natural killer T cells in mucosal immunity and inflammation.** *Am J Physiol Gastrointest Liver Physiol* 2008, **294**:G1-8.
- Kronenberg M, Gapin L: **The unconventional lifestyle of NKT cells.** *Nat Rev Immunol* 2002, **2**:557-68.
- Yang Y, Ueno A, Bao M, Wang Z, Im JS, Porcelli S, Yoon JW: **Control of NKT cell differentiation by tissue-specific microenvironments.** *J Immunol* 2003, **171**:5913-20.
- Kim CH, Johnston B, Butcher EC: **Trafficking machinery of NKT cells: shared and differential chemokine receptor expression among V alpha 24(+)V beta 11(+) NKT cell subsets with distinct cytokine-producing capacity.** *Blood* 2002, **100**:11-6.
- Chen H, Huang H, Paul WE: **NK1.1+ CD4+ T cells lose NK1.1 expression upon in vitro activation.** *J Immunol* 1997, **158**:5112-9.
- Schumann J, Voyle RB, Wei BY, MacDonald HR: **Cutting edge: influence of the TCR V beta domain on the avidity of CD1d:alpha-galactosylceramide binding by invariant V alpha 14 NKT cells.** *J Immunol* 2003, **170**:5815-9.
- Chun T, Page MJ, Gapin L, Matsuda JL, Xu H, Nguyen H, Kang HS, Stanic AK, Joyce S, Koltun WA, et al: **CD1d-expressing dendritic cells but not thymic epithelial cells can mediate negative selection of NKT cells.** *J Exp Med* 2003, **197**:907-18.
- Eberl G, MacDonald HR: **Rapid death and regeneration of NKT cells in anti-CD3epsilon- or IL-12-treated mice: a major role for bone marrow in NKT cell homeostasis.** *Immunity* 1998, **9**:345-53.
- Wilson MT, Johansson C, Olivares-Villagomez D, Singh AK, Stanic AK, Wang CR, Joyce S, Wick MJ, Van Kaer L: **The response of natural killer T cells to glycolipid antigens is characterized by surface receptor down-modulation and expansion.** *Proc Natl Acad Sci USA* 2003, **100**:10913-8.
- Harada M, Seino K, Wakao H, Sakata S, Ishizuka Y, Ito T, Kojo S, Nakayama T, Taniguchi M: **Down-regulation of the invariant Valpha14 antigen receptor in NKT cells upon activation.** *Int Immunol* 2004, **16**:241-7.
- Elgueta R, Benson MJ, de Vries VC, Wasiuk A, Guo Y, Noelle RJ: **Molecular mechanism and function of CD40/CD40L engagement in the immune system.** *Immunol Rev* 2009, **229**:152-72.
- Hornung M, Lindemann D, Kraus C, Peters A, Berberich I: **The CD40 TRAF family member interacting motif carries the information to rescue WEHI 231 cells from anti-IGM-induced growth arrest.** *Eur J Immunol* 1998, **28**:3812-23.
- Falschlehner C, Ganten TM, Koschny R, Schaefer U, Walczak H: **TRAIL and Other TRAIL Receptor Agonists as Novel Cancer Therapeutics.** *Adv Exp Med Biol* 2009, **647**:195-206.
- Nyambayar D, Iwabuchi K, Hedlund E, Murakawa S, Shirai K, Iwabuchi C, Kon Y, Miyazaki Y, Yanagawa Y, Onoe K: **Characterization of NKT-cell hybridomas expressing invariant T-cell antigen receptors.** *J Clin Exp Hematop* 2007, **47**:1-8.
- Schoenborn JR, Wilson CB: **Regulation of interferon-gamma during innate and adaptive immune responses.** *Adv Immunol* 2007, **96**:41-101.

38. Kronenberg M: **Toward an understanding of NKT cell biology: progress and paradoxes.** *Annu Rev Immunol* 2005, **23**:877-900.
39. Steinman L: **A brief history of T(H)17, the first major revision in the T(H)1/T(H)2 hypothesis of T cell-mediated tissue damage.** *Nat Med* 2007, **13**:139-45.
40. Mills KH: **Induction, function and regulation of IL-17-producing T cells.** *Eur J Immunol* 2008, **38**:2636-49.
41. Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M: **Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases.** *J Immunol* 1995, **155**:1151-64.
42. Suri-Payer E, Amar AZ, Thornton AM, Shevach EM: **CD4+CD25+ T cells inhibit both the induction and effector function of autoreactive T cells and represent a unique lineage of immunoregulatory cells.** *J Immunol* 1998, **160**:1212-8.
43. Shevach EM: **Certified professionals: CD4(+)CD25(+) suppressor T cells.** *J Exp Med* 2001, **193**:F41-6.
44. Papiernik M, de Moraes ML, Pontoux C, Vasseur F, Penit C: **Regulatory CD4 T cells: expression of IL-2R alpha chain, resistance to clonal deletion and IL-2 dependency.** *Int Immunol* 1998, **10**:371-8.
45. Glimcher LH, Townsend MJ, Sullivan BM, Lord GM: **Recent developments in the transcriptional regulation of cytolytic effector cells.** *Nat Rev Immunol* 2004, **4**:900-11.
46. Pearce EL, Mullen AC, Martins GA, Krawczyk CM, Hutchins AS, Zediak VP, Banica M, DiCioccio CB, Gross DA, Mao CA, *et al*: **Control of effector CD8+ T cell function by the transcription factor Eomesodermin.** *Science* 2003, **302**:1041-3.
47. Kassiotis G, Gray D, Kiafard Z, Zwirner J, Stockinger B: **Functional specialization of memory Th cells revealed by expression of integrin CD49b.** *J Immunol* 2006, **177**:968-75.

doi:10.1186/1471-2172-12-26

Cite this article as: Werner *et al.*: DX5⁺NKT cells display phenotypical and functional differences between spleen and liver as well as NK1.1⁻ Balb/c and NK1.1⁺ C57Bl/6 mice. *BMC Immunology* 2011 **12**:26.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

