

Research article

Open Access

Expression of HNF4alpha in the human and rat choroid plexus – Implications for drug transport across the blood-cerebrospinal-fluid (CSF) barrier

Monika Niehof¹ and Jürgen Borlak*^{1,2}

Address: ¹Fraunhofer Institute for Toxicology and Experimental Medicine, Center of Molecular Medicine and Medical Biotechnology, Nikolai-Fuchs-Str. 1, 30625 Hannover, Germany and ²Center of Pharmacology and Toxicology, Medical School of Hannover, Carl-Neuberg-Str. 1, 30625 Hannover, Germany

Email: Monika Niehof - niehof@item.fraunhofer.de; Jürgen Borlak* - juergen.borlak@item.fraunhofer.de

* Corresponding author

Published: 3 July 2009

Received: 1 December 2008

BMC Molecular Biology 2009, **10**:68 doi:10.1186/1471-2199-10-68

Accepted: 3 July 2009

This article is available from: <http://www.biomedcentral.com/1471-2199/10/68>

© 2009 Niehof and Borlak; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The choroid plexus consists of highly differentiated epithelium and functions as a barrier at the interface of the blood-cerebrospinal-fluid (CSF). This tissue may therefore determine the bioavailability and transport of drugs to the brain. Little is known about the expression of drug and xenobiotic metabolizing enzymes (DME) and of drug transporters in the human choroid plexus. Notably, the transcription factor and zinc finger protein HNF4alpha is a master regulator of DMEs and of drug transporters. As of today its activity in the blood-CSF barrier is unknown. Here we report our efforts in determining HNF4alpha activity in the regulation of ABC transporters in the human and rat choroid plexus.

Results: We report expression of HNF4alpha by qRT-PCR and by immunohistochemistry and evidence transcript expression of the ATP-binding cassette transporters ABCB1, ABCB4, ABCC1-6 in choroid plexus. Additionally, HNF4alpha DNA binding activity at regulatory sequences of ABCB4 and ABCC1 was determined by EMSA bandshift assays with a specific antibody. We then performed siRNA mediated functional knock down of HNF4alpha in Caco-2 cells and found ABCC1 gene expression to be repressed in cell culture experiments.

Conclusion: Our study evidences activity of HNF4alpha in human and rat choroid plexus. This transcription factor targets DMEs and drug transporters and may well determine availability of drugs at the blood-CSF barrier.

Background

Drug delivery to the brain is mediated by several factors, most notably transport across the blood brain (BB) and the choroid plexus barrier; the latter displays drug-metabolizing enzyme and drug transport activity. It may therefore determine the overall cerebral bioavailability of drugs [1]. Specifically, the choroid plexus is located within brain

vesicles. It is composed of a tight monolayer of polarized epithelial cells and forms the blood-cerebrospinal-fluid (CSF) barrier. Together with the blood-brain barrier, formed by endothelial cells of brain capillaries, it functions as the main interface between the central nervous system (CNS) and the peripheral circulation. Within the CNS this tissue is of great pharmacological interest, but

information on the expression of efflux transporters such as the ATP binding cassette (ABC) proteins is missing [2]. In contrast, their expression in liver, kidney, and intestine has been studied in considerable detail [3-5]. Indeed, the ABC drug transporters extrude a variety of structurally diverse drugs, drug conjugates and metabolites in an active, ATP-dependent manner and even against high concentration gradients. The three main ABC families considered to be involved in the disposition of xenobiotics include the ABCB family (MDR subfamily, multidrug resistance, e.g. ABCB1/MDR1), the ABCC family of multidrug resistance proteins (MRP subfamily, multidrug resistance related proteins, e.g. ABCC2/MRP2), and the breast cancer resistance protein (ABCG2/BCRP) of the ABCG family [3,4]. However, comprehensive studies on the expression levels of ATP transporters in the human choroid plexus have not been attempted.

Notably, there is clear evidence for HNF4 α to play a role in the transcriptional control of drug transporters. Specifically, HNF4 α is a member of the nuclear receptor superfamily and one of the key players in liver biology [6-8]. Among the genes regulated by HNF4 α are a broad range of xenobiotic-metabolizing cytochrome P450 isozymes [9,10], UDP-glucuronosyltransferases [11], sulfotransferases [12] and transporters including organic anion transporter 2 [13], organic cation transporter 1 [14], the ABC transporter ABCC2 [15], ABCC6 [16], ABCG5 [17] and ABCG8 [17]. Although there is clear evidence for HNF4 α to be of key importance in the control of drug metabolism it may also play a role in the regulation of transporters in the choroid plexus. Here we report our efforts in mapping HNF4 α to human and rat choroid plexus. We determined HNF4 α DNA binding activity and searched for transcript expression of various ABCB and ABCC transporters in the human choroid plexus. Apart from qRT-PCR and immunohistochemistry studies we evidence ABCC1 gene expression to be highly dependent on HNF4 α as determined in functional knock down studies. Overall, we provide evidence for HNF4 α to be an important regulator of ABC drug transporters in the choroid plexus and thus may impact efficacy of pharmacotherapy targeted to the brain.

Results

Initially, we searched for HNF4 α transcripts in individual samples of human and rat choroid plexus and confirmed gene expression of HNF4 α by quantitative real time RT-PCR (Figures 1A). We found HNF4 α transcript expression in human and rat choroid plexus to account for approximately a tenth of its expression in the liver (Figures 1A). It is of considerable importance that HNF4 α expression in the human and rat choroid plexus is restricted to P1 promoter driven isoforms (Table 1). Furthermore, we studied expression of the insulin-like growth factor 2 (IGF2), tran-

Table 1: HNF4 α isoform expression in human and rat choroid plexus.

Patient	HNF4 α P1	HNF4 α P2
Patient 100-04	0.0342	0
patient 44-04	0.0025	0
patient 62-04	0.4110	0
Animal	HNF4 α P1	HNF4 α P2
rat 1	0.097	0
rat 2	0.348	0
rat 3	0.158	0

HNF4 α P1 and P2 gene expression was measured in human and rat choroid plexus (n = 3, respectively) by real-time quantitative RT-PCR and computed relative to expression of mitATPase6, which served as a housekeeping gene. Patient characteristics are given in Table 5. Caco-2 cells and pancreatic tissue served as controls for HNF4 α P2 expression in human and rats, respectively.

thyretin (*TTR*) and the transcription factor *FOXJ1* to further qualify choroidal epithelial cells of the brain [18,19]. These transcripts are specifically enriched in choroid plexus. We observed abundant expression of *IGF2*, *TTR* and *FOXJ1* in human choroid plexus as compared to total brain RNA extracts (Figures 1B). There is the need to study histological well qualified tissue, as studies with total brain RNA extracts would render findings meaningless as will be discussed later on. Unfortunately, sufficient human choroid plexus tissue suitable for the harvest of nuclear protein and to perform western blotting as well as EMSA assay could not be obtained. We nonetheless demonstrate HNF4 α protein expression by immunohistochemistry by use of a specific HNF4 α antibody for human (Figures 2A) and rat choroid plexus (Figures 2C). To confirm specificity an excess of antigen preabsorbed to the antibody was used (Figures 2B, D).

We then analyzed expression of different members of the ABCB (MDR subfamily, multidrug resistance)/ABCC (MRP subfamily, multidrug resistance related) gene families in the human choroid plexus by quantitative real time RT-PCR and report results for n = 3 individual human choroid plexus samples. mRNA expression of ABCB4 (MDR2/3), ABCC1 (MRP1), ABCC2 (MRP2), ABCC3 (MRP3), ABCC4 (MRP4), ABCC5 (MRP5) and ABCC6 (MRP6) was comparable to its expression level in commercially available control total human brain RNA extracts (Figures 1C). mRNA expression of ABCB1 (MDR1) determined in choroidal epithelium was lower than in human liver (n = 4) and in brain (Figures 1C).

We then searched for HNF4 α binding sites in proximal promoter sequences (up to -7000 bp) of drug transporter coding genes. For this purpose, we used two different bioinformatic approaches (see Material and Methods section




Figure 1
Gene expression of HNF4 α and different ABC transporters in the choroid plexus. Gene expression of HNF4 α and different ABC transporters in the choroid plexus. A: HNF4 α gene expression in human (n = 3) and rat choroid plexus (n = 3) was measured by quantitative real-time RT-PCR and determined relative to expression of mitATPase6, which served as a housekeeping gene. Expression levels were compared to liver and total brain. Mean values + SD are shown. B and C: Gene expression of FOXJ1, IGF2 and TTR (B) and of different members of the ABCB and ABCC family (C) was analyzed in human choroid plexus by quantitative real-time RT-PCR and determined relative to expression of mitATPase6, which served as a housekeeping gene. Expression levels were compared to liver (n = 4) and total brain. Mean values + SD are shown.

for a description of the employed algorithms). We observed three binding sites within the *ABCB4* promoter, spaced approximately by 600 bp and 1600 bp and two recognition sites within the *ABCC1* promoter (Table 2). Predicted binding sites were confirmed by EMSA band shift assays. We used ³²P-labeled double-stranded DNA probes to specifically probe for HNF4 α -sites located in the human *ABCB4* (hABCB4_1, hABCB4_2 and hABCB4_3) and in the human *ABCC1* (hABCC1_1, hABCC1_2) gene. Note, DNA binding of nuclear extracts to the A-site of the *HNF1 α* -promoter (HNF1pro) served as a positive control.

This site is an established recognition site for HNF4 α . Unfortunately, sufficient amount of human choroid plexus suitable for the isolation of nuclear protein could not be obtained. Instead, we used nuclear extracts isolated from the human intestinal cell line Caco-2 which expresses several ABC transporter genes [20] and therefore is a rich source of HNF4 α nuclear protein. Indeed, HNF4 α protein expression in Caco-2 cells is comparable to organs such as the liver [21]. As depicted in Figures 3A we observed strong binding of HNF4 α to the A-site of the *HNF1 α* -promoter. We also observed strong binding of




Figure 2
Immunohistochemical detection of HNF4α in the choroid plexus. Slices of human (A, B) and rat (C, D) choroid plexus probes were stained with polyclonal antibodies against HNF4α. Arrows indicate representative HNF4α positive cells (A, C). To confirm specificity of the immunohistochemical localization antibodies were preabsorbed with excess of antigens for HNF4α (B, D). Patient identification numbers were indicated respectively and patient characteristics are given in Table 5. Magnification ×400.

HNF4α to the predicted sites in the promoters of *ABCB4* and *ABCC1* (Figures 3A); bands could be shifted with a specific HNF4α antibody therefore demonstrating selectivity and specificity of the assay. Alignment of human, rat and mouse *ABCB4* and *ABCC1* genes did not identify common HNF4α binding sites. This points to differences in the molecular organization of ABC promoters in orthologous genes. HNF4α binding sites for the rat and mouse *ABCB4* and *ABCC1* genes are given in Table 2,

whereas the sequences of oligonucleotides to confirm the predicted sites experimentally are shown in Table 3. As shown in Figures 3B and 3C EMSA band shift assays confirmed binding of HNF4α to rat and mouse *ABCB4* and *ABCC1* targeted sequences.

To further probe for the role of HNF4α in ABC gene regulation we employed an siRNA approach. Specifically, siRNA-mediated functional knock down of HNF4α in the

Table 2: Predicted binding sites for HNF4α in ABC genes.

Gene name	Accession number	bp relative to transcription start site	Score core/matrix	
Human				
ABCB4	NM_000443.3	-6274 to -6252	V\$zemlin13_11045	1.000/0.896
		-4680 to -4658	V\$HNF4_01	1.000/0.990
		-4063 to -4041	V\$zemlin13_1104	1.000/0.997
			V\$HNF4_01	1.000/0.954
ABCC1	NM_004996.3	-1756 to -1734	V\$zemlin13_11045	1.000/0.958
		-5266 to -5244	V\$HNF4_01	0.915/0.925
			V\$HNF4_01	1.000/0.944
			V\$zemlin13_11045	1.000/0.969
Rat				
ABCB4	NM_012690.1	-4584 to -4562	V\$zemlin13_11045	1.000/0.894
		-39 to -17	V\$HNF4_01	1.000/0.943
ABCC1	NM_022281.2	-2146 to -2124	V\$zemlin13_11045	1.000/0.978
			V\$HNF4_01	1.000/0.908
Mouse				
ABCB4	NM_008830.2	-5281 to -5259	V\$zemlin13_11045	1.000/0.907
ABCC1	NM_008576.2	+2626 to +2648 (Intron 1)	V\$HNF4_01	1.000/0.924
			V\$zemlin13_11045	1.000/0.910

Transfac matrix (V\$HNF4_01) or self-generated matrix (V\$zemlin13_11045) with a setting of core and matrix similarity 0.9 was used.

human Caco-2 cell line resulted in significantly decreased gene expression of ABCC1 (Table 4; for transfection efficiency and stable expression of mitATPase6 after transfection see Niehof and Borlak, 2008 [22]). These results confirm ABCC1 to be a gene target of HNF4α. ABCB4 expression in Caco-2 cells is near the limit of detection. Consequently, knockdown experiments are not meaningful.

Discussion

Our study aimed for a better understanding of the role of HNF4α in the regulation of drug transporters. Here we present evidence for expression of HNF4α in the epithelium of the CSF barrier. By applying position weight matrices to genomic sequences of ABC transporters we were able to predict HNF4α binding sites in the promoters of the ABCB4 and ABCC1 gene. The predicted binding sites were then confirmed by EMSA band shift assays. We propose a role for HNF4α in the regulation of drug trans-

Table 3: Shift-probes sequences.

Gene	Oligo name	Sequence
Human		
HNF1α	HNF1pro	AGGGCTGA AGTCCA A AGTTCA GTCCCTTC
ABCB4	hABCB4_1 (GS49)	AGAA AGGTCA A AGGCAA AGCA
	hABCB4_2 (GS50)	GTGC AGGGCA A AGGTCA GATT
	hABCB4_3 (GS51)	GCTA AGTCCA A AGATCA ATTCC
ABCC1	hABCC1_1 (GS48)	TTGAC AGTATA A AGGTCA AGCAA
	hABCC1_2 (GS171)	TTCAT AGAGCA A AGTACA GCAAC
Rat		
ABCB4	rABCB4_1 (GS166)	AGAAG CCCTCA A AGTCCA CCTAT
	rABCB4_2 (GS167)	CGCAG CGAGCA A AGTCCA GGTCT
ABCC1	rABCC1_1 (GS169)	GACAC GGGCCA A AGACCC ACAGA
Mouse		
ABCB4	mABCB4_1 (GS168)	ATAGC CATGCA A AGTCCA AAAGA
ABCC1	mABCC1_1 (GS170)	TCCCT TGTGCA A AGGTCT AGGAT

The central five nucleotides of the HNF4α binding site are highlighted in bold letters. The two direct repeats (DR1) separated by a spacer of one nucleotide are underlined.




Figure 3
HNF4α binding to promoters of ABCB4 and ABCC1. Electrophoretic mobility shift assays with 2,5 µg Caco-2 cell nuclear extract and oligonucleotides corresponding to the A-site of the HNF1α promoter (HNF1pro) and to putative HNF4α binding-sites within human (A), rat (B) and mouse (C) ABCB4 and ABCC1 as ³²P labeled probe. In supershift assays an antibody directed against HNF4α (+) was added. For oligonucleotide sequence information see Table 2 and 3.

Table 4: Functional knock down of HNF4α in cultures of human Caco-2 cells.

Gene	Gene expression 48 h after HNF4α siRNA transfection [%]	p-value*
HNF4α	24.0	0.0031
ABCC1	56.4	0.0478

Functional knock down of HNF4α was already reported [22]. * A student's t test was done to compare HNF4α siRNA transfection against negative siRNA transfected controls. The results were considered significant when the p value was less than 0.05. Gene expression was determined by real-time quantitative RT-PCR in triplicates; mitATPase6 served as housekeeping gene. The $\Delta\Delta C_T$ -method was used to determine repression of gene expression. Values for HNF4α siRNA transfection represent transcript abundance relative to negative siRNA transfected controls, which were set to 100%.

porters of the choroid plexus. Notably, HNF4α is a key player in the regulation of genes coding for various metabolic pathways and of xenobiotic metabolism [6-9,23,24]. This protein also promotes expression of an epithelial phenotype [25-27]. Specifically, epithelium of the choroid plexus cells is highly differentiated and functions as a blood-CSF barrier. The expression of HNF4α in the epithelium of the choroid plexus and its DNA binding to regulatory sequences of drug transporters is a novel finding. Its expression accounts for approximately a tenth of HNF4α expression found in liver. Furthermore, there is evidence for HNF4α to undergo alternative splicing with isoforms may arising from alternative splicing and/or usage of two different promoters [6]. Specifically, the P1 promoter generates six different isoforms (HNF4α1-α6); but activation of the P2 promoter results in isoforms HNF4α7-α9. P2 promoter-driven HNF4α isoforms are expressed throughout mouse liver development, but dis-

appear after birth, while P1 promoter-driven transcripts are abundantly expressed postnatally [28]. Additionally, P2 isoforms are induced in mouse and human hepatocellular carcinoma [29,30] and are primarily expressed in human pancreatic islets and exocrine cells [31]. In the case of the choroid plexus it appears that expression of HNF4α is restricted to P1 promoter-driven isoforms. HNF4α protein expression in human and rat choroid plexus could be clearly evidenced by immunohistochemistry. To further qualify tissue preparations of choroid plexus expression of *IGF2*, *TTR* and *FOXJ1* was investigated; their expression are accepted genetic markers of the choroid plexus [18,19]. We validated tissue preparations by morphological and genetic markers. Unlike previous studies where expression of *HNF4α* transcripts could not be evidenced in total RNA extracts of the brain [6], we were able to confirm expression of *HNF4α* in the choroid plexus of human and rat brain. Most certainly, total brain RNA extracts

Table 5: Patient characteristics.

Liver					
Patient Identification	Sex	Age	Information	Tissue	Application
P29	F	40	Colorectal liver metastasis	Healthy tissue from liver resection	Gene expression analysis, comparison to choroid plexus
P2	F	70	Hepatocellular carcinoma	Healthy tissue from liver resection	Gene expression analysis, comparison to choroid plexus
P3	F	57	Hepatocellular carcinoma	Healthy tissue from liver resection	Gene expression analysis, comparison to choroid plexus
P4	M	67	Hepatocellular carcinoma	Healthy tissue from liver resection	Gene expression analysis, comparison to choroid plexus
Choroid Plexus					
Patient Identification	Sex	Age	Cause of death	Tissue	Application
44-04	F	63	liver cirrhosis	38 h post mortem	Gene expression analysis
62-04	F	79	ovarial carcinoma	33 h post mortem	Gene expression analysis
100-04	M	74	Acute myocardial infarction	48 h post mortem	Gene expression analysis
H5	M	53	Circulatory failure, liver cirrhosis	Choroid plexus	Paraffin-embedded slices for immunohistochemistry
H40	M	32	Circulatory failure, endocarditis	Choroid plexus	Paraffin-embedded slices for immunohistochemistry
H54	M	59	Respiratory insufficiency, pneumonia	Choroid plexus	Paraffin-embedded slices for immunohistochemistry
H55	M	81	Circulatory failure, generalized arteriosclerosis	Choroid plexus	Paraffin-embedded slices for immunohistochemistry

Table 6: Real-time PCR primer sequences and amplification settings.

Gene	Accession number	Species	Primer sequence	Fragment length	Annealing	Extension	Fluorescence	Slope
HNF4 α	NM_000457	human	fwd:CTGCTCGGAGCCACCAAGAGATCCATG rev: ATCATCTGCCAGGTGATGCTCTGCA	371 bp	60°C	15 sec	88°C	-3.629
HNF4 α P1	NM_178849 NM_178850 NM_000457	human	fwd: GAATGCGACTCTCCAAAACC rev: GGCACTGGTTCTCTTGTCT	339 bp	68°C	14 sec	88°C	-3.058
HNF4 α P2	NM_001030003	human	fwd: GGGCTCCAGTGGAGAGTTC rev: CATAGCTTGACCTTCGAGTGC	342 bp	68°C	14 sec	88°C	-2.648
FOXJ1	NM_001454	human	fwd: CTACTIONGATGCCACGCTCA rev: CGAGGCACTTTGATGAAGC	183 bp	60°C	8 sec	86°C	-3.459
IGF2	NM_000162	human	fwd: GGTGCTTCTCACCTTCTTGG rev: GGGGTATCTGGGGAAGTTGT	298 bp	68°C	12 sec	90°C	-2.724
TTR	NM_000371	human	fwd: CAGAAAGGCTGCTGATGACA rev: GCCGTGGTGAATAGGAGTA	261 bp	60°C	11 sec	86°C	-3.438
ABCB1	NM_000927	human	fwd: AAAAAGATCAACTCGTAGGAGTG rev: GCACAAAATACACCAACAA	161 bp	60°C	7 sec	81°C	-3.331
ABCB4	NM_000443	human	fwd: CCACAGCGAACTGATGAAGA rev: CACGACAAAGTAGGGCCATT	301 bp	68°C	13 sec	83°C	-3.826
ABCC1	NM_004996	human	fwd: AGTCCCCACAGAAGGAGTGG rev: TCCCCGACCGTGGAGGATTT	358 bp	60°C	15 sec	87°C	-3.639
ABCC2	NM_000392	human	fwd: CCGTATCAGGTTTGCCAGTT rev: CAACAGCCACAATGTTGGTC	312 bp	60°C	13 sec	84°C	-3.756

Table 6: Real-time PCR primer sequences and amplification settings. (Continued)

ABCC3	NM_003786	human	fwd: CTCAATGTGGCAGACATCGG rev: GGGAGCTCACAACGTGTGC	177 bp	60°C	8 sec	87°C	-3.731
ABCC4	NM_005845 NM_001105515	human	fwd: CTTGGAGAGGAGTTGCAAGG rev: GCTGTGTTCAAAGCCACAGA	236 bp	68°C	10 sec	78°C	-4.479
ABCC5	NM_005688	human	fwd: CCTGTTTGGGAAGGAATATGA rev: CCTGTTTGGGAAGGAATATGA	203 bp	60°C	9 sec	88°C	-3.446
ABCC6	NM_001171	human	fwd: TTCTCTGTGTCGCTGGTGTC rev: GGCACTGTGTATGGTGATGC	309 bp	60°C	13 sec	87°C	-2.982
MitATPase6	NC_001807	human	fwd: CTAAAGGACGAACCTGA rev: TGGCCTGCAGTAATGTT	315 bp	55°C	13 sec	83°C	-3.608
HNF4 α	NM_022180	rat	fwd: GCCTGCCTCAAAGCCATCAT rev: GACCCTCCAAGCAGCATCTC	274 bp	55°C	11 sec	88°C	-3.697
HNF4 α P1	DI0554, EF193392	rat	fwd: AAATGTGCAGGTGTTGACCA rev: CACGCTCCTCCTGAAGAATC	178 bp	60°C	7 sec	87°C	-3.296
HNF4 α P2	AF329936 EF193390	rat	fwd: CTCCAGTGGCGAGTCCTTAT rev: TCACGCTCCTCCTGAAGAAT	171 bp	60°C	7 sec	87°C	-2.273
MitATPase6	NC_001665	rat	fwd: CTAAAGGACGAACCTGA rev: TGGCCTGCAGTAATGTT	315 bp	55°C	13 sec	83°C	-3.644

dilute copy number of *HNF4 α* transcripts to presumable levels below the limit of detection. Here, we evidence binding of *HNF4 α* to regulatory sequences of drug transporters expressed in the choroid plexus and analyzed gene expression of ABC transporters in patients with different causes of death, but the functional significance of the newly identified *HNF4 α* binding sites in activating the *ABCB4* and *ABCC1* promoters still needs to be established.

In the past attempts to detect *HNF4 α* DNA binding in a choroid plexus papilloma failed [32]. The investigators performed EMSA experiments with nuclear extracts of rat liver, kidney and intestine and of SV40-induced choroid plexus papilloma of transgenic mice, but unfortunately probed for *HNF4 α* binding with an oligonucleotide corresponding to the *HNF4 α* binding site in the mouse *TTR* (transthyretin) promoter. The authors did not employ an antibody in EMSA band shift assays; instead, competition with excess of unlabeled probes was done. Although the authors described a weak binding of *HNF4 α* with nuclear extracts from kidney they considered intestine as well as choroid plexus as deficient for *HNF4 α* binding. There is a need to consider tissue specific DNA binding activity. *HNF4 α* binding at the *TTR* promoter is much less as compared to the A-site in the *HNF1* promoter (not exceeding a tenth, data not shown). As detailed above, *HNF4 α* gene expression in human and rat choroid plexus is approximately one tenth of its expression in the liver (see Figures 1A). It is therefore not surprising that previous investigators [32] failed to detect *HNF4 α* protein in intestine and choroid plexus, even though the authors described weak expression of this protein in kidney. By now, it is well established that *HNF4 α* expression is not restricted to liver, but also functions in kidney and intestine [33,34]. Here we evidence by immunohistochemical staining *HNF4 α* to be expressed in epithelium of human and rat choroid plexus (see Figures 2).

Notably, the choroid plexus functions as a barrier for drug uptake to the brain. This tissue expresses drug metabolizing enzymes (DMEs) and some transporters [1]. Expression of DMEs in the choroid plexus is part of a defense program to prevent entry of xenobiotics into the brain. The blood-CSF barrier also regulates entry and distribution of various pharmacologically active compounds between the blood and the CSF interface and is basically involved in numerous exchange processes thereby determining the supply of the brain with nutrients and hormones [1]. Indeed, endogenous metabolites, as well as neurotransmitter and metabolites from the brain are cleared via this barrier [19]. In drug therapies efflux transporters of the ABC-family are of pivotal importance in determining therapeutic tissue levels. In the past research focused on their regulation in liver, kidney and intestine [3,4]. The knowledge on drug transporter in specialized

tissues of the brain is incomplete. Specifically, *ABCB* (*MDR*) proteins accept a broad range of substrates and may transport large lipophilic, neutral or cationic compounds. This includes a vast number of neuropharmacological drugs such as antiepileptic and antiviral drugs, antidepressants, opioids, antipsychotics and tranquilizer [35-40]. *ABCB1*, a transporter with highest expression in the gastrointestinal tract [4,41], is expressed at low levels in human choroid plexus, that is much lower than in liver and total brain. Our findings are consistent with results reported previously for rats [42]. Its apical expression in rat, mouse and human choroidal epithelium was shown previously [43]. Furthermore, *ABCB4* is highly expressed in the liver, where it is acting as a "flippase" in transporting phospholipids into the bile, but *ABCB4* can also bind and transport a subset of *ABCB1* substrates with an overlap in substrate specificity [44]. As reported for *ABCB1*, *ABCB4* is expressed at low levels in human choroid plexus, i.e. more than 500 fold lower than in liver. Low *ABCB4* expression was also evidenced in choroidal epithelium of the rat [42], but to the best of our knowledge our study is the first report on expression in human choroid plexus. An apical distribution of *ABCB1* in neonatal cultured rat choroid plexus cells imply a drug transport from blood into CSF [43]. In endothelial cells of small blood capillaries of the blood-brain barrier, apical located *ABCB1* pumps drugs back into the blood stream and therefore limits drug penetration to the brain [2].

Likewise, the *ABCC* (*MRP*) proteins are multispecific organic anion transporters and accept glucurono-, glutathione- and sulfo-conjugates. They transport physiological substrate conjugates as well as drug conjugates. Expression of *ABCC1-6* mRNA transcripts was reported for rat choroid plexus [42] as was expression of *ABCC1* for human choroid plexus [43,45,46]. *ABCC1*, *ABCC4* and *ABCC5* were expressed in human choroid plexus at least at 10 fold higher than in liver; whereas *ABCC2*, *ABCC3* and *ABCC6* were expressed up to 800 fold lower than liver. Similar results were reported for *ABCC* transporter expression in rat choroid plexus [42]. Notably, *ABCC1* is localized at the basolateral membrane of choroid plexus [43,45,46], but Gazzin et al [46] described a major difference in the localization of *ABCB1* and *ABCC1* proteins between the blood-brain and the blood-CSF barrier with strongest expression of *ABCC1* at the choroidal epithelium. Indeed, *ABCC* proteins contribute to the protective role of the choroid plexus and mediate basolateral efflux of conjugates resulting from choroidal drug metabolism into the blood. Although it is known that the choroid plexus is important in regulating the distribution of various pharmacologically active compounds between the blood and the CSF, the characterization of the involved human ABC transporters gives new insights into the function of the CSF barrier.

Furthermore, ABCBs (MDRs) and ABCCs (MRPs) are inducible transporters and are highly responsive to chemotherapeutics, carcinogens, inflammation, heat shock, hypoxia and irradiation [47]. They are regulated by a complex network of transcriptional cascades, such as by multiple ligand activated nuclear receptors like retinoid X receptor (RXR), farnesoid X receptor (FXR), constitutive androstane receptor (CAR) and the xenobiotic receptor pregnane X receptor (PXR) [47,48]. There is also evidence for the transcription factors AP-1, p53, Egr-1 and WT-1 to participate in their regulation with NF- κ B, Sp1 and Sp3 being involved in the constitutive expression [47]. Recently, an upregulation of *ABCB1*, *ABCB4* and *ABCC4* transcripts was reported in human embryonic kidney cells that conditionally expressed wild-type HNF4 α [33]. An important role of HNF4 α in the transcriptional control of drug transporters was reported for human hepatocytes as determined by adenoviral HNF4 α -siRNA mediated knockdown [49]. We also employed an siRNA mediated functional knockdown of HNF4 α and found *ABCC1* gene expression to be massively repressed. There is a need to improve an understanding of the mechanism by which transporters are regulated. This will impact the design of novel CNS therapeutics. Targeting transporters may thus be useful in achieving therapeutic tissue levels of CNS drugs.

Conclusion

We report expression of HNF4 α in choroid plexus of the human and rat brain. This factor might regulate expression of some ATP binding cassette transporters. Targeting of HNF4 α may impact efficacy of pharmacotherapy of CNS drugs.

Methods

Human and rat tissue

A total of $n = 7$ human and $n = 7$ rat tissues were analyzed. Samples of human choroid plexus ($n = 3$, gene expression analysis) were kindly provided by T. Arendt (Department of Neuroanatomy, Paul-Flechsig-Institute, University of Leipzig, Germany). Paraffin-embedded slices of human choroid plexus for immunohistochemistry ($n = 4$) were kindly provided by C. Grothe (Institute of Neuroanatomy, Hannover Medical School, Hannover, Germany). Human liver tissue (gene expression analysis) was obtained from patients undergoing hepatic resections and were kindly provided by J. Klempnauer (Department of Visceral and Transplantation Surgery, Hannover Medical School, Hannover, Germany). Patient characteristics are given in Table 5. Control human brain RNA was purchased from BD Biosciences (Heidelberg, Germany).

Samples of rat choroid plexus ($n = 3$, Sprague Dawley rats, gene expression analysis) were kindly provided by H. Hilbig and K. Spanel-Borowski (Department of Anatomy,

University of Leipzig, Germany). Samples of rat liver and brain (Sprague Dawley rats, gene expression analysis) were generated in-house. Paraffin-embedded slices of rat brain (Sprague Dawley rats) containing choroid plexus regions for immunohistochemistry ($n = 4$) were kindly provided by C. Grothe (Institute of Neuroanatomy, Hannover Medical School, Hannover, Germany).

Quantitative real-time RT-PCR

Analysis of human samples: Three human choroid plexus samples and four human liver samples were analyzed separately and used for calculation of the mean and standard deviation. Analysis of rat samples: Three rat choroid plexus samples, three rat liver and three rat brain samples were analyzed separately and used for calculation of the mean and standard deviation. Total RNA from choroid plexus and liver was isolated using the RNeasy Mini Kit (Qiagen, Hilden, Germany) according to the manufacturers recommendations. Subsequently to RNA isolation, a DNase I digest was performed. 4 μ g total RNA from each sample was used for reverse transcription (Omniscript Reverse Transcriptase, Qiagen, Hilden, Germany). Quantitative real-time RT-PCR measurement was done with the Lightcycler (Roche Diagnostics, Mannheim, Germany) with the following conditions: denaturation at 95°C, annealing at different temperatures for 8 sec, extension at 72°C for different times and detection of SYBR-Green I-fluorescence at different temperatures. Detailed primer specific conditions and oligonucleotide sequence information are given in Table 6. Relative quantification was performed using the "Fit Points Method" of the LightCycler3 Data Analysis Software version 3.5.28 (Roche Diagnostics, Mannheim, Germany) by comparing the sample values to a standard curve within the linear range of amplification. This comparison was performed during each LightCycler Run (for genes of interest as well as for the housekeeping gene, i.e. *mitATPase6*). The standardized sample values for each gene of interest were divided by the standardized values of the housekeeping gene. The slope of external standard curves are given in Table 6, indicating the PCR efficiency for each amplicon.

Caco-2 cell culture

Caco-2 cells are a valuable source for HNF4 α nuclear protein [21] and were obtained from and cultivated as recommended by DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany) and seeded with a density of 4×10^6 cells per 75 cm² flask and harvested after 11 days of culture.

Isolation of nuclear extracts

Nuclear extracts from Caco-2 cells were isolated by the modified method of Dignam et al [50]. Eleven days after seeding cells were washed twice with ice-cold PBS, scraped into microcentrifuge tubes and centrifuged for 5 min at

2000 × g, 4°C. Cell pellets were resuspended in lysis buffer (10 mM Tris pH 7.4, 2 mM MgCl₂, 140 mM NaCl, 1 mM DTT, 4 mM Pefabloc, 1% Aprotinin, 40 mM β-glycerophosphate, 1 mM sodiumorthovanadate and 0.5% TX100) at 4°C for 10 min (300 μl for 1 × 10⁷ cells), transferred onto one volume of 50% sucrose in lysis buffer and centrifuged at 14000 × g and 4°C for 10 min. Nuclei were resuspended in Dignam C buffer (20 mM Hepes pH 7.9, 25% glycerol, 420 mM NaCl, 1.5 mM MgCl₂, 0.2 mM EDTA, 1 mM DTT, 4 mM Pefabloc, 1% Aprotinin, 40 mM β-glycerophosphate, 1 mM sodiumorthovanadate, 30 μl for 1 × 10⁷ cells) and gently shaken at 4°C for 30 min. Nuclear debris was removed by centrifugation at 14000 × g at 4°C for 10 min. Protein concentrations were determined according to the method of Smith *et al* [51]. The extracts were aliquoted and stored at -70°C.

Electrophoretic mobility shift assays

Forward and reverse oligonucleotides (for sequence information see Table 3) were purchased from MWG Biotech (Ebersberg/Muenchen, Germany), annealed and ³²P-labeled using ³²PγATP and T4-kinase (New England Biolabs, Frankfurt, Germany). 2.5 μg Caco-2 cell nuclear extract and 10⁵ cpm (0.027 ng) radiolabeled probe were incubated in binding buffer consisted of 25 mM HEPES, pH 7.6, 5 mM MgCl₂, 34 mM KCl, 2 mM DTT, 2 mM Pefabloc, 2% Aprotinin, 40 ng of poly (dI-dC)/μl and 100 ng of bovine serum albumin/μl for 20 minutes on ice. Free DNA and DNA-protein complexes were resolved on a 6% polyacrylamide gel (acrylamide: bisacrylamide ratio = 37.5:1). Super shift experiments were done with a 1 μg HNF4α specific antibody (sc-6556x, Santa Cruz Biotechnology, Heidelberg, Germany). DNA binding of nuclear extracts to the A-site of the HNF1α-promoter (HNF1pro) served as a positive control.

Bioinformatic searches for HNF4α binding-sites

The transcription start site (TSS, +1) of the NCBI mRNA reference sequence (RefSeq) was aligned using the UCSC Genome Browser <http://genome.ucsc.edu/> for promoter annotation of the respective genes. Proximal promoters (up to -7000 bp) of human ABCB1, ABCB4 and ABCC1 to ABCC6 were checked for putative HNF4α binding-sites with the tool MATCH [52], <http://www.biobase.de> by employing two different weight matrix, i.e. V\$HNF4_01, Transfac matrix (generated by Biobase) and V\$zemplin13-11045 (self-generated). V\$zemplin13-11045 is based on a collection of 33 well known HNF4α sites and was generated using the matrix generation tool of Biobase <http://www.biobase.de>. At least one of both matrices has to exceed the cut-off of 0.9 for core similarity and 0.9 for matrix similarity (Table 2). Furthermore, proximal promoters (up to -7000 bp) of rat and mouse ABCB4 and ABCC1 were checked for putative HNF4α binding sites (Table 2).

Immunohistochemistry

The sections were deparaffinized, demasked by heating, incubated with 0,6% H₂O₂ in methanol for 30 min, and subsequently with protein block serum-free reagent (Dako, Glostrup, Denmark) for 10 min. Incubation with polyclonal antibody (Santa Cruz Biotechnology, Heidelberg, Germany) against HNF4α (sc-6556x, 1:400 dilution for rat sections, 1:200 dilution for human sections) was performed for 45 min. The sections were rinsed with Tris-buffered saline, incubated with biotinylated universal secondary antibodies (Dako, Glostrup, Denmark) for 15 min and subsequently with horseradish peroxidase-conjugated streptavidin solution (Dako, Glostrup, Denmark) for 15 min. Labeling was detected using a diaminobenzidine (DAB) chromogen solution (Dako, Glostrup, Denmark) for 5 min. The sections were counterstained with hematoxylin before examination under light microscope. To confirm the specificity of the immunohistochemical localization, antibodies preabsorbed 2 h with a twenty fold excess of antigen for HNF4α (sc-6556P, Santa Cruz Biotechnology, Heidelberg, Germany) were used.

siRNA silencing of HNF4α

Human HNF4α-specific siRNA probes were purchased from Qiagen (Hilden, Germany). Caco-2 cells (1,5 × 10⁵ cells/well in 24-well plate) were transfected in triplicate for 48 h with 25 nM of the siRNA duplex using HiPerFect transfection reagent (Qiagen, Hilden, Germany). Alexa-Fluor488 labeled siRNA (Qiagen, Hilden Germany) was used as negative siRNA and as positive control for transfection efficiency. For transfection efficiency and stable expression of mitATPase6 after transfection see Niehof and Borlak, 2008 [22].

Abbreviations

ABC transporter: ATP-binding cassette transporter; CSF: cerebrospinal fluid; DME: drug metabolizing enzymes; EMSA: electrophoretic mobility shift assay; HNF: hepatocyte nuclear factor; MDR: multidrug resistance gene family (ABCB); MRP: multidrug resistance related proteins gene family (ABCC); TTR: transthyretin.

Authors' contributions

JB designed the entire study, supervised the experimental works and is responsible for the final writing of the manuscript, MN supervised the experiments and prepared the initial draft of the manuscript. Both authors read and approved the final manuscript.

Acknowledgements

We thank S. Marschke, A. Pfanne and A. Schulmeyer for valuable technical assistance, S. Reymann and R. Zemplin for assistance in bioinformatics and advice on design of PCR primers, T. Arendt, U. Gärtner and C. Grothe for providing samples of human choroid plexus and H. Hilbig, K. Spanel-Borowski and C. Grothe for the preparation of rat choroid plexus. This

work was supported by a grant of the Lower Saxony Ministry of Culture and Science to J.B.

References

- Gherzi-Egea JF, Strazielle N: **Brain drug delivery, drug metabolism, and multidrug resistance at the choroid plexus.** *Microsc Res Tech* 2001, **52**:83-88.
- Graff CL, Pollack GM: **Drug transport at the blood-brain barrier and the choroid plexus.** *Curr Drug Metab* 2004, **5**:95-108.
- Schinkel AH, Jonker JW: **Mammalian drug efflux transporters of the ATP binding cassette (ABC) family: an overview.** *Adv Drug Deliv Rev* 2003, **55**:3-29.
- Glavinas H, Krajcsi P, Cserepes J, Sarkadi B: **The role of ABC transporters in drug resistance, metabolism and toxicity.** *Curr Drug Deliv* 2004, **1**:27-42.
- Toyoda Y, Hagiya Y, Adachi T, Hoshijima K, Kuo MT, Ishikawa T: **MRP class of human ATP binding cassette (ABC) transporters: historical background and new research directions.** *Xenobiotica* 2008, **38**:833-862.
- Sladek FM, Seidel SD: **Hepatocyte nuclear factor 4alpha.** In *Nuclear Receptors and Disease* Edited by: Burris T, McCabe ERB. Academic Press, London; 2001:309-361.
- Schrem H, Klemmner J, Borlak J: **Liver-enriched transcription factors in liver function and development. Part I: the hepatocyte nuclear factor network and liver-specific gene expression.** *Pharmacol Rev* 2002, **54**:129-158.
- Parviz F, Matullo C, Garrison WD, Savatski L, Adamson JW, Ning G, et al.: **Hepatocyte nuclear factor 4alpha controls the development of a hepatic epithelium and liver morphogenesis.** *Nat Genet* 2003, **34**:292-296.
- Jover R, Bort R, Gomez-Lechon MJ, Castell JV: **Cytochrome P450 regulation by hepatocyte nuclear factor 4 in human hepatocytes: a study using adenovirus-mediated antisense targeting.** *Hepatology* 2001, **33**:668-675.
- Gonzalez FJ: **Regulation of hepatocyte nuclear factor 4 alpha-mediated transcription.** *Drug Metab Pharmacokinet* 2008, **23**:2-7.
- Barbier O, Girard H, Inoue Y, Duez H, Villeneuve L, Kamiya A, et al.: **Hepatic expression of the UGT1A9 gene is governed by hepatocyte nuclear factor 4alpha.** *Mol Pharmacol* 2005, **67**:241-249.
- Echchgadda I, Song CS, Oh T, Ahmed M, De LCI, Chatterjee B: **The xenobiotic-sensing nuclear receptors pregnane X receptor, constitutive androstane receptor, and orphan nuclear receptor hepatocyte nuclear factor 4alpha in the regulation of human steroid/bile acid-sulfotransferase.** *Mol Endocrinol* 2007, **21**:2099-2111.
- Popowski K, Eloranta JJ, Saborowski M, Fried M, Meier PJ, Kullak-Ublick GA: **The human organic anion transporter 2 gene is transactivated by hepatocyte nuclear factor-4{alpha} and suppressed by bile acids.** *Mol Pharmacol* 2005, **67**:1629-1638.
- Saborowski M, Kullak-Ublick GA, Eloranta JJ: **The Human Organic Cation Transporter 1 Gene is Transactivated by Hepatocyte Nuclear Factor-4{alpha}.** *J Pharmacol Exp Ther* 2006, **317**:778-785.
- Qadri I, Hu LJ, Iwahashi M, Al Zuabi S, Quattrochi LC, Simon FR: **Interaction of hepatocyte nuclear factors in transcriptional regulation of tissue specific hormonal expression of human multidrug resistance-associated protein 2 (abcc2).** *Toxicol Appl Pharmacol* 2009, **234**:281-292.
- Douet V, VanWart CM, Heller MB, Reinhard S, Le Saux O: **HNF4alpha and NF-E2 are key transcriptional regulators of the murine Abcc6 gene expression.** *Biochim Biophys Acta* 2006, **1759**:426-436.
- Sumi K, Tanaka T, Uchida A, Magoori K, Urashima Y, Ohashi R, et al.: **Cooperative interaction between hepatocyte nuclear factor 4 alpha and GATA transcription factors regulates ATP-binding cassette sterol transporters ABCG5 and ABCG8.** *Mol Cell Biol* 2007, **27**:4248-4260.
- Lim L, Zhou H, Costa RH: **The winged helix transcription factor HFH-4 is expressed during choroid plexus epithelial development in the mouse embryo.** *Proc Natl Acad Sci USA* 1997, **94**:3094-3099.
- Strazielle N, Gherzi-Egea JF: **Choroid plexus in the central nervous system: biology and physiopathology.** *J Neuropathol Exp Neurol* 2000, **59**:561-574.
- Borlak J, Zwadlo C: **Expression of drug-metabolizing enzymes, nuclear transcription factors and ABC transporters in Caco-2 cells.** *Xenobiotica* 2003, **33**:927-943.
- Niehof M, Borlak J: **RSK4 and PAK5 Are Novel Candidate Genes in Diabetic Rat Kidney and Brain.** *Mol Pharmacol* 2005, **67**:604-611.
- Niehof M, Borlak J: **HNF4{alpha} and the Ca-Channel TRPC1 Are Novel Disease Candidate Genes in Diabetic Nephropathy.** *Diabetes* 2008, **57**:1069-1077.
- Hayhurst GP, Lee YH, Lambert G, Ward JM, Gonzalez FJ: **Hepatocyte nuclear factor 4alpha (nuclear receptor 2A1) is essential for maintenance of hepatic gene expression and lipid homeostasis.** *Mol Cell Biol* 2001, **21**:1393-1403.
- Odom DT, Zizlsperger N, Gordon DB, Bell GW, Rinaldi NJ, Murray HL, et al.: **Control of Pancreas and Liver Gene Expression by HNF Transcription Factors.** *Science* 2004, **303**:1378-1381.
- Chiba H, Gotoh T, Kojima T, Satohisa S, Kikuchi K, Osanai M, et al.: **Hepatocyte nuclear factor (HNF)-4alpha triggers formation of functional tight junctions and establishment of polarized epithelial morphology in F9 embryonal carcinoma cells.** *Exp Cell Res* 2003, **286**:288-297.
- Chiba H, Itoh T, Satohisa S, Sakai N, Noguchi H, Osanai M, et al.: **Activation of p21(CIP1/WAF1) gene expression and inhibition of cell proliferation by overexpression of hepatocyte nuclear factor-4alpha.** *Exp Cell Res* 2005, **302**:11-21.
- Battle MA, Konopka G, Parviz F, Gaggi AL, Yang C, Sladek FM, et al.: **Hepatocyte nuclear factor 4alpha orchestrates expression of cell adhesion proteins during the epithelial transformation of the developing liver.** *Proc Natl Acad Sci USA* 2006, **103**:8419-8424.
- Torres-Padilla ME, Fougere-Deschattre C, Weiss MC: **Expression of HNF4alpha isoforms in mouse liver development is regulated by sequential promoter usage and constitutive 3' end splicing.** *Mech Dev* 2001, **109**:183-193.
- Niehof M, Borlak J: **EPS15R, TASP1, and PRPF3 are novel disease candidate genes targeted by HNF4alpha splice variants in hepatocellular carcinomas.** *Gastroenterology* 2008, **134**:1191-1202.
- Tanaka T, Jiang S, Hotta H, Takano K, Iwanari H, Sumi K, et al.: **Dysregulated expression of P1 and P2 promoter-driven hepatocyte nuclear factor-4alpha in the pathogenesis of human cancer.** *J Pathol* 2006, **208**:662-672.
- Hansen SK, Parrizas M, Jensen ML, Pruhova S, Ek J, et al.: **Genetic evidence that HNF-1{alpha}-dependent transcriptional control of HNF-4{alpha} is essential for human pancreatic {beta} cell function.** *J Clin Invest* 2002, **110**:827-833.
- Costa RH, Van Dyke TA, Yan C, Kuo F, Darnell JE Jr: **Similarities in transthyretin gene expression and differences in transcription factors: liver and yolk sac compared to choroid plexus.** *Proc Natl Acad Sci USA* 1990, **87**:6589-6593.
- Lucas B, Grigo K, Erdmann S, Lausen J, Klein-Hitpass L, Ryffel GU: **HNF4alpha reduces proliferation of kidney cells and affects genes deregulated in renal cell carcinoma.** *Oncogene* 2005, **24**:6418-6431.
- Garrison WD, Battle MA, Yang C, Kaestner KH, Sladek FM, Duncan SA: **Hepatocyte nuclear factor 4alpha is essential for embryonic development of the mouse colon.** *Gastroenterology* 2006, **130**:1207-1220.
- Loscher W, Potschka H: **Role of multidrug transporters in pharmacoresistance to antiepileptic drugs.** *J Pharmacol Exp Ther* 2002, **301**:7-14.
- Potschka H, Fedrowitz M, Loscher W: **Multidrug resistance protein MRP2 contributes to blood-brain barrier function and restricts antiepileptic drug activity.** *J Pharmacol Exp Ther* 2003, **306**:124-131.
- Dagenais C, Graff CL, Pollack GM: **Variable modulation of opioid brain uptake by P-glycoprotein in mice.** *Biochem Pharmacol* 2004, **67**:269-276.
- El Ela AA, Hartter S, Schmitt U, Hiemke C, Spahn-Langguth H, Langguth P: **Identification of P-glycoprotein substrates and inhibitors among psychoactive compounds—implications for pharmacokinetics of selected substrates.** *J Pharm Pharmacol* 2004, **56**:967-975.
- Doran A, Obach RS, Smith BJ, Hosea NA, Becker S, Callegari E, et al.: **The impact of P-glycoprotein on the disposition of drugs targeted for indications of the central nervous system: evaluation**

- tion using the **MDR1A/1B** knockout mouse model. *Drug Metab Dispos* 2005, **33**:165-174.
40. Szakacs G, Varadi A, Ozvegy-Laczka C, Sarkadi B: **The role of ABC transporters in drug absorption, distribution, metabolism, excretion and toxicity (ADME-Tox)**. *Drug Discov Today* 2008, **13**:379-393.
 41. Brady JM, Cherrington NJ, Hartley DP, Buist SC, Li N, Klaassen CD: **Tissue distribution and chemical induction of multiple drug resistance genes in rats**. *Drug Metab Dispos* 2002, **30**:838-844.
 42. Choudhuri S, Cherrington NJ, Li N, Klaassen CD: **Constitutive expression of various xenobiotic and endobiotic transporter mRNAs in the choroid plexus of rats**. *Drug Metab Dispos* 2003, **31**:1337-1345.
 43. Rao VV, Dahlheimer JL, Bardgett ME, Snyder AZ, Finch RA, Sartorelli AC, et al.: **Choroid plexus epithelial expression of MDR1 P glycoprotein and multidrug resistance-associated protein contribute to the blood-cerebrospinal-fluid drug-permeability barrier**. *Proc Natl Acad Sci USA* 1999, **96**:3900-3905.
 44. Smith AJ, van Helvoort A, van Meer G, Szabo K, Welker E, Szakacs G, et al.: **MDR3 P-glycoprotein, a phosphatidylcholine translocase, transports several cytotoxic drugs and directly interacts with drugs as judged by interference with nucleotide trapping**. *J Biol Chem* 2000, **275**:23530-23539.
 45. Wijnhouds J, deLange EC, Scheffer GL, Berg DJ van den, Mol CA, van d V, et al.: **Multidrug resistance protein 1 protects the choroid plexus epithelium and contributes to the blood-cerebrospinal fluid barrier**. *J Clin Invest* 2000, **105**:279-285.
 46. Gazzin S, Strazielle N, Schmitt C, Fevre-Montange M, Ostrow JD, Tiribelli C, et al.: **Differential expression of the multidrug resistance-related proteins ABCB1 and ABCG1 between blood-brain interfaces**. *J Comp Neurol* 2008, **510**:497-507.
 47. Scotto KW: **Transcriptional regulation of ABC drug transporters**. *Oncogene* 2003, **22**:7496-7511.
 48. Eloranta JJ, Kullak-Ublick GA: **Coordinate transcriptional regulation of bile acid homeostasis and drug metabolism**. *Arch Biochem Biophys* 2005, **433**:397-412.
 49. Kamiyama Y, Matsubara T, Yoshinari K, Nagata K, Kamimura H, Yamazoe Y: **Role of human hepatocyte nuclear factor 4alpha in the expression of drug-metabolizing enzymes and transporters in human hepatocytes assessed by use of small interfering RNA**. *Drug Metab Pharmacokin* 2007, **22**:287-298.
 50. Dignam JD, Lebovitz RM, Roeder RG: **Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei**. *Nucleic Acids Res* 1983, **11**:1475-1489.
 51. Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, et al.: **Measurement of protein using bicinchoninic acid**. *Anal Biochem* 1985, **150**:76-85.
 52. Kel A, Reymann S, Matys V, Nettesheim P, Wingender E, Borlak J: **A novel computational approach for the prediction of networked transcription factors of aryl hydrocarbon-receptor-regulated genes**. *Mol Pharmacol* 2004, **66**:1557-1572.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

