



ORAL PRESENTATION

Open Access

EZH2 dependent epigenetic landscape in adult T cell leukemia and Tax immortalized cells

Dai Fujikawa¹, Makoto Yamagishi¹, Shota Nakagawa¹, Naoya Kurokawa¹, Ai Soejima¹, Seiichirou Kobayashi², Kaoru Uchimaru², Yuetsu Tanaka³, Kazumi Nakano¹, Toshiki Watanabe^{1*}

From 17th International Conference on Human Retroviruses: HTLV and Related Viruses
Trois Ilets, Martinique. 18-21 June 2015

Epigenetic regulations globally determine gene transcription. Recent studies have revealed that expression changes and genetic mutations of epigenetic factors cause epigenetic imbalance in cancers. We previously reported that aberrant expression of Polycomb repressive complex 2 (PRC2) components causes constitutive NF- κ B activation through silencing of miR-31 in ATL cells. However, the underlying mechanisms by which the epigenetic imbalance is induced and maintained remain to be elucidated. Here, we conducted ChIP-on-chip and transcriptome analyses of ATL and normal CD4+ T cells and found that the epigenetic reprogramming closely associates with ATL specific gene expression signature. Leukemic cell-specific silencing of cell cycle regulator, CDKN1A, was correlated with H3K27me3 level. In addition, orchestrated loss of microRNAs and multiple transcription factors appears to be mediated by the epigenetic mechanism. EZH2 upregulation and H3K27me3 accumulation were found in HTLV-1-infected populations derived from ATL patients as well as asymptomatic carriers. Furthermore, EZH2 inhibition blocked Tax-dependent cell growth and its immortalization in vitro. Intriguingly, the Tax-triggered immortalizing cells partially mimicked the methylation pattern observed in ATL cells, suggesting that the epigenetic alterations are closely involved in immortalization of infected cells and disease progression. In parallel, we found that over expression of PRC2 core components was induced by active signaling cascades including NF- κ B in ATL cells. Collectively, our results suggest a coherent positive feedback mechanism comprised of PRC2 and NF- κ B signaling, which stabilizes the epigenetic rearrangement and phenotypic

outcomes in HTLV-1 infected cells. Since pharmacological inhibition of EZH2 selectively killed ATL and HTLV-1 infected cells in ex vivo culture, targeting the epigenetic elements will hold great promise in treatment and prevention of ATL and HTLV-1-related diseases.

Authors' details

¹Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan.

²Institute of Medical Science, The University of Tokyo, Tokyo, Japan.

³Graduate School and Faculty of Medicine, University of the Ryukyus, Okinawa, Japan.

Published: 28 August 2015

doi:10.1186/1742-4690-12-S1-O13

Cite this article as: Fujikawa *et al.*: EZH2 dependent epigenetic landscape in adult T cell leukemia and Tax immortalized cells. *Retrovirology* 2015 **12**(Suppl 1):O13.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: tnabe@ims.u-tokyo.ac.jp

¹Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan
Full list of author information is available at the end of the article

