

Poster presentation

Open Access

Suicide risk, aggression and violence in major psychiatric disorder Seyed Mousavi* and Hamid Tallaei

Address: Behavioural Sciences Research Center, Isfahan University of Medical Sciences, Noor Hospital, Iran

* Corresponding author

from International Society on Brain and Behaviour: 2nd International Congress on Brain and Behaviour
Thessaloniki, Greece. 17–20 November 2005

Published: 28 February 2006

Annals of General Psychiatry 2006, **5**(Suppl 1):S298 doi:10.1186/1744-859X-5-S1-S298

Background

Aggression, violence and Suicide are important problems of mental health and society, they almost always cause disability, death, or other social problems recognizing distribution of behaviors, and assessing suicide risk in disorders can guide us to appropriate measures.

Materials and methods

This is a cross-sectional, and, descriptive-analytic study. We studied 801 psychiatric patients who admitted in a psychiatric emergency unit, in Isfahan (Iran), about aggression, Violence and risk of suicide, by a 30-items questionnaire, filled by researcher. Then data collected and analysed by statistical tools

Results

About one-third of patients had aggression (and) or Violence in admission, or, during hours before it. It was found most in men, 12–26 years old ages, and in Bipolar mood disorder patients. "High suicide Risk" measures were most found in patients with major depressive disorder patients. Differences of these phenomenon in disorders were statistically meaningful.

Discussion

These finding shows more aggression and violence in emergency psychiatric patients than previous studies in other countries. It may be due to more prevalence of bipolar patients in the study field. The up most finding of "High suicidal risk" patients in major depression make an impression for systematic preventive program of it, again. The data show a small epidemiologic map like of the mentioned phenomenon.

References

1. Sadock VA, Sadock B: *Synopsis of Psychiatry Philadelphia: Williams and Wilkins*; 2003:154-158. 288, 920, 901–921

2. Mann JJ, Waternaux C, Haas GL, Malone KM: **Toward a clinical model of suicidal behaviour in psychiatric patients.** *Am J Psychiatry* 1999, **156**:181-189.
3. Brodsky BS, Oquendo M, Ellis SP, Haas GL, Malone KM, Mann JJ: **The relationship of childhood abuse to impulsivity and suicidal behaviour in adults with major depression.** *Am J Psychiatry* 2001, **158**:1871-1877.
4. Steinert T, Wiebe C, Gebhardt RP: **Aggressive behavior against self and others among first admission patients with schizophrenia.** *Psychiatr Serv* 1999, **50**:85-89.