## CORRECTION

## **Open Access**



# Correction to: Maternal personality disorder symptoms in primary health care: associations with mother-toddler interactions at one-year follow-up

Magnhild Singstad Høivik<sup>1,3\*</sup>, Stian Lydersen<sup>2</sup>, Ingunn Ranøyen<sup>2</sup> and Turid Suzanne Berg-Nielsen<sup>4</sup>

## Correction to: BMC Psychiatry (2018) 18:198 DOI: 10.1186/s12888-018-1789-5

Following publication of the original article [1], the authors notified us of an error in the data presented in Table 2: the DIP-Q subscales show an incorrect scale.

The corrected Table 2 is presented below.

### Author details

<sup>1</sup>Department of Mental Health, Faculty of Medicine and Health Sciences, the Norwegian University of Science and Technology (NTNU), N-7491 Trondheim, Norway. <sup>2</sup>Regional Centre for Child and Youth Mental Health and Child Welfare – Central Norway, Faculty of Medicine and Health Sciences, The Norwegian University of Science and Technology (NTNU), Trondheim, Norway. <sup>3</sup>Division of Psychiatry, St Olavs Hospital, Trondheim University Hospital, Trondheim, Norway. <sup>4</sup>The Centre for Child and Adolescent Mental Health, Eastern and Southern Norway, Oslo, Norway.

### Received: 23 November 2018 Accepted: 23 November 2018 Published online: 13 December 2018

#### Reference

 Høivik, et al. Maternal personality disorder symptoms in primary health care: associations with mother–toddler interactions at one-year follow-up. 2018; 18:198. https://doi.org/10.1186/s12888-018-1789-5.

\* Correspondence: magnhild.s.hoivik@ntnu.no

<sup>3</sup>Division of Psychiatry, St Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

Full list of author information is available at the end of the article



© The Author(s). 2018 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

<sup>&</sup>lt;sup>1</sup>Department of Mental Health, Faculty of Medicine and Health Sciences, the Norwegian University of Science and Technology (NTNU), N-7491 Trondheim, Norway

| Baseline                         |                  |       |      |                 |         |       |                 |         |      |     | Follow up |      |                        |                                     |                |  |
|----------------------------------|------------------|-------|------|-----------------|---------|-------|-----------------|---------|------|-----|-----------|------|------------------------|-------------------------------------|----------------|--|
| Personality disorder<br>symptoms | All participants |       |      | Attrition group |         |       | Remaining group |         |      |     |           |      |                        |                                     |                |  |
|                                  | n                | mean  | sd   | n               | mean    | sd    | n               | mean    | sd   | n   | mean      | sd   | cut off<br>(diagnosis) | n (%) with symptoms<br>over cut off | Scale<br>range |  |
| Avoidant                         | 122              | 1.93  | 1.95 | 28              | 2.29    | 2.56  | 94              | 1.82    | 1.85 | -   | _         | -    | ≥ 4                    | 26 (21.3)                           | 0–7            |  |
| Dependent                        | 122              | 1.81  | 1.88 | 28              | 2.61*** | 2.39  | 94              | 1.57*** | 1.64 | -   | -         | -    | ≥ 5                    | 13 (10.7)                           | 0–9            |  |
| Obsessive-compulsive             | 122              | 3.88  | 1.74 | 28              | 4.00    | 1.89  | 94              | 3.84    | 1.70 | -   | -         | -    | ≥ 4                    | 69 (56.6)                           | 0–9            |  |
| Paranoid                         | 122              | 1.34  | 1.60 | 28              | 2.04*** | 2.03  | 94              | 1.14*** | 1.40 | -   | _         | -    | ≥ 4                    | 8 (6.6)                             | 0–8            |  |
| Schizoid                         | 122              | 0.73  | 0.97 | 28              | 0.68    | 0.86  | 94              | 0.75    | 1.00 | -   | -         | -    | ≥ 4                    | 1 (0.8)                             | 0–8            |  |
| Schizotypal                      | 122              | 0.41  | 1.66 | 28              | 2.04    | 1.88  | 94              | 1.22    | 1.55 | -   | -         | -    | ≥ 5                    | 10 (8.2)                            | 0-11           |  |
| Antisocial                       | 122              | 0.85  | 0.85 | 28              | 0.86    | 0.85  | 94              | 0.85    | 0.94 | -   | _         | -    | ≥ 3                    | 6 (4.9)                             | 0–10           |  |
| Borderline                       | 122              | 2.48  | 2.15 | 28              | 3.61*** | 2.69  | 94              | 2.14*** | 1.85 | -   | _         | -    | ≥ 5                    | 19 (15.6)                           | 0–17           |  |
| Histrionic                       | 122              | 1.25  | 1.22 | 28              | 1.39    | 1.17  | 94              | 1.21    | 1.24 | -   | -         | -    | ≥ 5                    | 1 (0.8)                             | 0–8            |  |
| Narcissistic                     | 122              | 0.86  | 1.05 | 28              | 1.29**  | 1.33  | 94              | 0.73**  | 0.87 | -   | _         | -    | ≥ 5                    | 1 (0.8)                             | 0–10           |  |
| Impairment and distress          | 122              | 0.66  | 1.03 | 28              | 0.80*   | 1.26  | 94              | 0.63*   | 0.96 | -   | _         | -    | ≥ 2                    | 20 (16.4)                           | 0–5            |  |
| Depressive symptoms              | 118              | 12.11 | 8.64 | 24              | 15.96*  | 11.23 | 94              | 10.99** | 7.84 | 85  | 8.74***   | 7.05 | -                      | -                                   | 0–63           |  |
| EAS subscales                    |                  |       |      |                 |         |       |                 |         |      |     |           |      |                        |                                     |                |  |
| Maternal sensitivity             | 152              | 22.41 | 5.12 | 42              | 21.48   | 4.94  | 110             | 22.77   | 5.16 | 110 | 25.29***  | 3.92 | -                      | -                                   | 7–29           |  |
| Maternal structuring             | 152              | 23.26 | 4.50 | 42              | 22.38   | 4.29  | 110             | 23.60   | 4.55 | 110 | 25.90***  | 3.39 | -                      | -                                   | 7–29           |  |
| Maternal non-hostility           | 152              | 26.01 | 3.58 | 42              | 24.92*  | 4.18  | 110             | 26.44*  | 3.24 | 110 | 27.32***  | 2.50 | -                      | -                                   | 7–29           |  |
| Maternal<br>non-intrusiveness    | 152              | 22.24 | 5.72 | 42              | 21.26   | 6.16  | 110             | 22.63   | 5.52 | 110 | 25.27***  | 4.34 | -                      | -                                   | 7–29           |  |
| Child responsiveness             | 152              | 22.66 | 5.36 | 42              | 21.59   | 5.06  | 110             | 23.07   | 5.43 | 110 | 25.70***  | 4.08 | -                      | -                                   | 7–29           |  |
| Child involvement                | 152              | 22.43 | 5.94 | 42              | 20.50   | 5.97  | 110             | 21.79   | 5.92 | 110 | 25.11***  | 4.71 | -                      | -                                   | 7–29           |  |

Table 2 Sample clinical characteristics at baseline and at one-year follow-up

Maternal personality disorder symptoms measured with the DIP-Q (DSM IV and ICD-10 Personality Questionnaire) \* = p < 0.05, \*\* = p < 0.01, (independent samples t-tests of characteristics in attrition compared to remaining groups in baseline sample, and paired sample t-tests comparing characteristics of remaining and follow up groups)