

RESEARCH ARTICLE

Open Access



Effects on postgraduate-year-I residents of simulation-based learning compared to traditional lecture-style education led by postgraduate-year-II residents: a pilot study

Akira Yamamoto¹, Mikako Obika^{1*} , Yasuhiro Mandai², Taku Murakami¹, Tomoko Miyoshi², Hideo Ino², Hitomi Kataoka³ and Fumio Otsuka¹

Abstract

Background: Simulation-based learning plays an important role in contemporary medical education, although there are problems providing tutors. Peer-assisted learning has begun being formally adopted in medical education. Although it is considered useful for simulation-based learning, its effectiveness remains unclear. This study was designed to compare the effect of simulation-based learning with that of traditional lectures conducted by postgraduate-year (PGY)-II residents on PGY-I residents.

Methods: This study was conducted at Okayama University Hospital over three years, for one week each year, before residents entered clinical practice. The study enrolled 76 PGY-I residents, who were randomized into two groups: simulation and lecture groups. PGY-II residents volunteered to conduct simulations and lectures. Knowledge evaluation was performed using pre- and post-tests, and self-evaluation of competence and behaviour-change and program evaluations were conducted using questionnaires.

Results: In both groups, knowledge test scores were found to improve significantly, and the score difference between pre- and post-tests in both the groups was not significant. Self-evaluation of competence and behaviour-change was found to be higher in the simulation group than the lecture group. The trainees in the simulation group valued the program and the PGY-II residents as teaching staff more than those in the lecture group.

Conclusions: The combination of simulation-based learning and peer-assisted learning led by PGY-II residents is potentially more effective in improving the postgraduate education of PGY-I residents than the combination of lecture and peer-assisted learning.

Keywords: Simulation-based learning, Peer-assisted learning, Lecture, Postgraduate education, Junior residents

Background

Previous medical training using the philosophy of ‘see one, do one, teach one’ posed inherent risks to both patients and trainees. Simulation-based learning can provide a risk-free environment where trainees are permitted to make mistakes, which will reduce the occurrence of errors at clinical sites [1, 2]. A meta-analysis showed that simulation-based

learning was more effective than traditional education for trainees to acquire medical technical skills [3]. However, the effectiveness of simulation-based learning for the acquisition of initial case management skills, which junior residents are required to learn, has not yet been confirmed. This study aims to examine this research gap.

On the practical side of postgraduate medical education in Japan, postgraduate-year-I (PGY-I) residents often receive formal and informal ‘near’ peer-assisted learning (PAL) from PGY-II residents. PAL is defined as follows: ‘trainees from similar social groupings who are not professional teachers helping each other to learn and

* Correspondence: obika-m@cc.okayama-u.ac.jp

¹Department of General Medicine, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan

Full list of author information is available at the end of the article



learning themselves by teaching'. [4] To the best of our knowledge, there are no studies that report the effects of simulation-based learning conducted by PGY-II residents for PGY-I residents, although many studies have revealed that PAL and simulation-based learning have educational benefits [5, 6].

Therefore, we offered a simulation-based learning and traditional lecture course conducted by PGY-II residents for PGY-I residents. We evaluated this program using three measures (knowledge evaluation, self-evaluation of competence and behaviour-change, and program evaluation) and compared the benefits of simulation-based learning with those of traditional lectures.

Methods

This interventional education study was conducted at Okayama University Hospital for one week each year for three years on PGY-I residents before they began clinical practice. During the specified week, new employee training was provided to teach the residents what they needed to know about working in the hospital, such as how to use electronic medical records. We randomized the residents by the order of their names in the Japanese syllabary and divided them into two groups (Fig. 1): simulation-based

learning (simulation group, $n = 38$) and traditional lecture style (lecture group, $n = 37$). One resident was excluded because of insufficient data. Each year, we evaluated the PGY-I residents who had agreed to participate.

To develop case scenarios and goals, in the year prior to conducting this study, we asked PGY-I residents about the type of cases they should experience in a PGY-I internship. Based on their responses, we conducted an alpha test comprising simulations of candidate cases and evaluated the difficulty, validity, accuracy, and effectiveness of the cases. Following the alpha test, we conducted a beta test consisting of experimental simulations with the residents to evaluate and calibrate cases. Finally, we determined three case scenarios (acute myocardial infarction, multiple injuries, and aspiration pneumonia) that were considered feasible and effective for PGY-I residents (Table 1). All teaching materials were closely examined by experts.

Each year, volunteer PGY-II residents conducted simulations and lectures as teaching staff. In the simulation group, PGY-I residents were divided into two groups by name. A few minutes were spent explaining the equipment, following which case presentation, team discussion, simulation training, and debriefing took 1, 3, 5, and





Fig. 1 Selection and categorisation of participants in the study

Table 1 Details of scenarios

	Simulation group (n = 38)	Lecture group (n = 37)
Cases and goals	<p>Case 1. An outpatient with acute myocardial infarction</p> <p>Understanding basic physical examination methods of walk-in patients.</p> <p>Understanding the initial evaluation and examination of patients with chest pain.</p> <p>Understanding differential diagnosis of fatal chest pain.</p> <p>Understanding acute myocardial infarction.</p> <p>Case 2. A patient with multiple injury coming by an ambulance</p> <p>Understanding the initial evaluation and treatment of high energy trauma.</p> <p>Understanding imaging findings of intraabdominal bleeding and multiple fractures.</p> <p>Case 3. An inpatient with pneumonia</p> <p>Understanding the initial evaluation and treatment at the time of sudden change to a fatal disease of a hospitalized patient.</p> <p>Understanding the method of report to an advanced doctor.</p> <p>Understanding pneumonia.</p>	
total time of each case	one hour	one hour
detail of time coarse (minutes)	explanation of equipment (5) case presentation (1) † team discussion (3) † simulation training (5) † debriefing (8) † †Repeating 3 times	lecture (60)
Number of each group	12–14†† ††divided to two groups in this group	11–14
Teaching staff	Volunteer postgraduate-year-II residents	Volunteer postgraduate-year-II residents

8 min, respectively. Since debriefing is considered the most important part of a simulation, we allocated more time to it, and the teaching staff received debriefing education before participating in the program [7]. They repeated the scenario, excluding the explanation of the equipment, three times while changing the roles of PGY-I residents as a doctor, a nurse, and a secretary writing down their orders or the treatment for the debriefing. Each scenario took a total of one hour. They were provided simulation tools such as *SimMan 3G*, a monitor, an emergency cart with medications, an intravenous drip, a cardiac defibrillator, an intubation instrument, and a white board to record the scenario's progress. The teaching staff presented the results of blood, X-ray, CT scan, electro-cardiogram, and ultrasound sonography tests when requested by trainees. The lecture group received a one-hour, one-way communication lecture based on the same case and goals that were used in simulation-based learning. Both groups had three sessions per week. Every year, after finishing the program, all residents were given the opportunity to participate in simulation-based learning and lectures.

For knowledge evaluation (Fig. 2), we prepared a pre- and a post-test, including 24 multiple-choice and 28 free-response questions based on the three scenarios (adjusted to a total of 0–100 points). For both groups, we administered a questionnaire immediately after each simulation and lecture for the self-evaluation of competence and program evaluation. Regarding the self-evaluation of competence, the questionnaire asked the degree to which the participants' knowledge, judgement, and skills had improved and their degree of confidence in seeing a patient (1–10 points each; 4–40 points in total). For program evaluation, the questionnaire asked about the difficulty of the case and meaningfulness of the program (1–10 points) (Table 2). At the end of the program, a questionnaire was used in all groups for the self-evaluation of behaviour-change and program evaluation (Table 3). Regarding the self-evaluation of behaviour-change, the questionnaire asked about the residents' self-learning time after each training session (1–5 points) and the degree of behaviour-change. For program evaluation, questions were asked about the degree of stress that resulted from this program, residents' expectations that this program would relieve stress in clinical situations, the



appropriateness of the program’s timing, the competency of teaching staff, and the residents’ recommendation whether this program should be continued (1–10 points). The questionnaires included adequate space for free responses about the program and teaching staff, as well. The questionnaires were created after literature review and expert validation [8, 9].

The statistical analysis of the population was conducted using a chi-squared test. Further, Student’s t-test was used to compare knowledge test scores and the total scores of the questionnaires after each simulation and lecture. In addition, a paired t-test was used to determine the improvement in knowledge test scores and self-evaluations after cases. For other data that required nonparametric tests, the Mann-Whitney test was used. In addition, we controlled the family-wise error rate using the Benjamini-Hochberg’s method for the adjustment of multiple comparisons [10]. The statistical analyses were performed using SPSS v.24.0 software for Windows.

This study was approved by the ethics committee of the Okayama Medical School.

Results

Population

There were no significant differences in the baseline demographic data of the two groups in the following characteristics: gender, experience in simulation-based learning as a trainee or as teaching staff, motivation to become teaching staff for simulation education, being a graduate of the Okayama Medical School, and self-study time each day ($p > 0.05$) (Table 4).

Knowledge evaluation

The pre-test scores of the two groups were not statistically different (mean score of simulation group, 52.3; mean score of lecture group, 54.5; 95% confidence interval [CI], - 9.3 to 5; p value, 0.50). A comparison of their pre-tests and post-tests revealed that both the groups improved their scores ($p < 0.001$) (Table 5). Further, no

Table 2 Questionnaire administered immediately after each simulation/lecture (depicted using black arrowheads in Fig. 2)

Category	Item	Points
Self-evaluation	Do you think your skill improved by receiving simulation-based learning/lecture?	Extremely true Non at all 10 9 8 7 6 5 4 3 2 1
Self-evaluation	Do you think your knowledge improved by receiving simulation-based learning/lecture?	Extremely true Non at all 10 9 8 7 6 5 4 3 2 1
Self-evaluation	Do you think your judgment improved by receiving simulation-based learning/lecture?	Extremely true Non at all 10 9 8 7 6 5 4 3 2 1
Self-evaluation	Do you think you can see a patient similar to this case in clinical sites?	Extremely true Non at all 10 9 8 7 6 5 4 3 2 1
Self-evaluation	Total score	Max Minimum 40 39 38 6 5 4
Program evaluation	How difficult was simulation-based learning/lecture education for you?	Easy Difficult 10 9 8 7 6 5 4 3 2 1
Program evaluation	Was simulation/lecture education meaningful for you?	Extremely true Non at all 10 9 8 7 6 5 4 3 2 1

Table 3 Questionnaire administered at the end of the program (depicted using white arrowheads in Fig. 2)

Category	Item	Points
Program evaluation	Did you get stress by receiving simulation/lecture?	Extremely true 10 9 8 7 6 5 4 3 2 1 Non at all
Program evaluation	Was it appropriate timing to participate in simulation/lecture for residents?	Extremely true 10 9 8 7 6 5 4 3 2 1 Non at all
Program evaluation	Should residents continue simulation/lecture education?	Extremely true 10 9 8 7 6 5 4 3 2 1 Non at all
Program evaluation	Will simulation/lecture education relieve stress in clinical sites for residents?	Extremely true 10 9 8 7 6 5 4 3 2 1 Non at all
Program evaluation	Are the 2nd-year residents appropriate leaders in simulation/lecture?	Extremely true 10 9 8 7 6 5 4 3 2 1 Non at all
Behaviour-change evaluation	Did you improve your behaviour by receiving simulation/lecture?	Extremely true 10 9 8 7 6 5 4 3 2 1 Non at all
Behaviour-change evaluation	How long did you study before simulation/lecture per day?	None, <30min, 30-60min, 1-2hours, >2hours 1 2 3 4 5
Behaviour-change evaluation	How long did you study after the 1st case in simulation/lecture?	None, <30min, 30-60min, 1-2hours, >2hours 1 2 3 4 5
Behaviour-change evaluation	How long did you study after the 2nd case in simulation/lecture?	None, <30min, 30-60min, 1-2hours, >2hours 1 2 3 4 5
Behaviour-change evaluation	How long did you study after the 3rd case in simulation/lecture?	None, <30min, 30-60min, 1-2hours, >2hours 1 2 3 4 5

difference was detected in score improvement between the groups ($p > 0.05$).

Self-evaluation of competence and behaviour-change

The total self-evaluation of competence scores after each scenario in the simulation group were higher than those in the lecture group (Case, simulation group mean points (SD) - lecture group mean points (SD), p value: Case.1, 31.1 (4.3) - 28.2 (6.8), 0.04; Case.2, 32.2 (4.7) - 28.9 (5.4), 0.02; Case.3, 33.2 (4.3) - 29.9 (6.5), 0.02) (Table 6).

Following simulation-based learning and lectures, the residents in the simulation group thought that their behaviour improved more than the behaviour of those in the lecture group (simulation group median points interquartile range (IQR), lecture group median points (IQR), p value: 9 (1)- 7.5 (2.75), < 0.001 , respectively) (Table 7). Regarding specific behaviour changes, the simulation group spent more time on self-learning than the lecture group, with a significant difference after each scenario (Case, simulation group median points (IQR) - lecture group median

points (IQR), p value: Case.1, 2 (0) - 2 (1), 0.001; Case.2, 2 (1.5) - 1 (1), 0.02; Case.3, 2 (2) - 1 (1), 0.02).

Program evaluation

The residents in the simulation group felt that Case.1 was more difficult than did those in the lecture group (simulation group median points (IQR) - lecture group median points (IQR), p value: 5 (2) - 6 (2), 0.01) (Table 8). In a free response questionnaire about this program (Table 9), a few residents in the simulation group felt nervous about Case.1. However, after Case.2, there were no differences in the scores of difficulty ($p > 0.05$) or in stress from each educational intervention ($p > 0.05$), and no comments were made in the free response questionnaire about being nervous. Further, there were no differences in responses regarding the appropriate timing of the simulation and lecture ($p > 0.05$).

The simulation group felt the training they participated in was more meaningful than did the lecture group (Case, simulation group median points (IQR) - lecture group median points (IQR), p value: Case.1, 9.5

Table 4 Baseline demographic data

Resident characteristics	Simulation	Lecture	p value
Male: Number of affirmation (%)	21 (55)	22 (60)	0.71
Have you ever had simulation-based learning in medical school?: Number of affirmation (%)	32 (84)	27 (73)	0.24
Have you ever been a tutor of simulation-based learning?: Number of affirmation (%)	7 (18)	6 (16)	0.84
Would you like to instruct simulation training?: Number of affirmation (%)	22 (18)	19 (51)	0.57
Did you belong to a study group or training club in medical school?: Number of affirmation (%)	15 (40)	12 (32)	0.53
Did you graduate from Okayama medical school?: Number of affirmation (%)	19 (50)	20 (54)	0.73
How long do you conventionally study everyday?: Median pointst (interquartile range) [points]	1 (1)	1 (1)	0.32

tNone, 1; < 30 min, 2; 30-60 min, 3; 1-2 h, 4; > 2 h, 5

Table 5 Knowledge evaluation using pre- and post-test scores (points)

	Pre-test		Post-test		difference	<i>p</i> value
	Mean	SD	Mean	SD		
Simulation group	52.3	12.3	69.6	9.3	17.3	<0.001**
Lecture group	54.5	13.9	70.9	11.1	16.7	<0.001**

p < 0.05, adjusted for multiple tests

***p* ≤ 0.001, adjusted for multiple tests

(1) - 8 (2), 0.01; Case.2, 10 (1) - 9 (2), 0.01; Case.3, 9.5 (2) - 9 (2), 0.03). Further, those in the simulation group felt more strongly that this training should continue in the future than did the participants in the lecture group (simulation group median points (IQR) - lecture group median points (IQR), *p* value: 10 (1) - 8 (2), 0.001, respectively). In a free response questionnaire about this program, they commented that they were motivated to learn more, learned a lot from the simulation-based training, had meaningful training, and so on.

The simulation group felt that participating in this training would relieve their stress in clinical situations more than did the lecture group (simulation group median points (IQR) - lecture group median points (IQR), *p* value: 9 (1.5) - 8 (2), 0.001, respectively) with comments in the free response portion of the questionnaire that they felt able to assume responsibilities at actual clinical sites.

Further, in the free response questionnaire about the teaching staff, both groups valued PGY-II, and no comments mentioned that the PGY-II residents were not appropriate teaching staff (Table 10). They commented that the reasons PGY-II residents were appropriate teaching staff included teaching them specifically what junior residents should and could do and having no boundaries regarding the questions they could ask. There was a significant difference between the two groups in the questionnaire scoring; the simulation group was more positive about PGY-II residents as appropriate teaching staff than the lecture group (simulation group median points (IQR) - lecture group median points (IQR), *p* value: 10 (1) - 9 (2), 0.02, respectively).

Table 6 Self-evaluation using questionnaire (1–10 points)

Item	Simulation		Lecture		<i>p</i> value
	Mean	SD	Mean	SD	
Total score of self-evaluation					
Case.1	31.1	4.3	28.2	6.8	0.04*
Case.2	32.2	4.7	28.9	5.4	0.02*
Case.3	33.2	4.3	29.9	6.5	0.02*

SD, standard deviation

**p* < 0.05, adjusted for multiple tests

Discussion

Compared to lectures incorporating PAL, simulation-based learning led by PGY-II had remarkable benefits for PGY-I residents from the perspective of the latter's knowledge acquisition and self-evaluation. Based on these data, we categorized the effects of this study into three themes: knowledge evaluation, self-evaluation of competence and behaviour-change, and program evaluation.

Knowledge evaluation

In terms of acquiring knowledge, the effect of simulation-based training on PGY-I residents was equivalent to that of lectures. Since the pre- and post-tests were self-derived, our data had limitations; however, these results have already been revealed by previous research [11]. A simulation is a process through which knowledge is translated into reasoned action [2]. In other words, a simulation translates 'knows' and 'knows how' into 'shows how' in the framework for clinical assessment specified by Miller [12]. The simulation group's improvement in clinical skills was evaluated by themselves and also by the PGY-II residents using checklists evaluating 'shows how' when it was detected (data not shown). Therefore, this suggests that simulation-based learning has educational effects on not only acquisition of knowledge but also improvement in 'shows how', whereas lectures are limited to the acquisition of knowledge.

Self-evaluation of competence and behaviour-change

Many studies have reported that simulation-based learning or PAL assists in the learning of medical procedures [5, 13]. Researchers have evaluated the educational effects on trainees using procedure time, checklists, written examinations, and other methods. In this study, we focused more attention on and evaluated trainees' awareness and behavioural changes, rather than their medical skills themselves. We chose this approach because educational methods that do not promote awareness, called 'spoon-feeding education', are not persistently effective for trainees, even if a trainee acquires knowledge or skills.

Self-efficacy affects individual behaviour to achieve goals. In the self-evaluations, it was suggested that simulation-based learning improved trainees' self-efficacy. Further, the simulation-based learning program was developed to include the concept of self-efficacy. Self-efficacy is known to have a substantial effect on a trainee's performance, particularly in nursing education [14]. It is derived from four principal sources of information: performance accomplishments, various experiences, verbal persuasion, and physiological states [15]. We provided all these sources in the simulation-based learning program. Trainees achieved their own successful experience (performance accomplishments) and saw what the others did

Table 7 Behaviour-change evaluation

Item	Simulation		Lecture		<i>p</i> value
	Median	IQR	Median	IQR	
Did you improve your behaviour by receiving simulation/lecture?	9	1	7.5	2.75	< 0.001**
Self-learning time after each case.					
Case.1	2	0	2	1	0.01*
Case.2	2	1.5	1	1	0.03*
Case.3	2	2	1	1	0.02*

Questionnaire of behaviour change (1–10 points) and self-learning time (1–5 points)

IQR, interquartile range

p < 0.05, adjusted for multiple tests

***p* ≤ 0.001, adjusted for multiple tests

in the same scenario (various experiences). In the debriefing, they could learn from their mutual reviews and the facilitator (verbal persuasion). In addition, a facilitator asked about their feelings when the training was done to objectively make the trainees aware of their feelings and revise them through mutual reviews (physiological states). Further, receiving simulation-based training while being observed by their peers seemed to differ from being trained by specialists whose positions were completely different from the perspectives of affinity and tension. The PAL conducted by PGY-II residents created a less stressful and more relaxed educational environment, as they commented in the free response questionnaire (physiological states). We detected that their self-efficacy gradually improved in the simulation group. We used high-fidelity patient simulators, such as *Simman* and real equipment, since previous research has shown that self-efficacy improves after the use of high-fidelity patient simulator scenarios [14]. In contrast, lecture-style education has only a

few of these elements. Therefore, we concluded that PGY-II residents were better teaching staff in simulation-based learning than in lectures from the perspective of self-efficacy.

The time spent on self-learning by residents increased, which, as a rule of thumb in Japan, seems to reflect motivation. It is known that self-efficacy interacts with motivation and behavior [16]. The residents in the simulation group felt that their behaviour improved more than that of the residents in the lecture group, as shown by the comments that they were motivated to learn more. Undoubtedly, they had other behaviour or awareness improvements that we were unable to detect in this study.

Program evaluation

In terms of the difficulty of cases, only Case.1 was evaluated as being more difficult by the simulation group compared to the lecture group. However, there was no

Table 8 Program evaluation by questionnaire (1–10 points)

Item	Simulation		Lecture		<i>p</i> value
	Median	IQR	Median	IQR	
How difficult was simulation-based learning/lecture education for you?					
Case.1	5	2	6	2	0.01*
Case.2	5	2	6	1	0.27
Case.3	5	1	6	2	0.09
Was simulation/lecture education meaningful for you?					
Case.1	9.5	1	8	2	0.01*
Case.2	10	1	9	2	0.01*
Case.3	9.5	2	9	2	0.04*
Did you get stress by receiving simulation/lecture?	4	3	4	3.75	0.17
Is it appropriate timing to participate in simulation/lecture for residents?	9	1.5	8	3	0.06
Should residents continue simulation/lecture education?	10	1	8	2	0.001**
Will simulation/lecture education relieve stress in clinical sites for residents?	9	1.5	8	2	0.001**
Are PGY-II residents appropriate teaching staffs in simulation/lecture?	10	1	9	2	0.02*

IQR, interquartile range

**p* < 0.05, adjusted for multiple tests

***p* ≤ 0.001, adjusted for multiple tests

Table 9 Program evaluation by free description questionnaire of the program

Simulation group	Lecture group
Context	Context
I thank to teaching staffs.	I learned a lot from this case.
I was motivated to learn more.	This case was easy for me to understand.
I learned a lot from this training.	I thank to teaching staffs.
This training was meaningful for me.	I understood the lack of my medical knowledge and skills.
I understood the lack of my medical knowledge and skills.	I was motivated to learn more.
I was able to assume the actual clinical site.	The teaching staffs are good at teaching.
This case was difficult for me to understand.	This case was difficult for me to understand.
I enjoyed this training.	
I felt my growth from Case.1 in Case. 2.	
I took this training using what I learned in the previous training.	
I was nervous in this training in Case. 1.	
I learned an importance of this training.	
By receiving this training, I would see outpatients in emergency department at my ease.	

difference in the stress resulting from each training session; hence, case difficulty did not seem to be a problem for the trainees. They also expected that the simulation would relieve more stress in clinical situations than the lecture. McMillan et al. [17] have already reported that simulation training relieved PGY-I residents' anxiety; hence, the PGY-I residents who participated in this simulation-based training were expected to feel less stress at clinical sites.

The residents were satisfied with the time spent in simulation-based learning and lectures during the orientation period. This is thought to have been derived from their anxiety to join clinical practice and their desire to rehearse clinical practice before starting it. In general, they will have similar experiences during simulation training in

the future. On the other hand, lectures are inferior to simulation-based learning in terms of the depiction of reality. Therefore, conducting a simulation-based learning program for residents before starting clinical training might resolve the residents' anxiety, satisfy their desire for practice, and be more meaningful than lectures.

Compared to the trainees in the lecture group, those in the simulation group more strongly felt that the PGY-II residents were appropriate teaching staff. In the free response questionnaire, they felt an affinity toward the PGY-II residents and the PGY-II residents had more empathy for them, having been in their shoes. Although it is a mainstream practice to have senior residents as teachers of other junior residents in Japanese clinical sites, there have been few studies which have focused on

Table 10 Program evaluation by free description questionnaire of teaching staff

Simulation group	Lecture group
Context	Context
PGY-II residents were proper teaching staffs, because they taught what junior residents should and could do concretely.	PGY-II residents were proper teaching staffs, because they taught us in the same position as us.
they had no boarder to ask a question.	what they taught was easy to understand.
they provided a relaxing environment.	they had no boarder to ask a question.
they knew PGY-I residents' feelings and sympathized with us.	they taught us based on their experience in PGY-I residents days.
they were good at teaching.	they knew what we did not know and should learn.
they taught us based on their experience in PGY-I residents days.	what they taught was packed with materials we wanted to learn.
what they taught was likely to be practiced soon in clinical sites.	
they taught us kindly and carefully.	
they were passionate.	
they were our goal one year from now.	
they motivated us to learn more.	

PAL using simulation-based education. In addition, there are general concerns that PGY-II residents may have insufficient knowledge and incorrectly teach topics. This negative aspect of PAL seemed more pronounced in simulation education than in lectures where tutors only teach prepared content. However, our program comprising only three hours of simulation training conducted by PGY-II residents revealed an equivalent effect on knowledge acquisition and a number of positive outcomes in the residents' self-evaluation of competence and behaviour-change and program evaluation over the traditional lecture-style training conducted by the same residents. There is a positive correlation between learning time and learning effect in simulation-based education [18]. In conclusion, it is suggested that simulation-based education conducted by PGY-II residents was more effective for PGY-I residents than traditional lectures.

Often, the staff who provide some of the special simulation-based trainings for the acquisition of specific medical skills are experts who have already received special training [13]. It is difficult for all hospitals to prepare such teaching staff. While PAL can potentially resolve the problem posed by the shortage of simulation-based learning tutors, from an ethical perspective, we must not force PGY-II residents to become tutors without demonstrating the associated advantages to them.

It is noted that our study has a few limitations. First, this study only suggests that PGY-II residents are more appropriate teaching staff for simulations than for lectures based on the comparison of data and free response questionnaires. Graham et al. [19] previously suggested that the use of PAL techniques and medical students to teach physical examinations is a comparable level of training to that of training by experts; a comparison study should be performed on simulation-based learning by PGY-II residents' PAL and experts teaching from the perspective of educational effects on trainees. Second, since we evaluated only the educational effects on PGY-I residents, it is necessary to consider the educational effects on PGY-II residents when they teach simulation-based learning. Third, since this study was conducted in a single facility and the questionnaires were limited in terms of validity in the course of their creation, this study should be considered a preliminary study, and further studies should be conducted in more facilities. Fourth, we examined only short-term educational effects and did not investigate the outcome in clinical situations. Currently, we are analysing data on PGY-I residents after they began clinical practice, focusing on the longer-term educational effects of this program. In addition, we evaluated their behaviour-change subjectively, rather than objectively, using questionnaires during the simulation and lecture. Future studies should examine the topic both subjectively and objectively.

Conclusions

Simulation-based learning and PAL are known to be effective educational methods. This study clarifies that a combination of these methods led by PGY-II residents, rather than specialists, can potentially better improve postgraduate education for PGY-I residents compared to a combination of lecture and PAL.

Abbreviations

PGY: Postgraduate-year; PAL: Peer-assisted learning.; SD: Standard deviation; IQR: Interquartile range.

Acknowledgements

Not applicable.

Funding

This work was supported by JSPS KAKENHI Grant Number 18 K10002. We used the grant for the proofreading of this paper in English.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

AY was the central in conceiving the research, coordinating activities, collecting the data, analysing it, interpreting it and drafting the initial manuscript. MO contributed to funding, analysing the data, interpreting it and drafting the initial manuscript. YM designed the research and interpreted the data. TM (Taku Murakami) was the central in conceiving the research, coordinating activities and collecting the data. TM (Tomoko Miyoshi), HI and HK designed the research. FO analysed the data. All authors contributed to the revision of the manuscript, approved the final version and are accountable for the accuracy and integrity of the content.

Ethics approval and consent to participate

Informed consent was obtained from all participants. This study was approved by the ethics committee of Okayama Medical School (No. 1602-039).

Consent for publication

Consent for publication was obtained from all participants.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of General Medicine, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan. ²Center for Education in Medicine and Health Sciences, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan. ³Department of Primary Care and Medical Education, Okayama University Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan.

Received: 18 December 2018 Accepted: 27 February 2019

Published online: 20 March 2019

References

1. Ziv A, Ben-David S, Ziv M. Simulation based medical education: an opportunity to learn from errors. *Med Teach*. 2005;27:193–9.
2. Lateef F. Simulation-based learning: just like the real thing. *J Emerg Trauma Shock*. 2010;3:348–52.
3. McGaghie WC, Issenberg SB, Cohen ER, Barsuk JH, Wayne DB. Does simulation-based medical education with deliberate practice yield better

- results than traditional clinical education? A meta-analytic comparative review of the evidence. *Acad Med*. 2011;86:706–11.
4. Topping KJ. The effectiveness of peer tutoring in further and higher education: a typology and review of the literature. *High Educ*. 1996;32:321–45.
 5. Ten Cate O, Durning S. Peer teaching in medical education: twelve reasons to move from theory to practice. *Med Teach*. 2007;29:591–9.
 6. Burgess A, McGregor D, Mellis C. Medical students as peer tutors: a systematic review. *BMC Med Educ*. 2014;14:115.
 7. Fanning RM, Gaba DM. The role of debriefing in simulation-based learning. *Simul Healthc*. 2007;2:115–25.
 8. Langan TS, Rigby IJ, Walker IW, Howes D, Donnon T, Lord JA. Simulation-based training in critical resuscitation procedures improves residents' competence. *Can J Emerg Med*. 2009;11(6):535–9.
 9. Weller JM. Simulation in undergraduate medical education: bridging the gap between theory and practice. *Med Educ*. 2004;38(1):32–8.
 10. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol*. 1995;57:289–300.
 11. Gordon JA, Shaffer DW, Raemer DB, Pawlowski J, Hurford WE, Cooper JB. A randomized controlled trial of simulation-based teaching versus traditional instruction in medicine: a pilot study among clinical medical students. *Adv Health Sci Educ*. 2006;11:33–9.
 12. Miller GE. The assessment of clinical skills/competence/performance. *Acad Medicine*. 1990;65:S63–7.
 13. Dayal AK, Fisher N, Magrane D, Goffman D, Bernstein PS, Katz NT. Simulation training improves medical students' learning experiences when performing real vaginal deliveries. *Simulation Healthc*. 2009;4:155–9.
 14. Weiler DT, Saleem JJ. Role assignment in high fidelity patient simulation and the effects on nursing students' self-efficacy development. *Proc Hum Factors Ergon Soc Annu Meet*. 2017;61:671–5.
 15. Bandura A. Self-efficacy: toward a unifying theory of behavioral change. *Adv Behav Res Ther*. 1978;1:139–61.
 16. Schunk DH. Self-efficacy, motivation, and performance. *J Appl Sport Psychol*. 1995;7:112–37.
 17. McMillan HJ, Writer H, Moreau KA, Eady K, Sell E, Lobos A-T, Grabowski J, Doja A. Lumbar puncture simulation in pediatric residency training: improving procedural competence and decreasing anxiety. *BMC Med Educ*. 2016;16:198.
 18. McGaghie WC, Issenberg SB, Petrusa ER, Scalese RJ. Effect of practice on standardised learning outcomes in simulation-based medical education. *Med Educ*. 2006;40:792–7.
 19. Graham K, Burke JM, Field M. Undergraduate rheumatology: can peer-assisted learning by medical students deliver equivalent training to that provided by specialist staff? *Rheumatology*. 2008;47:652–5.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

