

RESEARCH

Open Access



Approximate solutions of a sum-type fractional integro-differential equation by using Chebyshev and Legendre polynomials

Eisa Akbari Kojabad and Shahram Rezapour*

*Correspondence:
sh.rezapour@azaruniv.edu
Department of Mathematics,
Azarbaijan Shahid Madani
University, Tabriz, Iran

Abstract

We investigate the existence of solutions for a sum-type fractional integro-differential problem via the Caputo differentiation. By using the shifted Legendre and Chebyshev polynomials, we provide a numerical method for finding solutions for the problem. In this way, we give some examples to illustrate our results.

MSC: 26A33; 34A08; 34K37

Keywords: approximate fixed point; Chebyshev polynomial; Legendre polynomial; numerical solution; sum-type fractional integro-differential equation

1 Introduction

In 1969, Reinermann investigated some problems by using approximate fixed point property ([1]). In 1976, Yamamoto and Ohtsubo published a paper on subspace iteration accelerated by using Chebyshev polynomials for eigenvalue problems ([2]). There has been published some work about different fractional integro-differential equations by using Chebyshev polynomials ([3, 4] and [5]) or by using Legendre wavelets ([6–8] and [9]). Recently, different techniques for solving some fractional integro-differential equations have been used (see [6, 10–19]). In this paper by using an approximate fixed point result and the shifted Legendre and Chebyshev polynomials, we investigate the existence of solutions for a sum-type fractional integro-differential problem.

As is well known, the Caputo fractional derivative of order β for a continuous function $f : (0, \infty) \rightarrow \mathbb{R}$ is defined by ${}^C D^\beta f(t) = \frac{1}{\Gamma(n-\beta)} \int_0^t \frac{f^{(n)}(s)}{(t-s)^{\beta-n+1}} ds$, where $n = [\beta] + 1$ ([20, 21]). The fractional integral of order β for a function $f : (0, \infty) \rightarrow \mathbb{R}$ is defined by $I^\beta f(t) = \frac{1}{\Gamma(\beta)} \int_0^t (t-s)^{\beta-1} f(s) ds$ ([20, 21]). Let (X, d) be a metric space, T a selfmap on X and $\alpha : X \times X \rightarrow [0, \infty)$ a map. We say that T is α -admissible whenever $\alpha(x, y) \geq 1$ implies $\alpha(Tx, Ty) \geq 1$. Also, T is called α -contraction whenever there exists $\lambda \in (0, 1)$ such that $\alpha(x, y)d(Tx, Ty) \leq \lambda d(x, y)$ for all $x, y \in X$. We say that T has approximate fixed point property whenever there exists a sequence $\{x_n\}_{n \geq 1}$ in X such that $d(x_n, Tx_n) \rightarrow 0$. We need the following results.

Lemma 1.1 ([21]) *Let $q > 0$, $n = [q] + 1$ and $v \in C([0, 1], \mathbb{R})$. Then the fractional differential equation ${}^C D^q x(t) = v(t)$ has a solution in the form*

$$x(t) = I^q v(t) + c_0 + c_1 t + \cdots + c_{n-1} t^{n-1}.$$

© The Author(s) 2017. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Lemma 1.2 ([22]) Let (X, d) be a metric space and T an α -contractive and α -admissible selfmap on X such that $\alpha(x_0, Tx_0) \geq 1$ for some $x_0 \in X$. Then T has the approximate fixed point property. If X is complete and T is continuous, then T has fixed point.

2 Main result

Now, we are ready to study the existence of solution of the sum-type fractional integro-differential equation

$${}^cD^q x(t) = f(t, x(t), {}^cD^{\beta_1}x(t), \dots, {}^cD^{\beta_n}x(t)) + g(t, x(t), I^{\beta_1}x(t), \dots, I^{\beta_n}x(t)) \quad (2.1)$$

with boundary value conditions $\sum_{i=1}^n (a_i {}^cD^{\beta_i}x(1)) = \alpha_1 x'(1)$ and $\sum_{i=1}^n (b_i I^{\beta_i}x(1)) = \alpha_2 x'(0)$, where $1 < q < 2$, $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}$ and $f, g : [0, 1] \times \mathbb{R}^{n+2} \rightarrow \mathbb{R}$ are two maps.

Lemma 2.1 Let $1 < q < 2$ and $v \in C(I, \mathbb{R})$. Then the unique solution for the fractional differential equation ${}^cD^q x(t) = v(t)$ with boundary conditions $\sum_{i=1}^n (a_i {}^cD^{\beta_i}x(1)) = \alpha_1 x'(1)$ and $\sum_{i=1}^n (b_i I^{\beta_i}x(1)) = \alpha_2 x'(0)$ is given by

$$\begin{aligned} x(t) = & I^q v(t) - \frac{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1\right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1\right)\left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)}\right)\right)} \sum_{i=1}^n b_i I^{q+\beta_i} v(1) \\ & - \frac{\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2\right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1\right)\left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)}\right)\right)} \alpha_1 I^{q-1} v(1) \\ & + \frac{\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2\right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1\right)\left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)}\right)\right)} \sum_{i=1}^n a_i I^{q-\beta_i} v(1) \\ & + \frac{\alpha_1 t I^{q-1} v(1) - \sum_{i=1}^n (t a_i I^{q-\beta_i} v(1))}{\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1}, \end{aligned}$$

where $\alpha_1, \alpha_2, a_1, \dots, a_n, b_1, \dots, b_n$ are some real numbers.

Proof By using Lemma 1.1, general solution for the equation ${}^cD^q x(t) = v(t)$ is given by $x(t) = \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} v(s) ds + c_0 + c_1 t$, where $c_0, c_1 \in \mathbb{R}$. By applying the boundary condition $\sum_{i=1}^n (a_i {}^cD^{\beta_i}x(1)) = \alpha_1 x'(1)$, we get

$$\begin{aligned} & \sum_{i=0}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} v(s) ds + \frac{a_i c_1}{\Gamma(2-\beta_i)} + 0 \right) \\ & = \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} v(s) ds + \alpha_1 c_1 \end{aligned}$$

and by using the boundary condition $\sum_{i=1}^n (b_i I^{\beta_i}x(1)) = \alpha_2 x'(0)$, we get

$$\sum_{i=1}^n \left(\frac{b_i}{\Gamma(q+\beta_i)} \int_0^1 (1-s)^{q+\beta_i-1} v(s) ds + \frac{b_i c_0}{\Gamma(\beta_i+1)} + \frac{b_i c_1}{\Gamma(\beta_i+2)} \right) = \alpha_2 c_1.$$

This implies

$$\begin{aligned} c_1 & \left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \\ & = \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} v(s) ds - \sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} v(s) ds \right) \end{aligned}$$

and

$$\sum_{i=1}^n \left[c_1 \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right) + \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) c_0 \right] = - \sum_{i=1}^n \frac{b_i}{\Gamma(q+\beta_i)} \int_0^1 (1-s)^{q+\beta_i-1} v(s) ds.$$

Hence,

$$\begin{aligned} c_0 & = - \frac{\left(\sum_{i=1}^n \frac{b_i}{\Gamma(q+\beta_i)} \int_0^1 (1-s)^{q+\beta_i-1} v(s) ds \right) \left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \\ & \quad - \frac{\left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right) \right) \left(\frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} v(s) ds \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \\ & \quad + \frac{\left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right) \right) \left(\sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} v(s) ds \right) \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \end{aligned}$$

and

$$c_1 = \frac{\frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} v(s) ds - \sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} v(s) ds \right)}{\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1}.$$

Thus,

$$\begin{aligned} x(t) & = \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} v(s) ds \\ & \quad - \frac{\left(\sum_{i=1}^n \frac{b_i}{\Gamma(q+\beta_i)} \int_0^1 (1-s)^{q+\beta_i-1} v(s) ds \right) \left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \\ & \quad - \frac{\left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right) \right) \left(\frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} v(s) ds \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \\ & \quad + \frac{\left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right) \right) \left(\sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} v(s) ds \right) \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \\ & \quad + \frac{\frac{t\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} v(s) ds - t \sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} v(s) ds \right)}{\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1} \\ & = I^q v(t) - \frac{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \sum_{i=1}^n b_i I^{q+\beta_i} v(1) \\ & \quad - \frac{\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \alpha_1 I^{q-1} v(1) \end{aligned}$$

$$\begin{aligned}
& + \frac{\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \sum_{i=1}^n a_i I^{q-\beta_i} v(1) \\
& + \frac{\alpha_1 t I^{q-1} v(1) - \sum_{i=1}^n (t a_i I^{q-\beta_i} v(1))}{\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1}.
\end{aligned}$$

One can check that the given $x(t)$ is a solution for the problem ${}^c D^q x(t) = v(t)$ with the boundary conditions. This completes our proof. \square

Let $\mathcal{X} = \{x : x, {}^c D^{\beta_1} x, {}^c D^{\beta_2} x, \dots, {}^c D^{\beta_n} x \in C(I, \mathbb{R})\}$ be endowed with the metric

$$d(x, y) = \sup_{t \in I} |x(t) - y(t)| + \sup_{t \in I} |{}^c D^{\beta_1} x(t) - {}^c D^{\beta_1} y(t)| + \dots + \sup_{t \in I} |{}^c D^{\beta_n} x(t) - {}^c D^{\beta_n} y(t)|.$$

It is clear that (\mathcal{X}, d) is a complete metric space (see [23]). By using Lemma 2.1, a function $x \in \mathcal{X}$ is a solution for the fractional differential equation (2.1) whenever it satisfies the boundary conditions and there exist functions $v, v' \in L^1[0, 1]$ such that $v(t) = f(t, x(t), {}^c D^{\beta_1} x(t), \dots, {}^c D^{\beta_n} x(t)), v'(t) = g(t, x(t), I^{\beta_1} x(t), \dots, I^{\beta_n} x(t))$ and

$$\begin{aligned}
x(t) &= I^q (v(t) + v'(t)) \\
&- \frac{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \sum_{i=1}^n b_i I^{q+\beta_i} (v(1) + v'(1)) \\
&- \frac{\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \alpha_1 I^{q-1} (v(1) + v'(1)) \\
&+ \frac{\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+2)} - \alpha_2 \right)}{\left(\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1 \right) \left(\sum_{i=1}^n \left(\frac{b_i}{\Gamma(\beta_i+1)} \right) \right)} \sum_{i=1}^n a_i I^{q-\beta_i} (v(1) + v'(1)) \\
&+ \frac{\alpha_1 t I^{q-1} (v(1) + v'(1)) - \sum_{i=1}^n (t a_i I^{q-\beta_i} (v(1) + v'(1)))}{\sum_{i=1}^n \frac{a_i}{\Gamma(2-\beta_i)} - \alpha_1}
\end{aligned}$$

for all $t \in I$.

Theorem 2.2 Let $\xi : \mathbb{R}^{2(n+1)} \rightarrow \mathbb{R}$ be a map, $\lambda \in (0, 1)$ and $f, g : [0, 1] \times \mathbb{R}^{n+2} \rightarrow \mathbb{R}$ two functions such that

$$\begin{aligned}
& |f(t, x_1, x_2, \dots, x_{n+1}) - f(t, y_1, y_2, \dots, y_{n+1})| + |g(t, x_1, x_2, \dots, x_{n+1}) - g(t, y_1, y_2, \dots, y_{n+1})| \\
& \leq \frac{\lambda}{\Omega_1 + n\Omega_2} (|x_1 - y_1| + \dots + |x_{n+1} - y_{n+1}|)
\end{aligned}$$

for all $t \in I = [0, 1]$ and $x_1, \dots, x_n, y_1, \dots, y_n \in \mathbb{R}$ with

$$\xi(x_1, x_2, \dots, x_{n+1}, y_1, y_2, \dots, y_{n+1}) \geq 0,$$

where

$$\begin{aligned}
\Omega_1 &= \left[\left| \frac{1}{\Gamma(q+1)} \right| + \frac{\left| \frac{\alpha_1}{\Gamma(q)} \right| + \sum_{i=1}^n \left| \frac{5}{i} \lambda \right|}{\left| \sum_{i=1}^n \lambda_i^1 - \alpha_1 \right|} \right. \\
&\quad \left. + \frac{\left| \sum_{i=1}^n \lambda_i^1 - \alpha_1 \right| \left| \sum_{i=1}^n \frac{3}{i} \lambda \right| + \left| \sum_{i=1}^n \lambda_i^4 - \alpha_2 \right| \left(\left| \frac{\alpha_1}{\Gamma(q)} \right| + \left| \sum_{i=1}^n \frac{5}{i} \lambda \right| \right)}{\left| \left(\sum_{i=1}^n \lambda_i^1 - \alpha_1 \right) \sum_{i=1}^n \lambda_i^2 \right|} \right],
\end{aligned}$$

$$\Omega_2 = \max_{1 \leq j \leq n} \left(\left| \frac{1}{\Gamma(q - \beta_j + 1)} \right| + \left| \frac{\frac{\alpha_1}{\Gamma(q)}}{\Gamma(2 - \beta_j)(\sum_{i=1}^n \lambda_i^1 - \alpha_1)} \right| \right),$$

$$\lambda_i^1 = \frac{a_i}{1 - \Gamma(2 - \beta_i)}, \quad \lambda_i^2 = \frac{b_i}{\Gamma(\beta_i + 1)}, \quad \lambda_i^3 = \frac{b_i}{\Gamma(q + \beta_i)}, \quad {}_i^3\lambda = \frac{b_i}{\Gamma(q + \beta_i + 1)},$$

$$\lambda_i^4 = \frac{b_i}{\Gamma(\beta_i + 2)}, \quad \lambda_i^5 = \frac{a_i}{\Gamma(q - \beta_i)} \quad \text{and} \quad {}_i^5\lambda = \frac{a_i}{\Gamma(q - \beta_i + 1)}.$$

Assume that

$$\xi(u(t), {}^cD^{\beta_1}u(t), {}^cD^{\beta_2}u(t), \dots, {}^cD^{\beta_n}u(t), v(t), {}^cD^{\beta_1}v(t), {}^cD^{\beta_2}v(t), \dots, {}^cD^{\beta_n}v(t)) \geq 0$$

implies

$$\xi(Tu(t), {}^cD^{\beta_1}Tu(t), \dots, {}^cD^{\beta_n}Tu(t), Tv(t), {}^cD^{\beta_1}Tv(t), \dots, {}^cD^{\beta_n}Tv(t)) \geq 0,$$

where the operator $T : \mathcal{X} \rightarrow \mathcal{X}$ is defined by

$$\begin{aligned} Tu(t) &= \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \\ &\quad - \left[\frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\ &\quad \times \sum_{i=1}^n \left(\lambda_i^3 \int_0^1 (1-s)^{q+\beta_i-1} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \right) \\ &\quad - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) (\sum_{i=1}^n \lambda_i^2)} \right] \\ &\quad \times \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \\ &\quad - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\ &\quad \times \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \right) \\ &\quad + \frac{\frac{t\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\ &\quad - \frac{t \sum_{i=1}^n (\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds)}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\ &\quad + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} g(s, u(s), I^{\beta_1}u(s), \dots, I^{\beta_n}u(s)) ds \\ &\quad - \left[\frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\ &\quad \times \sum_{i=1}^n \left(\lambda_i^3 \int_0^1 (1-s)^{q+\beta_i-1} g(s, u(s), I^{\beta_1}u(s), \dots, I^{\beta_n}u(s)) ds \right) \\ &\quad - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) (\sum_{i=1}^n \lambda_i^2)} \right] \end{aligned}$$

$$\begin{aligned}
& \times \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\
& \times \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \right) \\
& + \frac{\frac{t\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\
& - \frac{t \sum_{i=1}^n (\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds)}{\sum_{i=1}^n \lambda_i^1 - \alpha_1}
\end{aligned}$$

for all $t \in I$. If there exists $u_1 \in \mathcal{X}$ such that

$$\xi(u_1(t), {}^cD^{\beta_1}u_1(t), \dots, {}^cD^{\beta_n}u_1(t), Tu_1(t), {}^cD^{\beta_1}Tu_1(t), \dots, {}^cD^{\beta_n}Tu_1(t)) \geq 0$$

for all $t \in [0, 1]$, then the problem (2.1) has an approximate solution.

Proof We define $\alpha : \mathcal{X} \times \mathcal{X} \rightarrow [0, \infty)$ by

$$\alpha(u, v) = \begin{cases} 1, & \xi(u(t), {}^cD^{\beta_1}u(t), \dots, {}^cD^{\beta_n}u(t), v(t), {}^cD^{\beta_1}v(t), \dots, {}^cD^{\beta_n}v(t)) \geq 0, \forall t \in I, \\ 0, & \text{else.} \end{cases}$$

We show that T is an α -admissible and α -contractive selfmap on \mathcal{X} . Let $u, v \in \mathcal{X}$ be such that $\xi(u(t), {}^cD^{\beta_1}u(t), \dots, {}^cD^{\beta_n}u(t), v(t), {}^cD^{\beta_1}v(t), \dots, {}^cD^{\beta_n}v(t)) \geq 0$ for all $t \in [0, 1]$. Then we have

$$\begin{aligned}
& |Tu(t) - Tv(t)| \\
& = \left| \left\{ \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \right. \right. \\
& \quad \left. \left. - \left[\frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \right. \right. \\
& \quad \left. \left. \times \sum_{i=1}^n \left(\lambda_i^3 \int_0^1 (1-s)^{q+\beta_i-1} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \right) \right. \right. \\
& \quad \left. \left. - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) (\sum_{i=1}^n \lambda_i^2)} \right] \right. \right. \\
& \quad \left. \left. \times \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \right. \right. \\
& \quad \left. \left. - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \right. \right. \\
& \quad \left. \left. \times \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} f(s, u(s), {}^cD^{\beta_1}u(s), \dots, {}^cD^{\beta_n}u(s)) ds \right) \right. \right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{\frac{t\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) ds}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\
& - \frac{t \sum_{i=1}^n (\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) ds)}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\
& + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\
& \times \sum_{i=1}^n \left(\lambda_i^3 \int_0^1 (1-s)^{q+\beta_i-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \right) \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) (\sum_{i=1}^n \lambda_i^2)} \right] \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\
& \times \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \right) \\
& + \frac{\frac{t\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\
& - \frac{t \sum_{i=1}^n (\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds)}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \Big\} \\
& - \left\{ \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds \right. \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\
& \times \sum_{i=1}^n \left(\lambda_i^3 \int_0^1 (1-s)^{q+\beta_i-1} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds \right) \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) (\sum_{i=1}^n \lambda_i^2)} \right] \\
& \times \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\
& \times \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds \right) \\
& + \frac{\frac{t\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\
& - \frac{t \sum_{i=1}^n (\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds)}{\sum_{i=1}^n \lambda_i^1 - \alpha_1}
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\
& \times \sum_{i=1}^n \left(\lambda_i^3 \int_0^1 (1-s)^{q+\beta_i-1} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds \right) \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) (\sum_{i=1}^n \lambda_i^2)} \right] \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds \\
& - \left[\frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right] \\
& \times \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds \right) \\
& + \frac{\frac{t\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \\
& - \frac{t \sum_{i=1}^n (\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds)}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \Bigg\} \\
& \leq \int_0^t \left| \frac{(t-s)^{q-1}}{\Gamma(q)} \right| \\
& \times |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \\
& + \left| \frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right| \\
& \times \sum_{i=1}^n \left(|\lambda_i^3| \int_0^1 (1-s)^{q+\beta_i-1} |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) \right. \\
& \left. - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \right) \\
& + \left| \frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) (\sum_{i=1}^n \lambda_i^2)} \right| \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) \right. \\
& \left. - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \right. \\
& + \left| \frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1) \sum_{i=1}^n \lambda_i^2} \right| \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) \right. \\
& \left. - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \right) \\
& + \left| \frac{\frac{t\alpha_1}{\Gamma(q-1)}}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \right| \int_0^1 (1-s)^{q-\beta_i-1} |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) \right. \\
& \left. - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \right. \\
& + \left| \frac{t \sum_{i=1}^n \lambda_i^5}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \right| \int_0^1 (1-s)^{q-\beta_i-1} |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) \right. \\
& \left. - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \right. \\
& + \int_0^t \left| \frac{(t-s)^{q-1}}{\Gamma(q)} \right| |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds
\end{aligned}$$

$$\begin{aligned}
& + \left| \frac{\sum_{i=1}^n \lambda_i^1 - \alpha_1}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1)(\sum_{i=1}^n \lambda_i^2)} \right| \sum_{i=1}^n \left(|\lambda_i^3| \int_0^1 (1-s)^{q+\beta_i-1} |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) \right. \\
& \quad \left. - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds \right) \\
& + \left| \frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1)(\sum_{i=1}^n \lambda_i^2)} \right| \frac{\alpha_1}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) \\
& \quad - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds \\
& + \left| \frac{\sum_{i=1}^n \lambda_i^4 - \alpha_2}{(\sum_{i=1}^n \lambda_i^1 - \alpha_1)(\sum_{i=1}^n \lambda_i^2)} \right| \sum_{i=1}^n \left(\lambda_i^5 \int_0^1 (1-s)^{q-\beta_i-1} |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) \right. \\
& \quad \left. - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds \right) \\
& + \left| \frac{\frac{t\alpha_1}{\Gamma(q-1)}}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \right| \int_0^1 (1-s)^{q-\beta_i-1} |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) \\
& \quad - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds \\
& + \left| \frac{t \sum_{i=1}^n \lambda_i^5}{\sum_{i=1}^n \lambda_i^1 - \alpha_1} \right| \int_0^1 (1-s)^{q-\beta_i-1} |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) \\
& \quad - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds \\
& \leq \left[\left| \frac{1}{\Gamma(q+1)} \right| + \frac{|\frac{\alpha_1}{\Gamma(q)}| + \sum_{i=1}^n |\lambda_i^5| \lambda_i}{|\sum_{i=1}^n \lambda_i^1 - \alpha_1|} \right. \\
& \quad \left. + \frac{|\sum_{i=1}^n \lambda_i^1 - \alpha_1| |\sum_{i=1}^n \lambda_i^3| + |\sum_{i=1}^n \lambda_i^4 - \alpha_2| (|\frac{\alpha_1}{\Gamma(q)}| + |\sum_{i=1}^n \lambda_i^5|)}{(|\sum_{i=1}^n \lambda_i^1 - \alpha_1| \sum_{i=1}^n \lambda_i^2)} \right] \\
& \quad \times \left(\sup_{t \in I} |f(t, u(t), {}^c D^{\beta_1} u(t), \dots, {}^c D^{\beta_n} u(t)) - f(t, v(t), {}^c D^{\beta_1} v(t), \dots, {}^c D^{\beta_n} v(t))| \right) \\
& \quad + \left[\left| \frac{1}{\Gamma(q+1)} \right| + \frac{|\frac{\alpha_1}{\Gamma(q)}| + \sum_{i=1}^n |\lambda_i^5| \lambda_i}{|\sum_{i=1}^n \lambda_i^1 - \alpha_1|} \right. \\
& \quad \left. + \frac{|\sum_{i=1}^n \lambda_i^1 - \alpha_1| |\sum_{i=1}^n \lambda_i^3| + |\sum_{i=1}^n \lambda_i^4 - \alpha_2| (|\frac{\alpha_1}{\Gamma(q)}| + |\sum_{i=1}^n \lambda_i^5|)}{(|\sum_{i=1}^n \lambda_i^1 - \alpha_1| \sum_{i=1}^n \lambda_i^2)} \right] \\
& \quad \times \left(\sup_{t \in I} |g(t, u(t), I^{\beta_1} u(t), \dots, I^{\beta_n} u(t)) - g(t, v(t), I^{\beta_1} v(t), \dots, I^{\beta_n} v(t))| \right) \\
& = \Omega_1 \left(\sup_{t \in I} |f(t, u(t), {}^c D^{\beta_1} u(t), \dots, {}^c D^{\beta_n} u(t)) - f(t, v(t), {}^c D^{\beta_1} v(t), \dots, {}^c D^{\beta_n} v(t))| \right. \\
& \quad \left. + \sup_{t \in I} |g(t, u(t), I^{\beta_1} u(t), \dots, I^{\beta_n} u(t)) - g(t, v(t), I^{\beta_1} v(t), \dots, I^{\beta_n} v(t))| \right).
\end{aligned}$$

Let $j \in \{1, 2, \dots, n\}$ be given. Then we have

$$\begin{aligned}
& |{}^c D^{\beta_j} T u(t) - {}^c D^{\beta_j} T v(t)| \\
& = \left| \left\{ \frac{1}{\Gamma(q-\beta_j)} \int_0^t (t-s)^{q-\beta_j-1} f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) ds \right. \right. \\
& \quad \left. \left. + \frac{\frac{\alpha_1 t^{1-\beta_j}}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) ds}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \right) \right|
\end{aligned}$$

$$\begin{aligned}
& - \frac{t^{1-\beta_j} \sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) ds \right)}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \Bigg\} \\
& + \left\{ \frac{1}{\Gamma(q-\beta_j)} \int_0^t (t-s)^{q-\beta_j-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \right. \\
& + \frac{\frac{\alpha_1 t^{1-\beta_j}}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \\
& - \frac{t^{1-\beta_j} \sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) ds \right)}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \Bigg\} \\
& - \left\{ \frac{1}{\Gamma(q-\beta_j)} \int_0^t (t-s)^{q-\beta_j-1} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds \right. \\
& + \frac{\frac{\alpha_1 t^{1-\beta_j}}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \\
& - \frac{t^{1-\beta_j} \sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s)) ds \right)}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \Bigg\} \\
& - \left\{ \frac{1}{\Gamma(q-\beta_j)} \int_0^t (t-s)^{q-\beta_j-1} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds \right. \\
& + \frac{\frac{\alpha_1 t^{1-\beta_j}}{\Gamma(q-1)} \int_0^1 (1-s)^{q-2} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \\
& - \frac{t^{1-\beta_j} \sum_{i=1}^n \left(\frac{a_i}{\Gamma(q-\beta_i)} \int_0^1 (1-s)^{q-\beta_i-1} g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s)) ds \right)}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \Bigg\} \\
& \leq \left| \frac{1}{\Gamma(q-\beta_j)} \int_0^t (t-s)^{q-\beta_j-1} |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) \right. \\
& \quad \left. - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \right. \\
& \quad \left. + \left| \frac{\frac{\alpha_1 t^{1-\beta_j}}{\Gamma(q-1)}}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \right| \right. \\
& \quad \times \int_0^1 (1-s)^{q-2} |f(s, u(s), {}^c D^{\beta_1} u(s), \dots, {}^c D^{\beta_n} u(s)) \\
& \quad - f(s, v(s), {}^c D^{\beta_1} v(s), \dots, {}^c D^{\beta_n} v(s))| ds \\
& \quad + \left| \frac{1}{\Gamma(q-\beta_j)} \int_0^t (t-s)^{q-\beta_j-1} |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) \right. \\
& \quad \left. - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds \right. \\
& \quad \left. + \left| \frac{\frac{\alpha_1 t^{1-\beta_j}}{\Gamma(q-1)}}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \right| \right. \\
& \quad \times \int_0^1 (1-s)^{q-2} |g(s, u(s), I^{\beta_1} u(s), \dots, I^{\beta_n} u(s)) - g(s, v(s), I^{\beta_1} v(s), \dots, I^{\beta_n} v(s))| ds
\end{aligned}$$

$$\begin{aligned}
&\leq \left(\left| \frac{1}{\Gamma(q-\beta_j+1)} \right| + \left| \frac{\frac{\alpha_1}{\Gamma(q)}}{\Gamma(2-\beta_j)(\sum_{i=1}^n \frac{a_i}{1-\Gamma(2-\beta_i)} - \alpha_1)(\sum_{i=1}^n (\frac{b_i}{\Gamma(\beta_i+1)}))} \right| \right) \\
&\quad \times \left(\sup_{t \in I} |f(t, u(t), {}^cD^{\beta_1} u(t), \dots, {}^cD^{\beta_n} u(t)) - f(t, v(t), {}^cD^{\beta_1} v(t), \dots, {}^cD^{\beta_n} v(t))| \right. \\
&\quad \left. + \sup_{t \in I} |g(t, u(t), I^{\beta_1} u(t), \dots, I^{\beta_n} u(t)) - g(t, v(t), I^{\beta_1} v(t), \dots, I^{\beta_n} v(t))| \right) \\
&= \Omega_2 \left(\sup_{t \in I} |f(t, u(t), {}^cD^{\beta_1} u(t), \dots, {}^cD^{\beta_n} u(t)) - f(t, v(t), {}^cD^{\beta_1} v(t), \dots, {}^cD^{\beta_n} v(t))| \right. \\
&\quad \left. + \sup_{t \in I} |g(t, u(t), I^{\beta_1} u(t), \dots, I^{\beta_n} u(t)) - g(t, v(t), I^{\beta_1} v(t), \dots, I^{\beta_n} v(t))| \right).
\end{aligned}$$

Thus, we get

$$\begin{aligned}
&d(Tu, Tv) \\
&= \sup_{t \in I} |Tu(t) - Tv(t)| + \sup_{t \in I} |{}^cD^{\beta_1} Tu(t) - {}^cD^{\beta_1} Tv(t)| + \dots \\
&\quad + \sup_{t \in I} |{}^cD^{\beta_n} Tu(t) - {}^cD^{\beta_n} Tv(t)| \\
&\leq \Omega_1 \left(\sup_{t \in I} |f(t, u(t), {}^cD^{\beta_1} u(t), \dots, {}^cD^{\beta_n} u(t)) - f(t, v(t), {}^cD^{\beta_1} v(t), \dots, {}^cD^{\beta_n} v(t))| \right. \\
&\quad \left. + \sup_{t \in I} |g(t, u(t), I^{\beta_1} u(t), \dots, I^{\beta_n} u(t)) - g(t, v(t), I^{\beta_1} v(t), \dots, I^{\beta_n} v(t))| \right) \\
&\quad + n\Omega_2 \left(\sup_{t \in I} |f(t, u(t), {}^cD^{\beta_1} u(t), \dots, {}^cD^{\beta_n} u(t)) - f(t, v(t), {}^cD^{\beta_1} v(t), \dots, {}^cD^{\beta_n} v(t))| \right. \\
&\quad \left. + \sup_{t \in I} |g(t, u(t), I^{\beta_1} u(t), \dots, I^{\beta_n} u(t)) - g(t, v(t), I^{\beta_1} v(t), \dots, I^{\beta_n} v(t))| \right) \\
&= (\Omega_1 + n\Omega_2) \left(\sup_{t \in I} |f(t, u(t), {}^cD^{\beta_1} u(t), \dots, {}^cD^{\beta_n} u(t)) - f(t, v(t), {}^cD^{\beta_1} v(t), \dots, {}^cD^{\beta_n} v(t))| \right. \\
&\quad \left. + \sup_{t \in I} |g(t, u(t), I^{\beta_1} u(t), \dots, I^{\beta_n} u(t)) - g(t, v(t), I^{\beta_1} v(t), \dots, I^{\beta_n} v(t))| \right) \\
&\leq \lambda \left(\sup_{t \in I} |u(t) - v(t)| + \sup_{t \in I} |{}^cD^{\beta_1} u(t) - {}^cD^{\beta_1} v(t)| + \dots + \sup_{t \in I} |{}^cD^{\beta_n} u(t) - {}^cD^{\beta_n} v(t)| \right) \\
&= \lambda d(u, v)
\end{aligned}$$

for all $u, v \in \mathcal{X}$. This implies that T is α -contraction. Let $u, v \in \mathcal{X}$ be such that $\alpha(u, v) \geq 1$. Then $\xi(u(t), {}^cD^{\beta_1} u(t), \dots, {}^cD^{\beta_n} u(t), v(t), {}^cD^{\beta_1} v(t), \dots, {}^cD^{\beta_n} v(t)) \geq 0$. Hence, $\xi(Tu(t), {}^cD^{\beta_1} Tu(t), \dots, {}^cD^{\beta_n} Tu(t), Tv(t), {}^cD^{\beta_1} Tv(t), \dots, {}^cD^{\beta_n} Tv(t)) \geq 0$ for all $t \in [0, 1]$ and so $\alpha(Tu, Tv) \geq 1$. It means that T is α -admissible. Finally, it is easy to check that $\alpha(u_1, Tu_1) \geq 1$. Now by using Lemma 1.2, T has approximate fixed point which is an approximate solution for the problem (2.1). \square

By using Lemma 1.2, one can easily check that the sum-type fractional integro-differential equation (2.1) has at least one exact solution whenever the functions f, g are continuous.

3 Numerical method

In this section, we use the Chebyshev and Legendre polynomials for finding approximate solutions of the problem (2.1). The shifted Chebyshev polynomials be defined

on $[0, 1]$ by $T_{n+1}^*(x) = 2(2x - 1)T_n^*(x) - T_{n-1}^*(x)$ for all $n \geq 1$, where $T_1^*(x) = 2x - 1$ and $T_0^*(x) = 1$ ([24]). The analytical form of the shifted Chebyshev polynomials $T_n^*(x)$ is given by $T_n^*(x) = n \sum_{i=0}^n (-1)^{n-i} \frac{2^{2i}(n+i-1)!}{(2i)!(n-i)!} x^i$ for all $n \geq 1$ ([24]). We have the orthogonality condition $\int_0^1 \frac{T_n^*(x)T_m^*(x)}{\sqrt{x-x^2}} dx = 0$ whenever $m \neq n$, $\int_0^1 \frac{T_n^*(x)T_m^*(x)}{\sqrt{x-x^2}} dx = \frac{\pi}{2}$ whenever $m = n \neq 0$ and $\int_0^1 \frac{T_n^*(x)T_m^*(x)}{\sqrt{x-x^2}} dx = \pi$ whenever $m = n = 0$ ([24]). Every function $u \in L^2([0, 1])$ can be expressed by the shifted Chebyshev polynomials as $u(x) = \sum_{i=0}^{\infty} c_i T_i^*(x)$, where $c_0 = \frac{1}{\pi} \int_0^1 \frac{u(t)T_0^*(t)}{\sqrt{t-t^2}} dt$ and $c_i = \frac{2}{\pi} \int_0^1 \frac{u(t)T_i^*(t)}{\sqrt{t-t^2}} dt$ for all $i \geq 1$ ([22]). Denote the first $(m+1)$ -terms of the shifted Chebyshev polynomials by $u_m(x) = \sum_{i=0}^m c_i T_i^*(x)$ for all $m \geq 1$ ([22]).

Theorem 3.1 Let $\alpha > 0$ be given. Then we have ${}^cD^\alpha(u_m(x)) = \sum_{i=\lceil \alpha \rceil}^m \sum_{k=\lceil \alpha \rceil}^i c_i w_{i,k}^{(\alpha)} x^{k-\alpha}$ and $I^\alpha(u_m(x)) = \sum_{i=0}^m \sum_{k=0}^i c_i \Theta_{i,k}^{(\alpha)} x^{k+\alpha}$, where $\Theta_{i,k}^{(\alpha)} = (-1)^{i-k} \frac{2^{2k} i(i+k-1)! \Gamma(k+1)}{(i-k)!(2k)! \Gamma(k+1+\alpha)}$, $\Theta_{0,0}^{(\alpha)} = \frac{1}{\Gamma(\alpha+1)}$ and $w_{i,k}^{(\alpha)} = (-1)^{i-k} \frac{2^{2k} i(i+k-1)! \Gamma(k+1)}{(i-k)!(2k)! \Gamma(k+1-\alpha)}$.

Proof By using the linear properties of the Caputo fractional derivative, we get

$$\begin{aligned} {}^cD^\alpha(u_m(x)) &= {}^cD^\alpha(c_0 T_0^*(x)) + \sum_{i=1}^m c_i {}^cD^\alpha(T_i^*)(x) \\ &= {}^cD^\alpha(c_0 T_0^*(x)) + \sum_{i=1}^m \sum_{k=0}^i c_i (-1)^{i-k} \frac{2^{2k} i(i+k-1)!}{(i-k)!(2k)!} {}^cD^\alpha(x^k). \end{aligned}$$

Since ${}^cD^\alpha(x^k) = 0$ whenever $k = 0, 1, \dots, \lceil \alpha \rceil - 1$ and ${}^cD^\alpha(x^k) = \frac{\Gamma(k+1)}{\Gamma(k+1-\alpha)} x^{k-\alpha}$ whenever $k \geq \lceil \alpha \rceil$, we have

$${}^cD^\alpha(u_m(x)) = \sum_{i=\lceil \alpha \rceil}^m \sum_{k=\lceil \alpha \rceil}^i c_i (-1)^{i-k} \frac{2^{2k} i(i+k-1)! \Gamma(k+1)}{(i-k)!(2k)! \Gamma(k+1+\alpha)} x^{k-\alpha} = \sum_{i=\lceil \alpha \rceil}^m \sum_{k=\lceil \alpha \rceil}^i c_i w_{i,k}^{(\alpha)} x^{k-\alpha}.$$

Also by using the linear properties of the Riemann-Liouville fractional integral, we get

$$\begin{aligned} I^\alpha(u_m(x)) &= I^\alpha(c_0 T_0^*(x)) + \sum_{i=1}^m c_i I^\alpha(T_i^*)(x) \\ &= I^\alpha(c_0 T_0^*(x)) + \sum_{i=1}^m \sum_{k=0}^i c_i (-1)^{i-k} \frac{2^{2k} i(i+k-1)!}{(i-k)!(2k)!} I^\alpha(x^k). \end{aligned}$$

Since $I^\alpha x^k = \frac{\Gamma(k+1)}{\Gamma(k+1+\alpha)} x^{k+\alpha}$, we obtain

$$\begin{aligned} I^\alpha(u_m(x)) &= \frac{c_0 x^\alpha}{\Gamma(\alpha+1)} + \sum_{i=1}^m \sum_{k=0}^i c_i (-1)^{i-k} \frac{2^{2k} i(i+k-1)! \Gamma(k+1)}{(i-k)!(2k)! \Gamma(k+1+\alpha)} x^{k+\alpha} \\ &= \sum_{i=0}^m \sum_{k=0}^i c_i \Theta_{i,k}^{(\alpha)} x^{k+\alpha}. \end{aligned}$$

This completes the proof. \square

For solving the problem (2.1) by using the Chebyshev method, we approximate $x(t)$ by

$$x(t) \cong \sum_{i=0}^m c_i T_i^*(t). \tag{3.1}$$

By substituting the estimates (3.1) in (2.1) and applying Theorem 3.1, we obtain

$$\begin{aligned}
 & \sum_{i=\lceil q \rceil}^m \sum_{s=\lceil q \rceil}^i c_i w_{i,s}^{(q)} t^{s-q} \\
 &= f\left(t, \sum_{i=0}^m c_i T_i^*(t), \sum_{i=\lceil \beta_1 \rceil}^m \sum_{s=\lceil \beta_1 \rceil}^i c_i w_{i,s}^{(\beta_1)} t^{s-\beta_1}, \dots, \sum_{i=\lceil \beta_n \rceil}^m \sum_{s=\lceil \beta_n \rceil}^i c_i w_{i,s}^{(\beta_n)} t^{s-\beta_n}\right) \\
 &+ g\left(t, \sum_{i=0}^m c_i T_i^*(t), \sum_{i=0}^m \sum_{s=0}^i c_i \Theta_{i,s}^{(\beta_1)} t^{s+\beta_1}, \sum_{i=0}^m \sum_{s=0}^i c_i \Theta_{i,s}^{(\beta_2)} t^{s+\beta_2}, \dots, \right. \\
 &\quad \left. \sum_{i=0}^m \sum_{s=0}^i c_i \Theta_{i,s}^{(\beta_n)} t^{s+\beta_n}\right). \tag{3.2}
 \end{aligned}$$

In equation (3.2) for $t = x_p$ and $p = 0, \dots, m + 1 - \lceil q \rceil$, we obtain

$$\begin{aligned}
 & \sum_{i=\lceil q \rceil}^m \sum_{s=\lceil q \rceil}^i c_i w_{i,s}^{(q)} x_p^{s-q} \\
 &= f\left(x_p, \sum_{i=0}^m c_i T_i^*(x_p), \sum_{i=\lceil \beta_1 \rceil}^m \sum_{s=\lceil \beta_1 \rceil}^i c_i w_{i,s}^{(\beta_1)} x_p^{s-\beta_1}, \dots, \sum_{i=\lceil \beta_n \rceil}^m \sum_{s=\lceil \beta_n \rceil}^i c_i w_{i,s}^{(\beta_n)} x_p^{s-\beta_n}\right) \\
 &+ g\left(x_p, \sum_{i=0}^m c_i T_i^*(x_p), \sum_{i=0}^m \sum_{s=0}^i c_i \Theta_{i,s}^{(\beta_1)} x_p^{s+\beta_1}, \sum_{i=0}^m \sum_{s=0}^i c_i \Theta_{i,s}^{(\beta_2)} x_p^{s+\beta_2}, \dots, \right. \\
 &\quad \left. \sum_{i=0}^m \sum_{s=0}^i c_i \Theta_{i,s}^{(\beta_n)} x_p^{s+\beta_n}\right). \tag{3.3}
 \end{aligned}$$

For calculating the unknowns c_0, \dots, c_m , we consider the roots of $T_{m+1-\lceil q \rceil}^*(t)$ and use the $\sum_{j=1}^n (a_j^c D^{\beta_j} x(1)) = \alpha_1 x'(1)$ and $\sum_{j=1}^n (b_j I^{\beta_j} x(1)) = \alpha_2 x'(0)$. Then we get

$$\sum_{j=1}^n a_j \sum_{i=\lceil \beta_j \rceil}^m \sum_{k=\lceil \beta_j \rceil}^i c_i w_{i,k}^{(\beta_j)} = \alpha_1 \sum_{i=1}^m \sum_{k=1}^i c_i w_{i,k}^{(1)} \tag{3.4}$$

and

$$\sum_{j=1}^n b_j \sum_{i=0}^m \sum_{s=0}^i c_i \Theta_{i,s}^{(\beta_j)} = 0. \tag{3.5}$$

Note that equations (3.3) and (3.4) and (3.5) generate $m + 1$ nonlinear equations which can be solved by using the Newton iterative method. Thus, we can find the unknowns c_0, \dots, c_m and so one can calculate $x(t)$. Similarly, the shifted Legendre polynomials on $[0, 1]$ defined by $L_{n+1}^*(x) = \frac{(2n+1)(2x-1)}{n+1} L_n^*(x) - \frac{n}{n+1} L_{n-1}^*(x)$ for all $n \geq 1$, where $L_0^*(x) = 1$ and $L_1^*(x) = 2x - 1$ ([25]). In fact, $L_n^*(x) = \sum_{i=0}^n (-1)^{n+i} \frac{(n+i)!}{(n-i)!(i)!^2} x^i$ for all $n \geq 1$, $\int_0^1 L_n^*(x) L_m^*(x) dx = 0$ whenever $m \neq n$ and $\int_0^1 L_n^*(x) L_m^*(x) dx = \frac{1}{2m+1}$ whenever $m = n$ ([25]). Every function $u \in L^2([0, 1])$ can be expressed by the shifted Legendre polynomials by $u(x) = \sum_{i=0}^{\infty} c_i L_i^*(x)$, where $c_i = (2i+1) \int_0^1 u(t) L_i^*(t) dt$ for $i \geq 1$ ([25]). Denote the first $(m+1)$ -terms shifted

Legendre polynomials by

$$u_m(x) = \sum_{i=0}^m c_i L_i^*(x). \quad (3.6)$$

By applying a similar proof of Theorem 3.1, one can prove next result.

Theorem 3.2 Let $\alpha > 0$ be given. Then we have ${}^cD^\alpha(u_m(x)) = \sum_{i=\lceil\alpha\rceil}^m \sum_{k=\lceil\alpha\rceil}^i c_i \mathcal{A}_{i,k}^{(\alpha)} x^{k-\alpha}$ and $I^\alpha(u_m(x)) = \sum_{i=0}^m \sum_{k=0}^i c_i \mathcal{B}_{i,k}^{(\alpha)} x^{k+\alpha}$, $\mathcal{A}_{i,k}^{(\alpha)} = (-1)^{i+k} \frac{(i+k)!}{(i-k)!(k)!\Gamma(k+1-\alpha)}$ and

$$\mathcal{B}_{i,k}^{(\alpha)} = (-1)^{i-k} \frac{(i+k)!}{(i-k)!(k)!\Gamma(k+1+\alpha)}.$$

Now, we approximate $x(t)$ by

$$x(t) \cong \sum_{i=0}^m d_i L_i^*(t). \quad (3.7)$$

By using estimates (3.7) in the problem (2.1) and applying Theorem 3.2, we obtain

$$\begin{aligned} & \sum_{i=\lceil q \rceil}^m \sum_{s=\lceil q \rceil}^i d_i \mathcal{A}_{i,s}^{(q)} t^{s-q} \\ &= f \left(t, \sum_{i=0}^m d_i L_i^*(t), \sum_{i=\lceil \beta_1 \rceil}^m \sum_{s=\lceil \beta_1 \rceil}^i d_i \mathcal{A}_{i,s}^{(\beta_1)} t^{s-\beta_1}, \dots, \sum_{i=\lceil \beta_n \rceil}^m \sum_{s=\lceil \beta_n \rceil}^i d_i \mathcal{A}_{i,s}^{(\beta_n)} t^{s-\beta_n} \right) \\ &+ g \left(t, \sum_{i=0}^m d_i L_i^*(t), \sum_{i=0}^m \sum_{s=0}^i d_i \mathcal{B}_{i,s}^{(\beta_1)} t^{s+\beta_1}, \sum_{i=0}^m \sum_{s=0}^i d_i \mathcal{B}_{i,s}^{(\beta_2)} t^{s+\beta_2}, \dots, \right. \\ & \left. \sum_{i=0}^m \sum_{s=0}^i d_i \mathcal{B}_{i,s}^{(\beta_n)} t^{s+\beta_n} \right). \end{aligned} \quad (3.8)$$

Now, we collocate (3.8) at $m+1-\lceil q \rceil$ points x_p ($p = 0, \dots, m+1-\lceil q \rceil$) as

$$\begin{aligned} & \sum_{i=\lceil q \rceil}^m \sum_{s=\lceil q \rceil}^i d_i \mathcal{A}_{i,s}^{(q)} x_p^{s-q} \\ &= f \left(t, \sum_{i=0}^m d_i L_i^*(x_p), \sum_{i=\lceil \beta_1 \rceil}^m \sum_{s=\lceil \beta_1 \rceil}^i d_i \mathcal{A}_{i,s}^{(\beta_1)} x_p^{s-\beta_1}, \dots, \sum_{i=\lceil \beta_n \rceil}^m \sum_{s=\lceil \beta_n \rceil}^i d_i \mathcal{A}_{i,s}^{(\beta_n)} x_p^{s-\beta_n} \right) \\ &+ g \left(x_p, \sum_{i=0}^m d_i L_i^*(x_p), \sum_{i=0}^m \sum_{s=0}^i d_i \mathcal{B}_{i,s}^{(\beta_1)} x_p^{s+\beta_1}, \sum_{i=0}^m \sum_{s=0}^i d_i \mathcal{B}_{i,s}^{(\beta_2)} x_p^{s+\beta_2}, \dots, \right. \\ & \left. \sum_{i=0}^m \sum_{s=0}^i d_i \mathcal{B}_{i,s}^{(\beta_n)} x_p^{s+\beta_n} \right), \end{aligned} \quad (3.9)$$

where x_p ($p = 0, \dots, m+1-\lceil q \rceil$) are roots of the polynomial $P_{m+1-\lceil q \rceil}^*(t)$. Also by substituting equation (3.7), Theorem 3.2 and the conditions $\sum_{j=1}^n (a_j {}^c D^{\beta_j} x(1)) = \alpha_1 x'(1)$ and

$\sum_{j=1}^n (b_j I^{\beta_j} x(1)) = \alpha_2 x'(0)$, we get

$$\sum_{j=1}^n a_j \sum_{i=\lceil \beta_j \rceil}^m \sum_{k=1}^i d_i \mathcal{A}_{i,k}^{(\beta_j)} = \alpha_1 \sum_{i=1}^m \sum_{k=1}^i d_i \mathcal{A}_{i,k}^{(1)} \quad (3.10)$$

and

$$\sum_{j=1}^n b_j \sum_{i=0}^m \sum_{s=0}^i d_i \mathcal{B}_{i,s}^{(\beta_j)} = 0. \quad (3.11)$$

Note that equations (3.9) and (3.10) and (3.11) generate $m+1$ nonlinear equations which can be solved by using the Newton iterative method to obtain the unknown d_0, \dots, d_m . Thus, one can calculate the solution $x(t)$ of the problem. Here, we provide two examples to illustrate our numerical methods. There is much work which provides some methods for numerical solutions of some types fractional differential equations (see [11, 14] and [18]). Our aim is not to introduce a method that can be answered with greater accuracy and speed. The following examples illustrate our main results and we show that numerical approximations could be exact sometimes.

Example 1 Consider the fractional differential equation

$$\begin{aligned} {}^c D^{\frac{3}{2}} x(t) &= [10t + \sin(t)] + \ln(|\sinh(t)| + 1) + \frac{1}{20}(x(t) + {}^c D^{\frac{1}{3}} x(t)) \\ &\quad + [{}^c D^{\frac{1}{2}} x(t) + 0.5] \end{aligned} \quad (3.12)$$

with the boundary conditions ${}^c D^{\frac{1}{2}} x(1) + {}^c D^{\frac{1}{3}} x(1) = x'(1)$ and $I^{\frac{1}{2}} x(1) + I^{\frac{1}{3}} x(1) = x'(0)$. Consider the function $f(t, x_1, x_2, x_3) = [10t + \sin(t)] + \ln(|\sinh(t)| + 1) + \frac{x_1}{20} + [x_2 + 0.5] + \frac{x_3}{20}$, $g(t, x_1, x_2, x_3) = 0$ and $\xi((x_1, x_2, x_3), (y_1, y_2, y_3)) = 1$ whenever $x_2 = 0$ and $y_2 = 0$ almost everywhere and $\xi((x_1, x_2, x_3), (y_1, y_2, y_3)) = -1$ otherwise. Put $n = 2$, $\lambda = 0.9$, $\alpha_1 = \alpha_2 = \alpha_3 = a_2 = b_1 = b_2 = 1$, $\beta_1 = \frac{1}{2}$, $\beta_2 = \frac{1}{3}$ and $q = \frac{3}{2}$. One can check that the problem (3.12) satisfy the conditions of Theorem (2.2), where, thus, the problem (3.12) has an approximate solution. Check Tables 1 and 2 and Figure 1.


One can find the coefficients c_i and d_i by using the explained Chebyshev and Legendre methods as in Table 1. Also, one can find difference of the numerical approximate solutions in Figure 1 and Table 2. Here, we denote the numerical solutions of the Chebyshev and Legendre methods by \tilde{x} and \hat{x} , respectively.

Table 1 Coefficients

<i>i</i>	Coefficient value of Chebyshev method c_i	Coefficient value of Legendre method d_i
1	-13.64912481	-13.89585847
2	32.68664752	32.61068585
3	0.739732907	0.983914814
4	0.123903993	0.188170898
5	0.004270165	0.012299691
6	0.011334926	0.02302969
7	-0.004503085	-0.009980864

Table 2 Differences

t	 $\tilde{x}(t) - \hat{x}(t)$
0.0	4.76063632959267e-13
0.1	3.90798504668055e-13
0.2	3.26849658449646e-13
0.3	2.62900812231237e-13
0.4	1.98951966012828e-13
0.5	1.36779476633819e-13
0.6	7.19424519957101e-14
0.7	8.88178419700125e-15
0.8	5.59552404411079e-14
0.9	1.26121335597418e-13
1	1.84741111297626e-13

Figure 1 Chebyshev and Legendre method.

Example 2 Consider the fractional integro-differential equation

$$\begin{aligned} {}^cD^{\frac{\sqrt{6}}{2}}x(t) &= e^{\cos(t)} + \ln(t+1) + \ln(t^2+1) + \frac{1}{30} \left((t+t^5)x(t) + t^{2c}D^{\frac{1}{2}}x(t) + \frac{\sqrt{t}}{t+1} {}^cD^{\frac{1}{3}}x(t) \right. \\ &\quad + \left(\frac{\sin x(t)}{\sin^2 x(t) + 1} \right) {}^cD^{\frac{\sqrt{3}}{2}}x(t) - \frac{2I^{\frac{\sqrt{3}}{2}}x(t)}{2|I^{\frac{\sqrt{3}}{2}}x(t)| + 3} I^{\frac{1}{2}}x(t) + \frac{\sqrt[5]{t}}{9} I^{\frac{1}{3}}x(t) \\ &\quad \left. - \frac{1}{4} I^{\frac{\sqrt{3}}{2}}x(t) \right) \end{aligned} \quad (3.13)$$

with boundary conditions $-{}^cD^{\frac{1}{2}}x(1) + {}^cD^{\frac{1}{3}}x(1) = 2x'(1)$ and $2I^{\frac{1}{2}}x(1) - 3I^{\frac{\sqrt{3}}{2}}x(1) = x'(0)$. Consider the continuous functions

$$f(t, x_1, x_2, x_3, x_4) = e^{\cos(t)} + \ln(t+1) + \frac{1}{30} \left(tx_1 + t^2x_2 + \frac{\sqrt{t}}{t+1}x_3 + \frac{\sin x(t)}{\sin^2 x(t) + 1}x_4 \right)$$


and $g(t, x_1, x_2, x_3, x_4) = \ln(t^2+1) + \frac{1}{30}(t^5x_1 - \frac{2x_3x_2}{2|x_3|+3} + \frac{\sqrt[5]{t}x_3}{9} - \frac{x_4}{2})$. Define the map $\xi((x_1, x_2, x_3, x_4), (y_1, y_2, y_3, y_4)) = 1$ for all $x_1, \dots, x_4, y_1, \dots, y_4 \in \mathbb{R}$. Put $n = 3$, $\lambda = 0.9$, $a_1 = -1$, $a_2 = 1$, $a_3 = 0$, $b_1 = 2$, $b_2 = 0$, $b_3 = -3$, $\alpha_1 = 2$, $\alpha_2 = 1$, $\beta_1 = \frac{1}{2}$, $\beta_2 = \frac{1}{3}$, $\beta_3 = \frac{\sqrt{3}}{2}$ and $q = \frac{\sqrt{6}}{2}$. Thus by using Theorem 2.2, the problem (3.13) has an exact solution. Check Tables 3 and 4 and Figure 2. We present the coefficients c_0, c_1, \dots, c_6 and d_0, d_1, \dots, d_6 (for $m = 6$) by using the Chebyshev and Legendre methods in Table 3. As one easily sees, the difference of the numerical approximate solutions by the Chebyshev and Legendre methods (which

Table 3 Coefficients

<i>i</i>	Coefficient value of Chebyshev method c_i	Coefficient value of Legendre method d_i
0	25.0832696245301	25.0923764343546
1	-100.27385020664	-100.245026343328
2	-0.023785620114085	-0.0164038330046405
3	-0.045007244673989	-0.0606901595074366
4	-0.0234838615675559	-0.0564629360395388
5	-0.0127366109683591	-0.0258775587930064
6	0.0135562287330535	0.030046706107713

Table 4 Differences

<i>t</i>	$ \tilde{x}(t) - \hat{x}(t) $
0	8.17294676380698E-10
0.1	6.89240664542012E-10
0.2	5.59495560992218E-10
0.3	4.26595647695649E-10
0.4	2.9262992029544E-10
0.5	1.60159885354005E-10
0.6	3.02238234439756E-11
0.7	9.81437153768638E-11
0.8	2.27224461468722E-10
0.9	3.58717500148487E-10
1	4.90501861349912E-10

Figure 2 Chebyshev and Legendre method.

has been provided in Figure 2) is inconsiderable. Denote the numerical solutions of the Chebyshev and Legendre methods by \tilde{x} and \hat{x} , respectively. In Table 4, we show that the difference of the approximate solutions obtained by Chebyshev and Legendre methods is negligible.

4 Conclusions

We first prove the existence of approximate solutions for a sum-type fractional integro-differential problem via Caputo differentiation. By using the shifted Legendre and Chebyshev polynomials, we provide a numerical method for finding solutions for the problem. Also, we give two examples to illustrate our results from a numerical point of view. Our aim is not to introduce a method that can be answered with greater accuracy and speed.

Acknowledgements

The authors would like to thank the esteemed referees for their important comments, which improved the final version of this manuscript. The research of the authors was supported by Azarbaijan Shahid Madani University.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors have made equal contributions. The whole work was carried out, read and approved by the authors.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 19 June 2017 Accepted: 17 October 2017 Published online: 02 November 2017

References

- Reinermann, J: Über Toeplitzsche Iterationsverfahren und einige ihrer Anwendungen in der konstruktiven Fixpunkttheorie. *Stud. Math.* **32**, 209-227 (1969) (in German)
- Yamamoto, Y, Ohtsubo, H: Subspace iteration accelerated by using Chebyshev polynomials for eigenvalue problems with symmetric matrices. *Int. J. Numer. Methods Eng.* **10**(4), 935-944 (1976)
- Khader, MM: Introducing an efficient modification of the variational iteration method by using Chebyshev polynomials. *Appl. Appl. Math.* **7**(1), 283-299 (2012)
- Khader, MM: Introducing an efficient modification of the homotopy perturbation method by using Chebyshev polynomials. *Arab J. Math. Sci.* **18**(1), 61-71 (2012)
- Maleknejad, K, Sohrabi, S, Rostami, Y: Numerical solution of nonlinear Volterra integral equations of the second kind by using Chebyshev polynomials. *Appl. Math. Comput.* **188**, 123-128 (2007)
- Chen, YM, Liu, LL, Sun, L, Sun, XH: Numerical solution of a nonlinear fractional integro-differential equation by using Legendre wavelets. *J. Hefei Univ. Technol. Nat. Sci.* **36**(8), 1019-1024 (2013) (in Chinese)
- Mirzaee, F, Fathi, S: Numerical solution of some class of integro-differential equations by using Legendre-Bernstein basis. *J. Hyperstruct.* **3**(2), 139-154 (2014)
- Yousefi, SA: Numerical solution of Abel's integral equation by using Legendre wavelets. *Appl. Math. Comput.* **175**, 574-580 (2006)
- Zhou, FY, Xu, XY: Numerical approximation of definite integrals by using Legendre wavelets and operator matrices. *Acta Math. Appl. Sin.* **38**(5), 862-873 (2015)
- Aydogan, SM, Baleanu, D, Mousalou, A, Rezapour, Sh: On approximate solutions for two higher-order Caputo-Fabrizio fractional integro-differential equations. *Adv. Differ. Equ.* **2017**, 221 (2017)
- Bhrawy, AH, Zaky, MA: Shifted fractional-order Jacobi orthogonal functions: application to a system of fractional differential equations. *Appl. Math. Model.* **40**(2), 832-845 (2016)
- Jafarian, A, Rostami, F, Golmankhaneh, AK, Baleanu, D: Using ANNs approach for solving fractional order Volterra integro-differential equations. *Int. J. Comput. Intell. Syst.* **10**(1), 470-480 (2017)
- Kumar, D, Singh, J, Baleanu, D: Modified Kawahara equation within a fractional derivative with non-singular kernel. *Therm. Sci.* (2017). doi:10.2298/TSCI160826008K
- Mashayekhi, S, Razzaghi, M: Numerical solution of distributed order fractional differential equations by hybrid functions. *J. Comput. Phys.* **315**, 169-181 (2016)
- Singh, J, Kumar, D, Nieto, JJ: A reliable algorithm for a local fractional Tricomi equation arising in fractal transonic flow. *Entropy* **2016**, 206 (2016)
- Singh, J, Kumar, D, Al Qurashi, M, Baleanu, D: A new fractional model for giving up smoking dynamics. *Adv. Differ. Equ.* **2017**, 88 (2017)
- Singh, J, Kumar, D, Al Qurashi, M, Baleanu, D: A novel numerical approach for a nonlinear fractional dynamical model of interpersonal and romantic relationships. *Entropy* **2017**, 375 (2017)
- Singh, H, Srivastava, HM, Kumar, D: A reliable numerical algorithm for the fractional vibration equation. *Chaos Solitons Fractals* **103**, 131-138 (2017)
- Suganya, S, Baleanu, D, Kalamani, P, Arjunan, MM: A note on non-instantaneous impulsive fractional neutral integro-differential systems with state-dependent delay in Banach spaces. *J. Comput. Anal. Appl.* **20**, 1302-1317 (2016)
- Kilbas, AA, Srivastava, HM, Trujillo, JJ: Theory and Applications of Fractional Differential Equations, vol. 204. Elsevier, Amsterdam (2006)
- Podlubny, I: Fractional Differential Equations. Academic Press, San Diego (1999)
- Miandaragh, MA, Postolache, M, Rezapour, Sh: Some approximate fixed point results for generalized α -contractive mappings. *Sci. Bull. "Politeh." Univ. Buchar. Ser. A, Appl. Math. Phys.* **75**(2), 3-10 (2013)
- Su, X: Boundary value problem for a coupled system of nonlinear fractional differential equations. *Appl. Math. Lett.* **22**, 64-69 (2009)
- Snyder, MA: Chebyshev Methods in Numerical Approximation. Prentice Hall, Englewood Cliffs (1966)
- Khader, MM: Numerical solution of nonlinear multi-order fractional differential equations by implementation of the operational matrix of fractional derivative. *Stud. Nonlinear Sci.* **2**(1), 5-12 (2011)