CORRECTION Open Access



Correction to: Huddling remodels gut microbiota to reduce energy requirements in a small mammal species during cold exposure

Xue-Ying Zhang^{1,2†}, Gansukh Sukhchuluun^{1,2†}, Ting-Bei Bo^{1,2†}, Qing-Sheng Chi¹, Jun-Jie Yang³, Bin Chen³, Lei Zhang^{3,4*} and De-Hua Wang^{1,2*}

Correction

Following publication of the original article [1], the authors reported an error in the caption of Fig. 4.

Originally, the incorrect figure caption was published as: **Fig. 4** Effects of huddling and cold on the concentrations of short-chain fatty acids (SCFAs). The concentrations of acetic acid (**a**), propionic acid (**b**), butyric acid (**c**), isobutyric acid (**d**), valeric acid (**e**), and isovaleric acid (**f**) in huddling and separated Brandt's voles at warm and cold T_a (n = 7-9/g group). *P < 0.05

The correct caption is:

Fig. 4 Effects of huddling and cold on the concentrations of short-chain fatty acids (SCFAs). The concentrations of acetic acid, propionic acid, butyric acid, isobutyric acid, valeric acid, and isovaleric acid in huddling and separated Brandt's voles at warm and cold $T_{\rm a}$ ($n=7-9/{\rm group}$). *P<0.05

Author details

¹State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China. ²University of Chinese Academy of Sciences, Beijing 100049, China. ³College of Life Science, Shandong Normal University, Ji'nan 250014, China. ⁴Microbiome Research Center, Shandong Institutes for Food and Drug Control, Ji'nan 250101, China

Published online: 09 July 2018

Reference

 Zhang, et al. Huddling remodels gut microbiota to reduce energy requirements in a small mammal species during cold exposure. Microbiome. 2018;6:103. https://doi.org/10.1186/s40168-018-0473-9.

[†]Xue-Ying Zhang, Gansukh Sukhchuluun and Ting-Bei Bo contributed equally to this work.



^{*} Correspondence: wangdh@ioz.ac.cn