

Index

Note: The letter *t* and *f* in the index refers to *tables* and *figures* respectively
For example 86*t* refers to *table* in page 86 and 215*f* refers to *figure* in page 215

A

- AAP, *see* American Academy of Pediatrics (AAP)
- Accreditation Council for Graduate Medical Education (ACGME), 33
- ACGME, *see* Accreditation Council for Graduate Medical Education (ACGME)
- Acute pain, definition, 183
- ADHD, *see* Attention-deficit hyperactivity disorder (ADHD)
- ADHD, clinical assessment tools
 - clinical considerations, 252–253
 - primary care toolkit, 252
 - standardized behavior rating scales, 251–252
- ADHD behavior rating scales, 252
- Adolescents with diabetes, 236–238
 - See also* Diabetes
- ADRESSING model, primary care setting, 67
- Adult daily functioning form, 115
- Adult-onset diabetes, 232
 - See also* Diabetes, type 1 diabetes
- Adult primary care, *see* Behavioral screening in adult primary care
- Agency, competing theories of, 56–57
- Agoraphobia, 170–171, 175
- AMA, *see* American Medical Association (AMA)
- American Academy of Pediatrics (AAP), 249, 251, 256, 289, 291, 292, 301
- American Diabetes Association, 231
- American Medical Association (AMA), 32, 125
- American Psychological Association (APA) Practice Directorate, 25
- Ancillary tests, CHD, *see* Coronary heart disease (CHD)
- Anxiety
 - considerations to support PCP, 177
 - consultation/liaison, 174–176
 - Cognitive-behavioral group therapy for social phobia*, 176
 - Cognitive-behavioral therapy for posttraumatic stress disorder*, 176
 - Cognitive-behavioral therapy for social phobia in adolescents: Stand up, speak out*, 176
 - Cognitive-behavioral treatment for generalized anxiety disorder*, 175
 - Compulsive hoarding and acquiring*, 176
 - Emotional processing of traumatic experiences*, 176
 - generalized anxiety disorder, 175
 - Managing social anxiety: A cognitive-behavioral therapy approach*, 176
 - Mastery of your anxiety and worry*, program (MAW-2), 175
 - Obsessive-compulsive disorder, 176
 - Overcoming obsessive-compulsive disorder*, 176
 - Overcoming the trauma of your motor vehicle accident: A cognitive-behavioral treatment program*, 176
 - panic disorder, 175

Anxiety (*cont.*)

- post-traumatic stress disorder, 176
- Prolonged exposure therapy*, 176
- Reclaiming your life from a traumatic experience: A prolonged exposure treatment program*, 176
- Reclaiming your life from PTSD: Teen Workbook*, 176
- social phobia/anxiety disorder, 175–176
- specific phobia, 176
- “Treatments That Work” series, 175
- facts
 - PTSD/GAD/OCD, 169
 - reason to seek help, 171
 - specific/social phobia, 169
- referral/stepped care, 178–179
 - ADAA website, 178
- screening/assessment, 172–174
 - diagnostic screening questionnaires, 172–174
 - IVR technology, 172
 - Mini International Neuropsychiatric Interview for DSM-IV, 172–173
 - PDSR, 173
 - PHQ, 172
 - PRIME-MD, 172
 - Social Phobia Diagnostic Questionnaire, 173
- self-help resources, 177
 - Anxiety disorders: Everything you need to know*, 177
 - generalized anxiety and worry, 178
 - obsessions/compulsions, 178
 - panic and phobias, on, 177–178
 - shyness and social anxiety, 178
 - trauma recovery, 178
- Anxiety disorders: Everything you need to know*, 177
- Anxiety subjective units of distress scale, 110
- APA, *see* American Psychological Association (APA)
- 5A’s approach, smoking, 277
- Atherosclerosis, 261
- Attention-deficit hyperactivity disorder (ADHD)
 - subtypes, 247
- Attention-deficit hyperactivity disorder in primary care
 - algorithm for treatment of school-aged children with ADHD, 255*f*
 - clinical assessment tools, *see* ADHD, clinical assessment tools
 - comorbidity, 249
 - developmental trends, 249
 - DSM-IV Diagnostic Criteria, current
 - hyperactivity, 248
 - impulsivity, 248
 - inattention, 247–248
 - epidemiology, 248–249
 - patient fact sheets/handouts, 249–250
 - handouts for providers/educators, 250
 - practice guidelines
 - publications, 256–258
 - stepped care treatments
 - books/workbooks, 258
 - videos and audio tips, 258–259
 - websites, 259–260
 - treatment manuals and resources for providers, 258

B

- BCP, *see* Behavioral health care providers (BCP)
- Behavioral change, competing theories of, 55–56
- Behavioral Consultation and Primary Care: A Guide to Integrating Services*, 7
- Behavioral health care providers (BCP), 21–26
- Behavioral health consultation for CHD, *see* Coronary heart disease (CHD)
- Behavioral health provider (BHP), 10, 19, 21, 26, 28, 36–39, 53–62, 147, 172, 174, 277, 279, 313
- “Behavioral Health Report Card,” 107
- Behavioral screening in adult primary care
 - characteristics, 125
 - college health centers, screening, 122
 - importance of, 121–122
 - instruments, recommended, 127–129
 - pediatric care, screening, 122
 - screeners, other, 129
 - screening, definition, 123–124
 - positive screen, 124
 - timing, 125–126
 - See also* Screening
- Behavior chain and solutions, 215*f*
- Behavior change
 - stages, 86*f*
 - strategies, stages of, 86*r*
- BHP, *see* Behavioral health provider (BHP)
- Biodyne model of integrated care
 - research-based psychotherapy, 20
- Biopsychosocial (BPS) model
 - training in integrated behavioral health, 78
- BPS, *see* Biopsychosocial (BPS) model

C

- Capitation, 28
- Carcinogen, 277–278
- Cardiology (101), 274
- Case conferences, mistakes in
 - extraneous issues, sidetracked by, 62
 - intimidation of physician, 61
 - making lengthy, 62
 - non-clarification, 61
 - non-involvement of patient, 62
 - office politics, stage for, 62
- Case management, 29
- CBT, *see* Cognitive behavioral therapy (CBT)
- CDC, *see* Center for disease control and prevention (CDC)
- Center for disease control and prevention (CDC), 289, 292
- Center for Epidemiologic Studies Depression Scale (CES-D), 128
- CES-D, *see* Center for Epidemiologic Studies Depression Scale (CES-D)
- CFO, *see* Chief Financial Officer (CFO)
- Chief Financial Officer (CFO), 107
- Child and adolescent clinic, 16
- Cholesterol, pediatric obesity, 290
 - See also* Pediatric obesity
- Chronic obstructive pulmonary disease (COPD), 177
- Chronic pain, 183
 - acute/chronic/widespread pain, definition, 183
 - consultation liaison
 - guiding patients towards pain acceptance, 188–189
 - managing pain-related psychological conditions, 189–190
 - goals of pain acceptance, 188
 - IASP, 183
 - messages, to reduce chronicity, 184
 - primary care setting, in, 183–184
 - screening/assessment, 184–187
 - information, parameters of pain, 184–187
 - over-medicalization, 185
 - self-help resources
 - books, 192–193
 - readings, 193–194
 - websites, 193
 - stepped care approach, 191–192
 - supporting primary care physician's efforts, 190–191
 - symptoms and behaviors, 186–187
- Client Satisfaction Questionnaire-(8), 99
- Cognitive-behavioral group therapy for social phobia*, 176
- Cognitive behavioral therapy (CBT), 37, 103, 176, 311, 313–314
 - See also* Somatization in primary care
- Cognitive-behavioral therapy for posttraumatic stress disorder*, 176
- Cognitive-behavioral therapy for social phobia in adolescents: Stand up, speak out*, 176
- Cognitive-behavioral treatment for generalized anxiety disorder*, 175
- Compulsive hoarding and acquiring*, 176
- Consultative liaison in primary care
 - case conference/treatment team meeting, schedule
 - common mistakes, *see* Case conferences, mistakes in
 - coordinated “team” approach, developing mindset of, 61
 - delineating roles, 60
 - key players, inviting, 60
 - treatment goals, prioritizing, 61
- issues in behavioral change
 - agency, competing theories of, 56–57
 - change, competing theories of, 55–56
 - human suffering, competing theories of, 55
- primary/specialty care consultation, differences, 53–54
- confidentiality includes PCP, 54
- pace, 53–54
- public health approach, emphasis on, 55
- treatment, physician control, 54
- skill sets, 57–59
 - mistakes, willingness to make reasonable, 58–59
 - pathophysiology for dummies, 59
 - psychopharmacology, knowledge of, 59
 - quick, being, 57
 - recommendations, relevant and doable, 58
 - referral question, clarifying and shaping, 58
- COPD, *see* Chronic obstructive pulmonary disease (COPD)
- Core competencies for primary care providers in an integrated clinic, 35†
- Coronary artery disease, *see* Coronary heart disease (CHD)
- Coronary heart disease (CHD)
 - additional resources, 274–275
 - cardiology 101, 274

- Coronary heart disease (*cont.*)
 heart disease guide, 275
 ancillary tests, 263
 AHA, 263
 cardiac stress imaging, 263
 EBCT, 263
 EKG, 263
 ETT, 263
 assessment tools, 263
 atherosclerosis, 261
 behavioral health, 261
 behavioral health assessment, 266
 behavioral treatments, 261
 consultation
 stress diary, 268
 deep breathing, 269
 definition, 261
 diamond forrester method, 263
 diet, exercise, and smoking
 HDL/LDL, 270
 lifestyle habits, 261
 muscle relaxation, progressive
 diet/exercise/smoking, 272–274
 psychological factors, 261
 risk factors, 262
 checklist, 266
 screening, 262–263
- Council on Medical Education (CME), 32
- D**
- Deep breathing, 269
- Department of Defense (DoD), 105
- Depression
 consultation/liaison
 assessment phase, 143–144
 behavioral consultant model, 140–141
 behavioral interventions, 145
 CBT procedures, advantages, 146
 clinical implications, 134
 cognitive strategy interventions,
 145–146
 consultation skills, 146–147
 DISCUS, template, 141
 drugs, associated with, 139
 follow-up visits, 146
 30-minute initial appointment,
 phases of, 143
 primary care consultation,
 elements of, 143
 psycho-education interventions,
 144–145
 treatment of depression, CBT, 141–143
 under-diagnosis/under-treatment,
 reasons, 134
 screening/assessment
 depression screening measures,
 137–138
 developing a team, 136–137
 diagnostic issues, 138–139
 “eighth vital sign,” 137
 self-report measures, 135
See also Primary care consultant toolkit
- Depression in primary care, facts, 133–134
- Depression subjective units of distress
 scale, 111
- DFBC, *see* Diabetes Family Behavior
 Checklist (DFBC)
- Diabetes
 adolescents with
 behavioral family systems therapy, 238
 coping skills training, 237
 ‘diabulimia,’ 236
 Multi systemic family therapy, 238
 adults with, 238–239
 cognitive behavioral strategies, 238
 SMBG, 239
 assessment measures, *see* Diabetes
 assessment measures
 books and workbooks, 242
 children with, 235–236
 primary caregivers of, 236
 computer-assisted intervention
 program, 241
 consultation/liaison, 233–234
 HbA1c%, 234
 definition, 231–232
 diagnosing methods, 231
 FPG, 231
 gestational, 232–233
 GlucoBoy, a blood sugar monitoring
 game, 241
 Health Hero video games series, 241
 practice guidelines, 234–235
 referral and stepped care, 242
 screening and assessment in primary
 care, 233
 self-help resources, 241–242
 symptoms of, 233
 type 1 diabetes, 232
 type 2 diabetes, 232
- Diabetes assessment measures
 Barriers to Diabetes Self-Care Scale, 239
 Diabetes Family Behavior Checklist
 (DFBC), 239
 Measurement of Diabetes Knowledge
 Scales (DKN), 239
 Self Care Inventory (SCI), 239

- Diabetes Family Behavior Checklist (DFBC), 239
- Diabetes Knowledge Scales (DKN), 239
- ‘Diabulimia,’ 236
- Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV)*, 247, 305
- Diamond forrester method, 263
- Diaphragmatic breathing, instructions for, 317–318
- DISCUS, 141
- See also* Primary care consultant toolkit
- DKN, *see* Diabetes Knowledge Scales (DKN)
- DoD, *see* Department of Defense (DoD)
- DSM-III somatization symptoms, 311
- DUKE Health Profile, 128
- Dynamic sizing, 64
- E**
- EBCT, *see* Electron-beam computerized tomography (EBCT)
- “Eighth vital sign,” 137
- EKG, *see* Electrocardiogram (EKG)
- Electrocardiogram (EKG), 263
- Electron-beam computerized tomography (EBCT), 263
- Emotional processing of traumatic experiences*, 176
- ETT, *see* Exercise treadmill testing (ETT)
- “Evidence-based teaching,” 81
- Exercise treadmill testing (ETT), 263
- F**
- Family clinic, 16
- Family therapy, 16
- Fasting plasma glucose test, 232*t*
- Fasting plasma glucose test (FPG), 231
- Fee-for-service (FFS), 21, 28
- Female smokers, 276
- Females with diabetes, 236
- FFS, *see* Fee-for-service (FFS)
- FPG, *see* Fasting plasma glucose test (FPG)
- G**
- GAD, *see* Generalized anxiety disorder (GAD)
- Gastrointestinal problems, 305, 312
- Generalized anxiety disorder (GAD), 169–170, 173, 175, 309
- Gestational diabetes, 232–233
- GlucoBoy, a blood sugar monitoring game, 241
- Glycated hemoglobin test (HbA1c%), 234
- H**
- Handbook of Psychological Assessment in Primary Care Settings, 122
- HARP, *see* Harvard/Brown Anxiety Research Program (HARP)
- Harvard/Brown Anxiety Research Program (HARP), 170
- HDL, *see* High-density lipoprotein (HDL)
- Headaches, 36, 293, 305, 312, 313, 317
- Headaches, gastrointestinal problems, 36, 293, 305, 312, 313, 317
- Health and Behavior Assessment Codes, 24–25
- Health Hero video games series, 241
- Heart disease guide, 275
- High-density lipoprotein (HDL), 270
- HPA, *see* Hypothalamic-pituitary-adrenal (HPA)
- Human suffering, competing theories of, 55
- Hyperactivity, 247–248
- Hyperglycemia and ketoacidosis, very high blood glucose, 233
- symptoms, 233
- Hypoglycemia, very low blood glucose symptoms, 233
- Hypothalamic-pituitary-adrenal (HPA), 91
- I**
- IASP, *see* International Association for the Study of Pain (IASP)
- IBS, *see* Irritable bowel syndrome (IBS)
- Impulsivity, 248
- Inattention, 247–248
- Initiative to Transform Medical Education (ITME), 32
- Insomnia, 171, 277
- ‘Institute of Medicine report’ (IOM, 2001), 13
- Insulin, pediatric obesity, 290
- See also* Pediatric obesity
- Integrated behavioral health care, financial models for, 19–29
- APA Practice Directorate, 25
- BCP, 21
- capitated models, 21–22
- control, 22
- effective function, areas to, 23
- FFS, 21
- “hallway hand-offs,” 21
- Health and Behavior Assessment Codes, 24–25
- health plan, obstacles, 23
- integrated care, aims/definition, 19

- Integrated behavioral health care (*cont.*)
 medical cost offset, 22, 27
 military health system, 25–27
 mixed payment models, 21–22
 models, biodyne/consultative, 19–20
 PCP, 21
 physician-leveraging effect, 21
 reimbursement/risk-sharing,
 for (Melek), 20
 See also Melek's models
 requirements, 23
 RVU, 26–27
 scale, 23
 scope of effort, 23
 time, element of, 22
 US Air Force health system
 planners, 25–26
 VR, 22
- Integrated care
 aims/definition, 31
 benefits of adopting
 gap analysis, 7–10
 See also Integrated health care system,
 characteristics of readiness
 hiring/training
 attitudes, negative signs, 44–45
 attitudes/behaviors, positive, 44
 bootstrap methods, 42
 doctorates vs. sub doctorates, 42
 hire, choice of, 42
 prior experience, type of, 44
 research agenda in training, 45
 skill sets, 41
 integrated care curriculum at University
 of Nevada, Reno, 46–47
 at post-graduate level, training, 47
 pre-graduate training, 45–46
 required readings, 48–50
 two-week training agenda/required
 reading, 47–48
 type of degree/behavioral health
 professional, 42–43
 doctorates vs. subdoctorates, 42
 psychologists vs. other health
 professionals, 42–43
 psychoreligion, 43
 values and prior training, 43
 requirement, 10
- Integrated Care: A Guide for Effective
 Intervention*, 7
- Integrated care curriculum at University of
 Nevada, Reno, 46–47
- Integrated care setting, measurement
 in, 96–99
 adult daily functioning form, 115
 anxiety subjective units of distress
 scale, 110
 Client Satisfaction Questionnaire-(8), 99
 consumer satisfaction, 98
 depression subjective units of distress
 scale, 111
 disease specific, mental/physical, 96–97
 general functioning, 98
 medical services, utilization of, 99
 non-disease-specific, 98
 patient satisfaction, 99
 primary care provider satisfaction survey,
 116–118
 provider satisfaction, 99
 quality of life, 98
 SF- 12v2TM Health Survey Acute Recall,
 98, 104–105, 112–114
 SUDS, 97
- Integrated care settings, medical personnel in,
see Medical personnel in integrated
 care settings
- Integrated health care setting, quality
 improvement, *see* Quality
 improvement, integrated health
 care setting
- Integrated health care system, characteristics
 of readiness, 5–7
 knowledge readiness, 5
 organizational readiness, 5
 physician adoption, 5
 physician adoption, patterns of, 5
 resource readiness, 5
 staff characteristics, 6
- Interactive voice response (IVR) technology,
 anxiety, 172
- 'Intermittent care strategy,' 282
- International Association for the Study of Pain
 (IASP), 183
- Irritable bowel syndrome (IBS), 171, 177
- IVR, *see* Interactive voice response (IVR)
 technology, anxiety
- J**
- Juvenileonset diabetes (or)
 insulin-dependent, 231
See also Diabetes
- L**
- LDL, *see* Low-density lipoprotein (LDL)
- Leveraging, physician, 27
- Low-density lipoprotein (LDL), 270

M

- Malignancies, smoking, 276
- Managing social anxiety: A cognitive-behavioral therapy approach*, 176
- Mastery of your anxiety and worry*, program (MAW-2), 175
- Maternal smoking cessation interventions, 286–287
- Medicaid, 21
- Medical cost offset, 28
- Medical personnel in integrated care settings
 - ACGME, 33
 - ADHD, 36
 - aims of primary care, 31
 - AMA, 32
 - clinical competencies, 37–38
 - CME, 32
 - competencies of, 31–39
 - competency development, 38–39
 - family practitioners, 32
 - formal didactic training, 39
 - ITME, 32
 - knowledge competencies, 35–36
 - NP, 32
 - OB/GYN, 32
 - primary care providers, 31
 - program-level competencies, 38
- Medical terminology, 24
- Medicare, 21
- Melek's models
 - Medicare/Medicaid, capitated, 21
- MI, *see* Motivational interviewing (MI)
- MI, issues in applying
 - ask-provide-ask, 207–208
 - decisional balance, 208–209
 - making the transition, 209–213
 - setting an agenda, 207
 - traps, 208–209
 - using menu of options, 207
 - using rulers, 208
- Mini International Neuropsychiatric Interview for DSM-IV, 172–173
- Motivational interviewing (MI)
 - behavior chain and solutions, 215*f*
 - change plan, creating, 213–215
 - skill building, 213–215
 - change talk, eliciting and reinforcing, 202
 - efficacy of, 201
 - patient-centered approach, 200
 - patient–provider relationship, 201
 - patient's readiness, 213

- patient worksheet, *see* Patient worksheet, MI
- and primary care, 201
- principles of, *see* Principles of MI
- provider references, *see* Provider references, MI
- reasons, usefulness, 201
- sample dialogue integrating MI and skills-building, 217*t*–219*t*
- and skill building, combining, 215–216
- skills of, *see* Skills of MI
- special issues in applying, 207–215

N

- National Initiative for Children's Healthcare Quality (NICHQ), 249
- National Institute of Diabetes and Digestive and Kidney Diseases, (NIDDK), 231, 233
- New World Medical Dictionary*, 31
- NICHQ, *see* National Initiative for Children's Healthcare Quality (NICHQ)
- Nicotine
 - dependence, 277
 - Fagerstrom Test for Nicotine Dependence, 280
 - replacement, 284
 - See also* Smoking
- Nicotine replacement therapy (NRT), 282
- NIDDK, *see* National Institute of Diabetes and Digestive and Kidney Diseases, (NIDDK)
- NP, *see* Nurse practitioners (NP)
- NRT, *see* Nicotine replacement therapy (NRT)
- Nurse practitioners (NP), 32

O

- Obesity, symptoms and signs of conditions associated with, 293*t*
- OB/GYN, *see* Obstetricians/gynecologists (OB/GYN)
- Objective structured clinical exams (OSCE), 82
- Obsessions/compulsions, 178
- Obsessive-compulsive disorder (OCD), 170, 176
- Obstetricians/gynecologists (OB/GYN), 32
- OCD, *see* Obsessive-compulsive disorder (OCD)
- Oregon Healthcare System, 5
- Organization Readiness for Integrating Behavioral Health into Primary Care instrument, 5

- OSCE, *see* Objective structured clinical exams (OSCE)
- Overcoming obsessive-compulsive disorder*, 176
- Overcoming the trauma of your motor vehicle accident: A cognitive-behavioral treatment program*, 176
- Overweight children, 289
See also Pediatric obesity
- P**
- Pain acceptance, 188
- Pain coping/relief model, 188
- Panic Control Treatment (PCT), 175
- Panic Disorder Self-Report (PDSR), 173
- Patient Health Questionnaire (PHQ), 127
- Patient-provider dialogue (sample), brief encounter without and with MI, 210*t*–212*t*
- Patient worksheet, MI
 appointment summary and change plan, 227
 decisional balance exercise, 224
 hints, treatment follow up, 225
 menu of options, 222
 rulers, 223
- PC, *see* Primary care (PC)
- PC model, 16
- PCP, *see* Primary care provider (PCP)
- PCP, keys to supporting
 psycho-education about medication, 147
 scripts for PCP to refer patients, 148
 tailor recommendations, 148
- PCT, *see* Panic Control Treatment (PCT)
- PDSR, *see* Panic Disorder Self-Report (PDSR)
- Pediatric obesity, 289–304
 assessment tools, clinical, 291
 activities, physical, 294
 activity level, 294
 body mass index, 292
 CDC, 292
 insulin resistance, laboratory testing, 295
 parental readiness to change, 296
 pediatric growth charts, 292
 pediatricians, 293
 pediatric practitioners, 292
 psychosocial adjustment, 295
 comorbidity, 290
 psychological, 290
 trends/developmental, 290
 definition, clinical, 289
 body mass index, 289
 Center for disease control and prevention (2007), 289, 292
 overweight children, 289
 epidemiology, 289
 African-American, 289
 childhood obesity, 289
 Mexican-American, 289
 native American, 289
 pediatric obesity, 289
 sedentary lifestyles, 289
 patient fact sheets/handouts, 290
 children/adolescents, 290
 tools for providers, 291
 practice guidelines
 clinical, 296
 obesity prevention, 296
 recommendation/implementation, 297
 treatment manuals
 clinical guides, 301
 handbooks, 300
 treatments, stepped care
 books, for parents, 301
 education and treatment websites, 301
 online assessment resources, 301
 other tools, 302
 websites, school/community-based action, 302
 workbooks, 301
- Pharmacotherapy
 chronic pain, in, 184
 evidence-based medicine for mental health, 35
 mood disorders and opioid medication, for, 190
 primary care settings, in, 149
 somatization, for, 310
 tobacco cessation treatment, for, 282
- PHQ, *see* Patient Health Questionnaire (PHQ)
- PHQ for anxiety, 172
- PHQ-2/PHQ-9/BDI-2/GDS, depression screening measures, 137
- Physician-leveraging effect, 21
- Post-traumatic stress disorder (PTSD), 169–170, 176, 307
- PPO, Preferred provider organizations (PPO)
- Preferred provider organizations (PPO), 21, 29
- Primary care consultant, toolkit, *see* Toolkit, primary care consultant
- Primary care consultant toolkit
 ABCs of stress management, 162
 behavioral health consultation, 167
 changing alarming self-talk, 165–166
 reassuring thoughts, 164

- thinking and stress, 160–161
 - 4 types of alarming thoughts, 163
 - Primary Care Evaluation of Mental Disorders (PRIME-MD), 127, 172
 - Primary care model, components of, 14*f*
 - Primary care (PC), 13
 - Primary care provider (PCP), 21
 - Primary care provider satisfaction survey, 116–118
 - Primary care setting
 - administrators knowledge, 13–17
 - administrator, choice of model, 16–17
 - clinical psychology services, integrated, 13
 - components of primary care model, 14*f*
 - ‘Institute of Medicine report’ (IOM, 2001), 13
 - patient access and costs
 - anecdotal study, reports, 14–15
 - PC model, 16
 - PC model, components of, 14*f*
 - psychotropic medications, 13
 - quality treatment/satisfaction/collaboration, 15–16
 - reducing costs, 15
 - cultural competency in
 - 10 clinical considerations in addressing cultural differences, 69
 - culturally competent practice, importance of, 65
 - dynamic sizing, 64
 - human diversity, 63
 - pain, common medical symptom, 65–66
 - race and ethnicity discussion, recommendations, 70
 - resources for PCP/behavioral health consultants, 70–74
 - books, 71–72
 - journals, 73–74
 - websites, 74
 - screening and assessment, 66–70
 - ADRESSING model, 67
 - Micro aggressions, 67–68
 - self-assessment results, 67
 - PRIME-MD, *see* Primary Care Evaluation of Mental Disorders (PRIME-MD)
 - Principles of MI
 - discrepancy, developing, 201
 - empathy, expressing, 201
 - provider reference, 229
 - rolling with resistance, 202
 - self-efficacy, supporting, 202
 - Productivity adjustments, 21, 28
 - Prolonged exposure therapy*, 176
 - Provider references, MI
 - decisional flowchart, 228
 - principles/skills of, 229
 - Psychological treatments, chronic pain, 190
 - Psychopharmacology, 24, 43, 46, 48, 59, 90
 - Psychotropic medications, 13, 34, 37, 59, 104, 122
 - PTSD, *see* Post-traumatic stress disorder (PTSD)
- Q**
- Quality improvement, integrated health care setting, 100*f*
 - data, using
 - individual-level, 101–104
 - system-level, 104–105
 - history of, 95
 - implementation, quality improvement process
 - resistance, overcoming, 105–106
 - resource limitations, overcoming, 106–107
 - solutions, 107–109
 - ingredients for creating quality improvement process, 95–96
 - measurement, *see* Integrated care setting, measurement in setting, 96
 - integrated care setting, 95–96
 - process, 99–101
 - psychological instruments, traditional vs. used in integrated care setting, 97*t*
- R**
- Readiness assessment, integrating behavioral health care, 8–9
 - Readiness assessment system, determination of need
 - adopting integrated care, benefits of gap analysis, 7–10
 - characteristics, 5–7
 - first vital steps issues, 4
 - integrated care, 5
 - Organization Readiness for Integrating Behavioral Health into Primary Care instrument, 5
 - lessons learned, 4–5
 - Oregon Healthcare System, 5
 - possibility of integrated health care, 3–4
 - goals, 3
 - implementation, 3
 - readiness assessment, 4

- Readiness assessment (*cont.*)
 readiness to adopt, 3
- Reclaiming your life from a traumatic experience: A prolonged exposure treatment program*, 176
- Reclaiming your life from PTSD: Teen Workbook*, 176
- Referral and stepped care
 anxiety in primary care, 178–179
 self-management of diabetes, 242
 smokers, special populations of, 285–286
- Relative value units (RVU), 26, 29
- Risk adjustments, 28
- RVU, *see* Relative value units (RVU)
- S**
- Sample contract, diabetes, 237*t*–238*t*
- Sample of a blood glucose monitoring form for patients, 240*t*
- School-aged children with ADHD, algorithm for treatment of, 255*f*
- SCI, *see* Self Care Inventory (SCI)
- Screening
 anxiety in primary care, 172–174
 chronic pain in primary care setting, 184–187
 coronary heart disease, for, 262–263
 cultural competence in primary care setting, 66–70
 depression in primary care, 135–139
 diabetes in primary care, 233
 instruments, behavior in adult primary care, 121–129
 DUKE Health Profile, 128–129
 PHQ, 127
 PRIME-MD, 127–128
 smoking in primary care setting, for, 276–282
 somatization in primary care, 308–309
- Secondhand smoke, 276
 infants, exposed to, 276
- “See one, do one, teach one,” approach to teaching in medicine, 79
- Self Care Inventory (SCI), 239
- Self help resources
 anxiety, patients struggling with, 177
 chronic pain, 192
 diabetes, individuals with, 241
 managing depression, 150
 smokers who wish to quit, for, 284
- Self-monitoring of one’s blood glucose (SMBG), 239
- SF-12v2TM Health Survey Acute Recall, 112–114
- Shyness and social anxiety, 178
- Skills of MI
 affirmations, 203
 open questions, 203
 reflective listening, 203–204
 summaries, 204
- SMBG, *see* Self-monitoring of one’s blood glucose (SMBG)
- Smoking, 277–287
 5A’s approach, 277
 consultation/liaison, 284–285
 care strategy, 282
 facts, 275–276
 carcinogen, 277–279
 malignancies, 276
 secondhand smoke, 276
 substance abuse and mental health services administration (2005), 276
 intensive counseling, 283
 links/tools for behavioral health providers, 285
 minimal counseling intervention, 283
 nicotine dependence
 cessation of nicotine, 277
 impaired concentration, 277
 restlessness, 277
 tolerance to nicotine, 277
 referral/stepped care, 286–287
 nicotine replacement, 284
 5R’s motivational intervention, 279
 screening, 5A’s approach, 278–284
 advise each smoker, 278
 arranging follow-up, 282
 asking each patient, 277
 assess each smoker, 278
 assist the patient, 283
 BHP, 277
 patient’s motivation, enhancing, 279
 PCP, 277
 smoking cessation, 276
 self-help resources, to quit, 285–286
 5-Stage trans-theoretical model of behavior change, 279
- Social anxiety disorder, 175–176
- Social Phobia Diagnostic Questionnaire, 173
- Social phobia disorder, 175–176
- Somatic Symptom Index (SSI), 311
- Somatization in primary care, 305–320
 assessment instruments, 310–311
 patient health questionnaire-15, 310
 somatic symptom index, 311
 symptom checklist, 311
 assessment of, 307

- cognitive behavioral therapy
 - behavioral activation, 313–314
 - cognitive restructuring, 313–314
 - relaxation training, 312–313
 - definition, of, 305–306
 - diaphragmatic breathing, instructions for, 317–318
 - evidence-based treatment, 311
 - facts, basic
 - comorbidity, 306–307
 - health care-seeking/utilization, 307
 - sociodemographic, 306
 - patient health questionnaire-(15), 317
 - pharmacotherapy, 312
 - provider consultation, 312
 - provider resources, 317–318
 - retribution training, 313
 - self-help treatment resources, 316–317
 - self-report rating scale – relaxation, 318
 - somatization, definition, 305–306
 - DSM-IV criteria, 306
 - headaches, gastrointestinal problems, 305
 - mental disorders, 305
 - somatization disorder, 306
 - Specialty behavioral health care, 19–20, 26, 28, 147
 - SSI, *see* Somatic Symptom Index (SSI)
 - STAI, *see* State Trait Anxiety Inventory (STAI)
 - State Trait Anxiety Inventory (STAI), 128
 - Statistical Manual of Mental Disorders (DSM-IV), 24
 - Stress management
 - HPA, research on, 91
 - stress, definition, 91
 - stress/coping, questions to assess, 91–92
 - training to patients, 92
 - Subject units of distress scales (SUDS), 97
 - SUDS, *see* Subject units of distress scales (SUDS)
- T**
- Toolkit, primary care consultant
 - behavior in medicine, 77–78
 - biomedical/biopsychosocial model, 78
 - training, reason for, 78
 - teaching and medical culture, 78–80
 - “curbside consult” and precepting sessions, 81
 - designing educational program, 80–81
 - “Evidence-based teaching,” 81
 - pedagogy, 81–82
 - “see one, do one, teach one,” 79
 - teaching evaluations, 82
 - teaching examples, content/process
 - behavior change, stages/strategies, 85–87
 - building relationship/collaborative environment, tips for, 83–84
 - meta-cognitive skills, 87–89
 - PCPs, work of, 82–83
 - PPI skills, 84–85
 - psychiatric disorders, 89–90
 - stress management, 91–92
 - Trauma recovery, 178
 - “Treatments That Work” series, 175
 - Type 1 diabetes, 232
 - Type 2 diabetes, 232
- U**
- US Air Force health system planners, 25
 - US Preventive Service Task Force, 135
- V**
- Veterans administration (VR), 22
 - VR, *see* Veterans administration (VR)
- W**
- Weight loss targets, 298
 - Widespread pain, 183, 191
 - “Worry times”, scheduling, 92