

# Index

## A

- Acute bacterial prostatitis, 289, 291
- Adenoma
  - adrenal cortical, 116, 127
    - CT, 124, 125, 128
    - MRI sequence, 124, 125
  - myelolipoma, 124, 125
- Adenomatoid tumour, 304, 305
- ADPKD. *See* Autosomal dominant polycystic kidney disease
- Adrenal gland
  - adrenal metastasis
    - contrast-enhanced axial CT, 145
    - FDG-PET/CT image, 144, 145
    - T2-weighted MR image, 144, 145
  - arterial supply and venous drainage, 121, 123
  - carcinoma
    - FDG-PET/CT image, 140–142
    - histological features, 287, 288
    - MRI sequence, 140, 142
    - phase contrast-enhanced CT image, 140, 142
  - ganglioneuromas
    - CT image, 134, 136
    - occurrence, 134, 136
    - pathological assessment, 134–136
  - hyperaldosteronism
    - causes of, 131, 133
    - Conn's adenoma and bilateral hyperplasia, 131–133
    - diagnosis, 131, 133
    - H&E stain, Conn's adenoma, 131–133
  - layers of, 121–123
  - lipid-poor adrenal cortical adenomas, 126, 127
  - lipid-rich adenoma
    - adrenal cortical adenomas, 116, 127
    - CT, 124, 125, 128
    - MRI sequence, 124, 125
    - myelolipoma, 124, 125

- myelolipoma
  - axial non-contrast CT, 128–130
  - complications of, 128, 130
  - composition of, 128, 130
  - MRI sequence, 128–130
- phaeochromocytoma
  - CT and T2-weighted MRI, 137, 139
  - histological features, 137–139
  - radionuclide-labelled MIBG, 137–139
  - size, 121–123
- Adrenal metastasis
  - contrast-enhanced axial CT, 145
  - FDG-PET/CT image, 144, 145
  - T2-weighted MR image, 144, 145
- Angiomyolipoma (AML), 56–58
- Askin sarcoma, 157, 158
- Atypical adenomatous hyperplasia (AAH), 248–250
- Atypical small acinar proliferation (ASAP), 248, 250
- Autosomal dominant polycystic kidney disease (ADPKD), 45, 47

## B

- Balanitis xerotica obliterans (BXO), 358–361
- Benign prostatic hyperplasia (BPH), 231, 232, 242, 245, 282, 284–286
- Benign viral warts, 381
- Bladder
  - anatomical relationship, 175–177
  - calcified mass, 223–224
  - cancer, 201, 203
  - 'cobra head' sign, 199–200
  - CT urogram, 185–187
  - cystitis, 228–230
  - cystitis cystica, 205–206, 209
  - cystitis follicularis, 205–206, 209
  - detrusor overactivity, 178, 180
  - detrusor pressure, 178–180
  - E. coli*, 225–226

**Bladder (cont.)**

- embryology, 175, 177
  - exstrophy, 190–191
  - extra-genitourinary associations, 192–193
  - fistula, 216–219
  - function of, 178, 180
  - haematuria, 201–204
  - haematuria and dysuria, 223–224
  - Hutch diverticulum, 196–198
  - innervation of, 181, 182
  - inverted papilloma, 205–209
  - KUB X-ray, 223–224
  - malakoplakia, 205–209
  - MRI urography, 185–187
  - pathology, 178–180
  - PBS, 192–193
  - PUNLMP, 205–209
  - PUV, 194–195
  - rupture, 220–222
  - stress urinary incontinence, 178, 180
  - telangiectasia, 181, 182
  - trabeculation, 181, 182
  - trauma, 220–222
  - ultrasound, 183–184
  - urachal anomalies, 188–189
  - ureterocele, 199–200
  - UTI, 225–226
  - wall structure, 175–177
- Bladder exstrophy, 190–191
- Bone scintigram, 270, 273
- Brachytherapy, 279, 281
- Bulbar urethral stricture, 347–349

**C**

- Carcinoma of the Prostate (CAP), 257, 258, 260
- Cavernosography, 377–378
- Cobra head sign, 199–200
- Computed tomography (CT)
  - adenoma, 124, 125, 128
  - fibrosis, 172, 174
  - ganglioneuromas, 134, 136
  - inguinal cannal, 330, 331
  - leiomyosarcoma, 151, 152
  - lipid-rich adenoma, 124, 125, 128
  - myelolipoma, 128–130
  - nodal staging, 326, 328
- Congenital adrenal hyperplasia (CAH), 306–308
- Conn's adenoma, 131–133
- Cryptorchidism, 362–363
- Cystic transformation, 309, 310
- Cystitis, 228–230
- Cystitis cystica, 205–206, 209
- Cystitis follicularis, 205–206, 209

**D**

- Detrusor pressure, 389–390
- Detrusor sphincter dyssynergia pattern, 403–404
- Digital rectal examination (DRE), 234, 236
- Digital single lens reflex (DSLRL), 16
- Dimercaptosuccinic acid (DMSA), 11
- Doppler image
  - hypervascular, 322, 324
  - intratesticular varicocele, 320, 331
  - mediastinum testis, 320, 331
- Dynamic sentinel lymph node biopsy, 374–376
- Dysuria, 223–224

**E**

- Emphysematous pyelonephritis, 43–44
- Endometriosis, 84–85
- Epididymitis, 298–300
- Ewing's sarcoma, 157, 158
- Extraprostatic tissue (EPT), 257, 258, 260

**F**

- Fibrosis
  - aetiological factors, 172, 174
  - CT and nephrostogram image, 172, 174
  - idiopathic RPF, 172–174
  - RPF complication, 172–174
- Fibrous septa, 295, 297
- Fistula, 216–219
- Fournier's gangrene, 382–384

**G**

- Ganglioneuromas
  - CT image, 134, 136
  - occurrence, 134, 136
  - pathological assessment, 134–136
- Genitourinary tuberculosis, 342, 343
- Gleason sum score, 254–256
- Glomerular filtration rate (GFR), 8, 11
- Gonococcal urethritis, 350–351
- Granulomatous prostatitis, 292, 391

**H**

- Haematuria, 201–204, 223–224
- High-grade prostatic intra-epithelial neoplasia (HGPIN), 237, 239, 248, 249
- Histopathological image
  - aim, 13
  - governance and storage, 18–19

- operative surgical specimen
    - bivalved orchidectomy, 15
    - examination and fixation, 13
    - hesitant cut, 14
    - nephrectomy, 14
    - ureterectomy, carcinoma, 15
    - Wegener's granulomatosis, 16
  - photography technique, 18
  - tissue and photography, 17–18
  - uses, 13
  - Horseshoe kidney, 94–96
  - Hutch diverticulum, 196–198
  - Hydrocele
    - anechoic fluid collection, 315, 317
    - inflammatory debris, 315–317
    - treatment, 315, 317–319
    - tunica vaginal, 315–317
  - Hyperaldosteronism
    - causes of, 131, 133
    - Conn's adenoma and bilateral hyperplasia, 131–133
    - diagnosis, 131, 133
    - H&E stain, Conn's adenoma, 131–133
  - Hypertrophic column of Bertin, 97–98
  - Hypospadias, 362–363
- I**
- Immunohistochemistry, 237, 239
  - Inferior vena cava leiomyosarcoma.
    - See* Leiomyosarcoma
  - Inverted papilloma:, 205–209
- K**
- Kidneys and ureters
    - aetiology, 43–44
    - angiomyolipoma, 56–58
    - autosomal dominant polycystic kidney disease, 45, 47
    - cortico-medullary phase, 25–27
    - CT KUB, 25–27
    - CT protocols, 25–27
    - CT urogram, 25–27
    - emphysematous pyelonephritis, 43–44
    - endometriosis, 84–85
    - grading system, 28, 31
    - horseshoe kidney, 94–96
    - hyperechoic cortex, 21–24
    - hypertrophic column of Bertin, 97–98
    - imaging modality, 25–27
    - intermittent flank pain, 59–60
    - macroscopic haematuria, 51–52
    - multicystic dysplastic kidney, 89–90
    - nephrocalcinosis, 73–76
    - nephrographic phase, 25–27
    - non-specific chronic back pain, 56–58
    - oncocytoma, 59–60
    - orientation, 21–24
    - papillary necrosis, 33–35
    - pelvicureteric obstruction, 33–35
    - post-renal transplant complications, 117–118
    - primary megaureter, 102–103
    - pyeloureteritis cystica, 38–39
    - radiolucent stones, 77–79
    - renal artery aneurysm, 106–108
    - renal cystic lesions, 45–47
    - renal infarction, 104–105
    - renal lymphoma, 70–72
    - retrocaval, 102–103
    - staghorn calculus, 82–83
    - transitional cell carcinoma, 53–55
    - tuberculosis, 40–42
    - ureteric calculus, 77–79
    - von Hippel Lindau (VHL) disease, 45–47
    - XGP, 33–37
- L**
- Leiomyosarcoma
    - central low attenuation area significance, 151, 152
    - CT image, 151, 152
    - gender predilection, 153, 154
    - hydronephrosis, 153, 154
    - inferior vena cava, 153, 154
    - metastasis frequency, 151, 152
    - venous leiomyosarcomata, 153, 154
  - Lichen sclerosis. *See* Balanitis xerotica obliterans (BXO)
  - Lower urinary tract symptoms
    - detrusor overactivity, 421–422
    - flow rate, 415–416, 425–426
    - flow signal, 421–422
    - obstruction/hypocontractile detrusor, 427–428
    - overactive bowel contractions, 421–422
    - pinching, 427–428
    - spike artefact, 415–416
    - urethral stricture, 425–426
    - uroflow study, 423–424
  - Lymphoedema, 364–365
  - Lymphoma
    - chronic lymphocytic leukaemia, 167, 169
    - retroperitoneal adenopathy, 167, 169
- M**
- Magnetic resonance imaging (MRI)
    - adenoma, 124, 125
    - adrenal gland, 140, 142

Magnetic resonance imaging (MRI) (*cont.*)

- bladder, 185–187
- importance, 302, 303
- lipid-rich adenoma, 124, 125
- myelolipoma, 128–130
- phaeochromocytoma, 137, 139
- prostate, 240, 241
  - abnormality, 240, 241
  - central gland, 242, 245
  - CT, 231, 232
  - diffuse abnormal low signal, 292, 293
  - image, feature, 234, 236
  - low signal, 254–256, 257, 260
  - marked improvement, 289–291
  - prostatic mass, 277, 278
  - rectoprostatic angle, 257, 260
  - right posterior PZ, 251, 253
  - staging, 234, 236
    - T2 and diffusion weighted, 257, 259, 261
  - testes and adnexae, 302, 303
  - undescended testis, 302, 303
  - urological disorder, 8, 10
  - uses of, 302, 303

## Malakoplakia, 205–209

## Malignant peripheral nerve sheath tumour (MPNST), 159, 160

## Marked hyperaemia, 298–300

## Meatal stenosis, 358, 360

## Mercaptoacetyltriglycine (MAG), 11

## Metaiodobenzylguanidine (MIBG), 137–139

## Micturating cystourethrogram (MCUG), 194–195

MPNST. *See* Malignant peripheral nerve sheath tumour

## Multicystic dysplastic kidney, 89–90

## Myelolipoma

- axial non-contrast CT, 128–130
- complications of, 128, 130
- composition of, 128, 130
- MRI sequence, 128–130

## N

## Nephrocalcinosis, 73–76

## Nocturia and occasional urge incontinence

- detrusor overactivity, 387–388
- leakage, 387–388
- voiding phase, 387–388

## Non-seminomatous germ cell tumour

- (NSGCT), 326, 328, 329
- axial CT, 163, 164
- metastasis, 16, 163
- orchidectomy relapse rate, 163, 164
- post-chemotherapy, 163, 164

## O

## Oncocytoma, 59–60

## Orchiectomy specimen

- transverse section, 330, 332
- white fibrous mas, 330, 332

## P

## Papillary necrosis, 33–35

## Papillary Urothelial Neoplasm of Low Malignant Potential (PUNLMP), 205–209

## Paraganglioma

- benign pathology, 155, 156
- malignancy, 155, 156
- organs of Zuckerkandl, 155, 156
- radionuclide tracer, 155, 156
- sign and symptoms, 155, 156

## Pelvicutereic obstruction, 33–35

## Penile Doppler study, 379–380

## Penis

- arteriography, 379–380
- benign viral warts, 381
- Bowenoid papulosis, 366, 368
- bulbar urethral stricture, 347–349
- cancers
  - risk factors, 370, 372
  - staging of, 370, 372
  - subtypes, 370–373
- cavernosography, 377–378
- cryptorchidism, 362–363
- dynamic sentinel lymph node biopsy, 374–376
- Fournier's gangrene, 382–384
- fractured, 355–357
  - surgical exploration, 355–357
  - tunical rupture, 355–357
  - urethral distraction injury, 355–357
- glans resurfacing, 358–361
- hypospadias, 362–363
- intra-epithelial neoplasia (IN)
  - keratin production, 366, 368
  - treatment, 366, 368
- lichen sclerosis, 358–361
- lymphoedema, 364–365
- meatal stenosis, 358, 360
- palpable lymph nodes, 374–376
- penile Doppler study, 379–380
- Peyronie's disease, 377–378
- reduced maximum flow rate, 347–349
- trauma
  - antegrade urethrogram, 352–354
  - diastasis, 352, 354
  - management and investigation, 352, 354

- pubic ramus fracture, 352, 354
- transection, 352–354
- urethrography, 347–349
  - epithelial irregularity, 350–351
  - gonococcal urethritis, 350–351
- urethroplasty, 347–349
- Zoon's balanitis, 366, 368
- Peyronie's disease, 377–378
- Phaeochromocytoma
  - CT and T2-weighted MRI, 137, 139
  - histological features, 137–139
  - radionuclide-labelled MIBG, 137–139
- Picture archiving and Communication Systems (PACS), 2
- Posterior urethral valves (PUV), 194–195
- Primary megoureter, 102–103
- Primitive neuroectodermal tumour (PNET)
  - coronal and axial CT, 157, 158
  - Ewing's sarcoma and Askin sarcoma, 157, 158
  - prognostic factor, 157, 158
- Prostate
  - AAH, 248, 249
  - abnormal intermediate signal, 282–284
  - acinus, 237, 239
  - acute bacterial prostatitis, 289, 291
  - adenocarcinoma cell, 270, 272, 273
  - advanced disease, 274, 276
  - ASAP, 248, 250
  - basal cell, 237, 239
  - biopsy core, 251, 253
  - bone metastases and lymphadenopathy, 266, 267
  - bone scintigram, 270, 273
  - bone turnover, 270, 273
  - brachytherapy, 279, 281
  - CAP, 257, 258, 260
  - clinical stage
    - T4N1 M1b, 266, 267
    - T4N1 M1c, 268, 269
  - compare with normal gland, 248, 249
  - cystic hyperplastic nodule, 242, 243, 245
  - differential diagnosis, 240, 241, 262, 264
  - EPT, 257, 258, 260
  - extensive retroperitoneal lymphadenopathy, 274–276
  - extraprostatic extension, 257, 260
  - fat-suppressed T1 weighted, 268, 269
  - Gleason grading system
    - glandular micro-architecture, 254, 256
    - pattern, 254, 256
  - granulomatous prostatitis, 292, 293
  - H, E and IH preparation, 248–250
  - hip, explanation, 266, 267
  - histological feature, 287, 288
  - hyperplasia
    - morphology, 242–245
    - nodular, 246, 247
  - immunohistochemical technique, 270, 272, 273
  - immunohistochemistry, 236, 238
  - invasive cancer, 233, 253
  - leiomyosarcoma, 285, 286
  - magnetic resonance imaging (MRI)
    - abnormality, 240, 241
    - central gland, 242, 245
    - CT, 231, 232
    - diffuse abnormal low signal, 292, 293
    - image, feature, 234, 236
    - low signal, 254–256, 257, 260
    - marked improvement, 289–291
    - prostatic mass, 277, 278
    - rectoprostatic angle, 257, 260
    - right posterior PZ, 251, 253
    - staging, 234, 236
    - T2 and diffusion weighted, 257, 259, 261
  - microscopic feature, 237, 239
  - necrosis, 246, 247
  - non-adenocarcinoma, 285, 286
  - non-caseating granuloma, 292, 293
  - pathological and clinical significance, 248, 249
  - pelvic node, 279–281
  - PET CT image, 282–284
  - positive surgical margin, 257, 258, 260
  - post-mortem coronal section, 274–276
  - prostatic adenocarcinoma, 277, 278
  - radical prostatectomy (RP)
    - coronal macroscopic section, 251–253
    - specimen, 298–300
  - radical radiotherapy, 279, 281
  - radiological feature, 266, 267
  - radiological staging, 251, 253
  - right external iliac node, 277, 278
  - right side of, 234–236
  - sagittal T2 and T1 weighted, 270, 271, 273
  - significant pathology, 266, 267
  - stroma, 237, 239
  - structure
    - axial and coronal high-resolution T2-weighted image, 231, 232
    - axial CT image, 231, 232
    - surgical resection specimen, 242, 243, 245
  - T2c prostate cancer, 254–256
  - transrectal ultrasound (TRUS)
    - abnormality, 240, 241
    - biopsy, 236, 236, 287, 288

- Prostate (*cont.*)
- echogenicity, 254, 256, 257, 260
  - irregular cavity, 287, 288
  - thinned PZ, 242, 243, 245
  - tumour stage, 262–264
  - T2 signal
    - low, 257, 260, 279, 281
  - TUR defect, 268, 269
  - ultrasound image, 234, 236
  - utricle cysts, 240, 241
  - well differentiated carcinoma, 248, 249
  - zones of, 231, 232
- Prostatic adenocarcinoma, 277, 278
- Prune belly syndrome (PBS), 192–193
- Pyeloureteritis cystica, 38–39
- R**
- Radical prostatectomy (RP)
- coronal macroscopic section, 251–253
  - specimen, 298–300
- Radical radiotherapy, 49, 51
- Radiological image
- computed tomography (CT), 3
  - contrast media
    - chemotoxic effect, 10
    - limitation, 9
    - MRI, 11
    - signal–noise ratio, 8
    - ultrasound, 10–11
  - contrast nephropathy, 10
  - hazards
    - effective radiation dose, 12
    - stochastic effect, 12
    - X-ray interact, 11
  - hounsfield or CT Unit, 3
  - magnetic resonance imaging (MRI)
    - physics, 7
    - signal, 8
    - T2 weighted, 9
    - urological disorder, 8, 10
  - nuclear medicine, 11
  - plain, 2
  - tissue–energy interaction, 1
  - ultrasound
    - grey–scale, 7
    - moving interface, 6
    - urological disorder, 6
  - window level and width
    - renal–specific protocol and CT technique, 4–5
    - screen setting, 3
    - urological disorder, 5
- X-rays
- PACS, 2
  - production, 2
- Radiological staging, 251, 253
- Radiolucent stones, 77–79
- Renal artery aneurysm, 106–108
- Renal cystic lesions
- Bosniak classification, 48, 50
  - malignancy, 48, 50
  - nephrectomy specimen, 28–31
  - USS and CT findings, 53–55
- Renal infarction, 104–105
- Renal lymphoma, 70–72
- Renal-specific protocol and CT technique, 4–5
- Renal trauma, 28, 31
- Resistive index (RI), 7
- Retrocaval ureter, 102–103
- Retroperitoneum
- de-differentiated liposarcoma
    - anatomical structure, 149, 150
    - CT scan, 149, 150
    - resection recurrence rate, 149, 150
    - symptoms, 149, 150
  - fibrosis
    - aetiological factors, 172, 174
    - CT and nephrostogram image, 172, 174
    - idiopathic RPF, 172–174
    - RPF complication, 172–174
  - leiomyosarcoma
    - central low attenuation area
      - significance, 151, 152
    - CT image, 151, 152
    - gender predilection, 153, 154
    - hydronephrosis, 153, 154
    - inferior vena cava, 153, 154
    - metastasis frequency, 151, 152
    - venous leiomyosarcomata, 153, 154
  - liposarcoma
    - characteristics, 147, 148
    - de-differentiation risk, 147, 148
    - radiological appearance, 147, 148
    - treatment modality, 147, 148
  - lymphoma
    - chronic lymphocytic leukaemia, 167, 169
    - retroperitoneal adenopathy, 167, 169
  - metastatic lymphadenopathy
    - differential diagnosis, 170, 171
    - histology, 170, 171
  - metastatic non-seminomatous germ cell tumour
    - $\alpha$ -FP and  $\beta$ -HCG, 165, 166
    - lymph node mass, 165, 166

- NSGCT  
 axial CT, 163, 164  
 orchidectomy relapse rate, 163, 164  
 post-chemotherapy, 163, 164
- paraganglioma  
 benign pathology, 155, 156  
 malignancy, 155, 156  
 organs of Zuckerkandl, 155, 156  
 radionuclide tracer, 155, 156  
 sign and symptoms, 155, 156
- plexiform neurofibroma  
 malignant transformation risk, 159, 160  
 MPNST, 159, 160  
 NF-1, 159, 160
- PNET  
 coronal and axial CT, 157, 158  
 Ewing's sarcoma and Askin sarcoma, 157, 158  
 prognostic factor, 157, 158
- tuberculosis  
 CT-guided aspiration/biopsy, 161, 162  
 diagnosis, 161, 162  
 psoas abscess, 161, 162
- S**
- Sonography  
 anechoic dilated structure, 320, 331  
 detection, 302, 303  
 epididymis, 295, 297–300  
 homogenous reflectivity, 295–297  
 inguinal region, 302, 303  
 mediastinum testis, 295, 297  
 tunica albuginea, 295, 297
- Sperm granulomata, 336, 337
- Spinal cord injury  
 detrusor sphincter dyssynergia pattern, 403–404  
 filling phase, 401–402  
 fluoroscopic image, 401–402  
 neurogenic detrusor overactivity, 401–402
- Squamous cell cancer (SCC), 370–372
- Staghorn calculus, 82–83
- T**
- Testes and adnexae  
 acute focal testicular infarction, 312, 314  
 adenomatoid tumour, 304, 305  
 anatomy  
 fibrous septa, 295, 297  
 tunica albuginea, 295, 297  
 tunica vaginalis, 295, 297  
 bi-valved orchidectomy specimen, 298, 300  
 blood supply, 295, 297  
 CAH, 306–308  
 CT  
 inguinal cannal, 330, 331  
 nodal staging, 326, 328  
 cystic dysplasia, 309, 311  
 cystic transformation, 309, 310  
 Doppler image  
 hypervascular, 322, 324  
 intratesticular varicocele, 320, 331  
 mediastinum testis, 320, 331  
 preserved vascularity, 340, 341  
 vascularity, 332, 335  
 epididymal tail  
 abnormality, 304, 305  
 heterogeneous, 298–300  
 epididymitis, 298–300  
 extratesticular collection, 340, 341  
 fibrous tissue, 342, 343  
 florid recent infarction, 312–314  
 genitourinary tuberculosis, 342, 343  
 haematological malignancy, 322–324  
 H and E section, 302, 303  
 heterogeneous testicular, 326, 328  
 hydrocele  
 anechoic fluid collection, 315, 317  
 inflammatory debris, 315–317  
 treatment, 315, 317–319  
 tunica vaginal, 315–317  
 intratesticular varicocele, 320, 321  
 lack of vascularity, 298–300  
 marked hyperaemia, 298–300  
 mediastinum testis, 306, 307  
 MRI  
 importance, 302, 303  
 undescended testis, 302, 303  
 uses of, 302, 303  
 multiple punctate echogenic foci, 338, 339  
 multitude condition, 338, 339  
 NSGCT, 326, 328, 329  
 orchietomy specimen  
 abnormality, 309–311  
 bivalved, replacement, 326–328  
 epididymis, 312–314, 322–324  
 risk management, 342, 343  
 risk of condition, 342, 343  
 transverse section, 330, 332  
 white fibrous mass, 330, 332  
 poor vascularity, 326, 328  
 sonography  
 anechoic dilated structure, 320, 331  
 detection, 302, 303  
 epididymis, 295, 297–300  
 homogenous reflectivity, 295–297

- Testes and adnexae (*cont.*)
- inguinal region, 302, 303
  - mediastinum testis, 295, 297
  - tunica albuginea, 295, 297
  - spermatic cord, tumour, 330, 331
  - sperm granulomata, 336, 337
  - surgical exploration, 340, 341
  - symphysis, 322–324
  - testicular cancer, 302, 303
  - testicular lymphoma, 322–324
  - testicular microlithiasis, 338, 339
  - TGCT, 333, 335
  - tunica albuginea, 295–297, 340, 341, 344
  - UDT, 302, 303
  - ultrasound technique
    - undescended testis, 302, 303
    - upper pole, 336, 337, 340–343
  - Testicular appendages, 295, 297
  - Testicular germ cell tumor (TGCT), 333, 335
  - Testicular microlithiasis, 338, 339
  - Tissue–energy interaction, 1, 3
  - Transitional cell carcinoma (TCC), 53–55, 203–204, 214
  - Tuberculosis, 40–42
    - CT-guided aspiration/biopsy, 161, 162
    - diagnosis, 161, 162
    - psoas abscess, 161, 162
  - Tumour stage, 262–264
  - Tunica albuginea, 295–297, 340, 341, 344
  - Tunica vaginalis, 295, 297
  - TUR defect, 268, 269
- U**
- Ultrasound technique, 302, 303
  - Undescended right testis (UDT), 302, 303
  - Ureteric calculus, 77–79
  - Ureterocoele, 199–200
  - Urethral stricture, 425–426
  - Urinary tract infection (UTI), 225–226
  - Urodynamics
    - frequency and urgency
      - anticholinergic medications, 391–392
      - cough action, 385–386
      - detrusor overactivity, 393–396, 405–406
      - detrusor pressure, 405–406
      - filling phase, 385–386
      - nocturia and occasional urge incontinence, 387–388
      - obstruction, 395–396
      - overactive contraction, 395–396
      - rectal pressure, 417–418
      - T8 level spinal cord, 399–400
      - urge-related incontinence, 390–391
      - voiding phase, 393–394
      - voiding study, 385–386
    - incomplete emptying
      - detrusor pressure, 389–390
      - filling, 389–390
      - fluoroscopic image, 407–408
      - overactivity, 389–390
      - prostatic urethra trapping, 407–408
      - voiding phase, 407–408
    - incomplete voiding, 409–410
    - indwelling suprapubic catheter, 419–420
    - lower urinary tract symptoms, 413–414
      - detrusor overactivity, 421–422
      - flow rate, 415–416, 425–426
      - flow signal, 421–422
      - obstruction/hypocontractile detrusor, 427–428
      - overactive bowel contractions, 421–422
      - pinching, 427–428
      - spike artefact, 415–416
      - urethral stricture, 425–426
      - uroflow study, 423–424
    - obstructive voiding symptoms, 411–412
    - spinal cord injury, 396–397
      - detrusor sphincter dyssynergia pattern, 403–404
      - filling phase, 401–402
      - fluoroscopic image, 401–402
      - neurogenic detrusor overactivity, 401–402
- Urological disorder**
- computed tomography, 5
  - MRI, 8, 10
  - ultrasound, 6
- V**
- Von Hippel Lindau (VHL) disease, 45–47
- X**
- Xanthogranulomatous pyelonephritis (XGP), 33–37
- Z**
- Zoon's balanitis, 366, 368