

Index

A

- Accordion®, 41
- Age-related renal considerations, 180–181
- Amplatz serial dilators, 6
- Angled-tip angiographic catheter, 6
- Antiretropulsion devices, 41
- Avulsion. *See* Ureteral avulsion

B

- Ballistic lithotripsy
 - advantages and disadvantages, 33
 - electrokinetic energy, 34
 - pneumatic energy, 33–34
- Balloon dilator, 25
- Balloon retention catheters, 48–49
- Bleeding, URS
 - anticoagulation, 139
 - bladder irrigation, 138
 - fluoroscopic image, 139
 - and UPJ, 139
 - ureteral stent, 139
- BMI. *See* Body mass index (BMI)
- Body mass index (BMI)
 - categories, 66
 - dose-related effect, 69
 - stone location, 166
 - and stone size, 162
- “Bulls-eye” technique, 4, 8

C

- Calculi, staghorn, 52
- Caliceal diverticulum (CD)
 - CT reconstructions, 2
 - retrograde injection, 8
 - stones in, 7
- CD. *See* Caliceal diverticulum (CD)
- Clinical Research Office of the Endourologic Society (CROES), 63
- Coated stents
 - biofilm formation and bacterial adhesion, 132–133
 - drug eluting, 133
 - glycosaminoglycans, 133
 - heparin, 133
 - Triclosan, 133

Combilith® Walz, 34

Complications

PCNL

- aggregated stones, 67
- calix and pelvis, 62
- Clavien classification, 61–62
- complication grades, Clavien, 62
- and CROES, 63
- intraoperative(*see* Intraoperative complications, PCNL)
- Kumpe catheter, 75
- mid/lower pole site, 76
- multivariate analysis, 62
- and PCNLGS, 63
- postoperative(*see* Postoperative complications, PCNL)
- prevention, identification and treatment, 77–78
- renal pelvic stone and superior pole calyx, 75

URS

- intraoperative, 138–144
- postoperative complications, 144–147

Computed tomography (CT)

- CT-guided access, 6
- noncontrast, 7
- retrorenal colon, demonstration, 2

Contemporary access sheaths, 22

Contemporary ureterosopes, 22

Coupling, 155–156

CROES. *See* Clinical Research Office of the Endourologic Society (CROES)

CT. *See* Computed tomography (CT)

CyberWand®, 37

D

DECT. *See* Dual energy CT (DECT)

Delayed complications

- hemorrhage, 74
- infundibular stenosis, 74–75
- nephrocutaneous fistula, 74
- perinephric hematoma, urinoma and abscess, 75
- stricture, 74

Designs, stent

- dual durometer, 128–129
- loop tail, 128
- tail, 128

- Diabetes mellitus, 180
- Diathermy coagulation, 56
- Digital rigid nephroscope, 40
- Dilation and tract placement, 6–7
- Direct ureteroscopic vision, 21, 25
- Drainage tubes, types, 48–49
- Dual energy CT (DECT)
- surgical management, 161
 - uric acid and non-uric acid, 161
- E**
- ECIRS. *See* Endoscopic Combined IntraRenal Surgery (ECIRS)
- Electrohydraulic lithotripsy (EHL)
- combined probes, 37–39
 - generators, 153
 - portable devices, 39–40
- Electrokinetic lithotripters, 34
- Electromagnetic lithotripsy (EML) generators, 153–154
- Endo-irrigation systems, 99
- Endopyelotomy tubes, 48–49
- Endoscopic Combined IntraRenal Surgery (ECIRS)
- complication rate, 18
 - double J stent, 18
 - GMSV position, 14
 - retrograde access (*see* Retrograde ureteroscopy)
 - supine technique (*see* Supine technique)
- Endoscopic PCNL
- access sheath, deployment, 22
 - access tract, creation, 25–26
 - advantages, 26
 - balloon dilator, 25
 - challenges, 27–28
 - direct endoscopic visualization, 21
 - flexible cystoscope, 23
 - instrument setup guide, 24
 - large renal stones, 21
 - needle puncture site, 25
 - positioning, 22–23
 - potential advantages, 26
 - renal access, 24–25
 - renal collection, method, 26–27
 - retrograde access, 21–22
 - stone retrieval, 27
 - tract dilation, 25
 - ureteral access, 23–24
 - well-organized back table, 24
- ESWL. *See* Extracorporeal shock wave lithotripsy (ESWL)
- Extracorporeal shock wave lithotripsy (ESWL)
- acoustic pulse, 151, 152
 - cavitation, 152–153
 - coupling, 155, 156
 - description, 151
 - dynamic squeezing, 153
 - focal zone, 154–155
 - generators (*see* Generators)
 - imaging, 157
 - monitoring, 155–156
 - quasi-static squeezing, 152
 - rate, 156–157
 - spallation, 152
 - tear/shear forces, 152
 - urinary stones, 151
- Extrarenal damage
- cardiac and vascular injury, 184–185
 - fetal and reproductive organ injury, 185
 - pain, 183
 - pulmonary injury, 185
 - UTI and sepsis, 183–184
 - visceral organ injury, 184
- F**
- False passage, URS, 140
- Fibrin glue, 54–55
- Flexible cystoscope, 23
- Fluoroscopy, 4–5, 25
- Focal zone, ESWL, 154–155
- Fragmentation
- bubble-free ultrasound gel, 155
 - shock waves, 152
 - stone, 152
- Fragment clearance, SWL
- mechanical maneuvers, 168
 - pharmacologic measures, 168–170
- Fragment side effects
- Brushite stones, 183
 - multiple treatments, 182
 - steinstrasse, 181–182
 - stone recurrence, 183
 - ureteral stricture disease, 182
- 18 Fr Councill catheter, 8–9
- Functional renal injury
- categorization, 178
 - cavitation forces, 178
 - ¹³¹I-Hippuran, 179
 - parenchymal changes, 178
 - renal biopsies, 178
 - renal scarring, 179
 - SWL therapy, 179
- G**
- Galdakao-modified supine Valdivia (GMSV) position, 14–15
- Gelatin matrix, 55
- Generators
- EHL, 153
 - EML, 153–154
 - PZL, 154
- GMSV position. *See* Supine technique
- Guidewire placement, 83–84
- H**
- Hematoma/hemorrhage
- acute renal damage, 177

- bleeding diathesis, 178
 - CT and MRI, 177
 - description, 177
 - perirenal hematoma, 177, 178
 - subcapsular hematoma, 177
 - Hemostatic agents
 - FDA definition, 54
 - fibrin glue, 54–55
 - gelatin matrix, 55
 - Holmium laser
 - advantages and disadvantages, 33
 - fiber, 35
 - PNL, key technique, 36
 - Horseshoe kidney, percutaneous stone removal
 - risk, 41
 - SWL and, 43
 - technique, 31–32
 - ureteroscopy, 43
 - Hounsfield units (HU), 161
 - HU. *See* Hounsfield units (HU)
 - Hydrothorax
 - non-staghorn calculi, 68
 - procedure-related complications, 63
 - Hypertension, 179–180
- I**
- Infection. *See* Sepsis
 - Informed consent
 - laparoscopic ureterolithotomy, 96
 - observation/medical expulsive therapy, 95
 - PCNL, 95–96
 - SWL, 95
 - ureteral stent, 95
 - URS, 95
 - Infundibulo-pelvic angle (IPA)
 - IVP and, 116
 - lower pole calices, 116
 - Injury
 - antegrade nephrostogram, 143
 - mucosal (*see* Mucosal Injury)
 - thermal, 140
 - Instrumentation. *See also* Techniques
 - horse-shoe kidney PNL, 43
 - list, 42
 - stone fragment retrieval, 40
 - Internal drainage, types
 - antegrade stent placement, 53
 - tethers, 53
 - Intracorporeal lithotripsy, 31, 32
 - Intracorporeal lithotripters, 97
 - Intraoperative complications, PCNL
 - air embolism, 73
 - bowel injury, 71
 - collecting system and fluid extravasation, 72
 - duodenal injury, 71
 - hemorrhage, 70
 - hypothermia, 72
 - lung and pleura injury, 70–71
 - nerve injuries, 72–73
 - spleen, liver and gall bladder injury, 71–72
 - ureteral avulsion, 73
 - Intraoperative retrograde ureteroscopy, 17–18
 - Intraoperative risk factors, PCNL
 - access, 68–69
 - dilation, 69
 - patient positioning, 68
 - radiation exposure, 69
 - tubeless PCNL, 69–70
 - IPA. *See* Infundibulo-pelvic angle (IPA)
- K**
- Kidney. *See* Horseshoe kidney, percutaneous stone removal
- L**
- LASER. *See* Light amplification by stimulated emission of radiation (LASER)
 - Laser lithotripsy, 35–36
 - Light amplification by stimulated emission of radiation (LASER), 35
 - Lithobreaker®, 40
 - LithoClast Ultra®
 - in vitro* studies, 39
 - limitations, 38
 - pneumatic probe, 38
 - Lithopump®, 38–39
 - Lithotripsy
 - ballistic (*see* Ballistic lithotripsy)
 - EHL (*see* Electrohydraulic lithotripsy (EHL))
 - holmium laser, 120
 - Ho:YAG, 115
 - intracorporeal, 31, 32
 - laser, 35–36
 - SWL (*see* Shock wave lithotripsy (SWL))
 - tubeless procedure, 49
 - ultrasonic, 32–33
 - Lithovac device, 33
 - Lower pole. *See also* Ureteroscopy (URS)
 - active deflection, 116
 - description, 116
 - flexible ureteroscopes, 116
 - IPA and IVP, 116
 - and SFR, 116
 - stone displacement (*see* Stone distraction)
 - SWL and PCNL, 116
- M**
- Management
 - airway, 66
 - conservative management, 71
 - peritoneal signs/sepsis, 71
 - Medical expulsive therapy
 - analgesia, 92
 - patient decision-making, 92
 - patient factors, 92
 - surgical intervention, 92, 93
 - Mucosal injury, 139–140

N

NCCT. *See* Non-contrast computed tomography (NCCT)

Nephrolithotomy, 2

Nephrostomy

- drainage, 49
- tube, size, 47–48

Nomogram

- logistic regression analysis, 161–162
- stone attenuation, 163
- stone length, location and number, 159

Non-contrast computed tomography (NCCT), 31

O

Operative management

- ancillary instrumentation, 96
- balloon dilators, 97
- C-arm monitors, 99
- catheters, 97
- endo-irrigation systems, 99
- flexible ureteroscopes, 98
- guidewires, 96
- intracorporeal lithotripters, 97
- orifice cannulation, 99
- placement, safety wire, 100–101
- preoperative antibiotics, 99
- semi-rigid ureteroscopes, 98
- stone fragmentation/extraction, 102
- stone retrieval devices, 97–98
- ureteral access sheaths, 97
- ureteral occluding devices, 97
- ureteroscope insertion, 101
- URS algorithm, ureteral stone, 99, 100

Organ injury. *See* Extrarenal damage

P

Patient positioning, 3–4

Patient selection, SWL

- and BMI, 162
- and DECT, 161
- fragility, 160–161
- literature review, predictors, 162, 163
- lower pole stone study, 160
- meta-analysis, 159–160
- middle and distal ureter, 160
- nomogram, 161–162
- PCNL and RCTS, 160
- renal and ureteral calculi, 161
- ROC curves, 162
- and SSD, 161

PCNL. *See* Percutaneous nephrolithotomy (PCNL)

PCNL Global Study (PCNLGS), 63–64

Perc N-Circler, 41

Percutaneous

- access, 31
- pyelolithotomy technique, 13
- retrograde approach, 13–14
- stone removal, 13
- tract dilation, 17

Percutaneous nephrolithotomy (PCNL)

- anatomical considerations, 2–3
- angled-tip angiographic catheter, 6
- antegrade percutaneous semirigid URS, 94
- antiretropulsion devices, 41–42
- ballistic lithotripsy (*see* Ballistic lithotripsy)
- C-arm rotation, 4–5
- CD stones, 9
- classification, complications, 61–63
- contemporary indications, 2
- CT guided access, 6
- description, 61
- digital rigid nephroscope, 40
- dilation and tract placement, 6–7
- and EHL (*see* Electrohydraulic lithotripsy (EHL))
- endoscopic (*see* Endoscopic PCNL)
- equipment, 4
- fascial incising needle, 7
- fluoroscopic access techniques, 3–4
- fluoroscopy, 4–5
- hydronephrosis, 93
- instrument list, 4
- intraoperative complications, 70–73
- intraoperative risk factors, 68–70
- laboratory investigations, 2
- and laparoscopic ureterolithotomy, 94
- laser lithotripsy, 35–36
- lung injury, 95–96
- management of renal calculi, 113
- modifications, 21–22
- morbidity, 61
- obese/morbidly obese patient, 114
- patient positioning, 3–4
- pre-operative evaluation and risk factors, 63–68
- pre-operative planning, indications, 2
- prevention, identification, and treatment, 77–78
- renal collecting system, 21, 26
- retrorenal colon. CT demonstration, 2
- rigid nephroscopy, 7
- safety principles, 10
- second needle puncture and dilated tracts, 27
- stone-free rates, 160
- stone retrieval devices, 40–41
- techniques (*see* Techniques)
- treatment modalities, 160
- ultrasonic lithotripsy, 32–33
- ultrasound guided puncture, 5–6
- ureteral stones, 61
- URS vs., 115

Percutaneous stone removal. *See* Percutaneous nephrolithotomy (PCNL)

Perforation. *See* Ureteral perforation

Photo acoustic effect, 35

Piezoelectric (PZL) generators, 154

Placement, stents

- description, 130
- distal tip to deploy, 131
- fluoroscopic guidance, 130
- guidewire, kidney, 131
- proximal curl, deploy, 131

- pubic symphysis, 131
 - right ureteral calculus, 130
 - Pneumatic energy, 33–34
 - Pneumatic lithotripters, 33–34
 - Polyurethane
 - description, 126
 - distal curl, 130
 - radial coil reinforcement, 126
 - Positioning of patient
 - comfortable, 14
 - operating room organization, 14–16
 - Postoperative complications, PCNL
 - death, 74
 - fever and sepsis, 73–74
 - Postoperative renal drainage, 13
 - Post PCNL drainage
 - diathermy coagulation, 56
 - drainage tubes, types, 48–49
 - hemostatic agents (*see* Hemostatic agents)
 - inclusion criteria, 56
 - internal drainage, types, 53
 - nephrosotomy drainage, evolution, 49
 - nephrostomy tube, size, 47–48
 - tubeless modifications (*see* Tubeless modifications)
 - tubeless PCNL, 49, 52–53
 - Preoperative measures, SWL
 - antibiotics, 164
 - antioxidants and calcium channel blockers, 164–165
 - Pre-operative risk factors, PCNL
 - anti-coagulation medications, 64
 - and BMI, 64
 - coagulopathy, 63–64
 - negative urine culture, 64
 - obesity, 65–66
 - operative volume and surgeon experience, 64–65
 - patient age and co-morbidity, 65
 - radiographic investigation, 64
 - renal anomalies, 66–67
 - renal surgery, 67
 - stone burden, 67–68
 - Prone percutaneous access. *See* Percutaneous nephrolithotomy (PCNL)
 - Prone position, patient
 - drapes, 23
 - endoscopic access, 23
 - PCNL, 22–23
 - Proximal ureteric stones, 31–32
 - PZL generators. *See* Piezoelectric (PZL) generators
- R**
- Randomized clinical trials (RCTs)
 - post-SWL UTIs, 164
 - renal calculi, 160
 - shock wave rate, 167
 - RCTs. *See* Randomized clinical trials (RCTs)
- Renal calculi
 - indications, ureteroscopic management, 113, 114
 - lower pole (*see* Lower pole)
 - SFRs, 122
 - SWL and PCNL, 113, 114
 - SWL vs. URS, 113, 114
 - URS, 114, 115
 - Retrograde ureteroscopy
 - evaluation, preliminary, 16–17
 - intraoperative, 17–18
 - Reverse lithotomy position, 13
 - Rigid nephroscopy, 7
 - Risk factors, SWL
 - age-related, 80–181
 - diabetes mellitus, 180
 - functional renal injury, 178–179
 - hematoma/hemorrhage, 177–178
 - hypertension, 179–180
 - Roth-Net retrieval device®, 41
- S**
- Sepsis
 - bacteremia and, 145
 - cephalosporins, 145
 - nitrite negative urine, 145
 - preoperative antibiotics, 144
 - stent/percutaneous nephrostomy, 145
 - urea-splitting bacteria, 145
 - urine culture, 144–145
 - SFR. *See* Stone-free rates (SFR)
 - Shock wave delivery
 - pattern, administration, 166–167
 - rate, 167
 - Shock wave lithotripsy (SWL)
 - CD stones, 7
 - contemporary surgical techniques, 116
 - extrarenal damage, 183–185
 - fragment clearance (*see* Fragment clearance, SWL)
 - fragment side effects, 181–183
 - informed consent, 95
 - locations, renal calculi, 113
 - patient selection (*see* Patient selection, SWL)
 - preoperative measures, 164–165
 - protective pretreatment/treatment protocols, 186–187
 - proximal ureteral stones, 93
 - residual fragments, 170
 - risk factors (*see* Risk factors, SWL)
 - stone type and anatomic characteristics, 113
 - treatment, 9, 165–167
 - upper urinary tract stones, 159
 - and URS, 93, 159
 - vs. URS management, 114, 115
 - URS vs. PCNL, 115
 - Skin-to-stone distance (SSD)
 - description, 161
 - logistic regression, 161–162
 - mean stone density, 162
 - SSD. *See* Skin-to-stone distance (SSD)

- Steinstrasse
 asymptomatic patients, 182
 description, 181
 left proximal and mid-ureter, 181
 stone size, 181–182
 ureteral perforation, 182
 ureteral stent, 182
- Stent
 biodegradable, 133–134
 designs (*see* Designs, stent)
 discomfort, 132
 double J-type stents, 125
 drug elution, 133
 endopylotomy, 128
 evolution, 125
 glycosaminoglycans, 133
 heparin-and Triclosan-coated, 133
 injuries, 129–130
 lithotripsy/stone extraction, 125
 medical treatment, discomfort, 132
 placement (*see* Placement, stents)
 polymers, 126
 reinforced/metal, 126–127
 renal colic, 125
 retrograde injection, 127–128
 tether, 130
 variable length types, 128
- Stent morbidity
 designs (*see* Designs, stent)
 phenazopyridine, 132
- Sterile draping, 16
- Stone Breaker®, 39–40
- Stone distraction
 basket extraction, 121
 fluoroscopic and intraoperative images, 119, 120
 upper pole calyx, 119–120
- Stone-free rates (SFR)
 CD treatment, 9
 gravitation effect, 116
 laparoscopic techniques, 9
 lower pole and renal calculi, 116, 122
 morbidity, 116
 and PCNL, 1, 121
- Stone retrieval devices, 40–41
- Stricture. *See* Ureteral stricture
- Submucosal/lost stone
 “silent stricture”, 143
 stone granuloma/ureteral stenosis, 42
 ureter/retroperitoneal urinoma, 142
- Supine technique
 endovision-assisted renal puncture, 17
 GMSV position, 14–15
 intraoperative retrograde ureteroscopy, 17–18
 “kebab” patient, preparation, 17
 management advantages, 14
 patient positioning and organization, 14–16
 percutaneous and retrograde accesses, 16
 postoperative renal drainage, 13
 retrograde ureteroscopy evaluation, 16–17
 rigid nephroscope, movement, 17
 sterile draping, 16
 urological advantages, 14
 Swiss LithoClasT®, 33–34
 SWL. *See* Shock wave lithotripsy (SWL)
- T**
- TDPN. *See* Transdiverticular puncture and neoinfundibulotomy (TDPN)
- Techniques
 ballistic lithotripters, 33–34
 digital rigid nephroscope, 40
 electrohydraulic lithotripsy, 36–37
 laser lithotripsy, 35–36
 ultrasonic lithotripsy, 32–33
- Technology failures, URS, 144
- Thermal injuries, 140
- Transdiverticular puncture and neoinfundibulotomy (TDPN), 9–10
- Treatment, SWL
 anesthesia, 165–166
 coupling, 166
 positioning, patient, 165
 shock wave delivery (*see* Shock wave delivery)
- Triangulation techniques, 4, 9
- Tubeless modifications
 in children, 52–53
 complex PCNL, 51–52
 conservative modification, 51
 meta-analysis, 50–51
 nonrandomized trials, 51
 pain scale results, 49
 randomized controlled trials, 49–50
 visual analog scale pain score, 50
- Tubeless PCNL
 bilateral, 52
 obese patients, 52
 patient, supine position, 52
 totally, 53
- U**
- Ultrasonic lithotripsy, 32–33
- Ultrasonic lithotripter, 31
- Ultrasound (US)
 guided puncture, 5–6
 lithotripsy (*see* Ultrasonic lithotripsy)
- UPJ. *See* Uretero-pelvic junction (UPJ)
- Ureteral access sheath
 clinical utility, 87, 88
 description, 86–87
 diameter, 87
 disposable devices, 86
 kink-resistant sheath, 86
 PTFE, 84
 technique, 87
- Ureteral avulsion
 antegrade nephrostogram, 143
 auto-transplantation, 144
 basket extraction, 143

- description, 143
 - endourologic basket, 143
 - stricture formation, 144
 - transureteroureterostomy, 144
 - Ureteral balloon, 85
 - Ureteral catheter displacement, 4
 - Ureteral perforation
 - description, 140
 - hydrophilic wire, 141
 - laser fibers, 141
 - percutaneous nephrostomy tube, 142
 - rigid/flexible URS, 141
 - treatment, 142
 - ureteroscopic laser lithotripsy, 141
 - ureteroscopic treatment, 141
 - Ureteral stent. *See* Stent
 - Ureteral stricture
 - asymptomatic patients, 146
 - avulsion, 146
 - description, 145
 - endopyleotomy, 146
 - length and location, 146
 - retrograde pyelogram, 146
 - retrograde ureterogram and antegrade nephrostogram, 146, 147
 - Uretero-pelvic junction (UPJ), 3
 - Ureteroscopic access
 - balloon dilation, 85–86
 - guidewire placement, 83–84
 - placement, 84–85
 - reporting investigations, 87, 88
 - ureteral access sheath, 86–87
 - “wireless” approaches, 86
 - Ureteroscopy (URS)
 - access tract, creation, 25–26
 - basket extraction, stone, 121
 - calices, 27
 - classification, 138
 - current flexible ureteroscopes, 117, 118
 - description, 137
 - endoscopic PCNL, 27–28
 - flexible vs. rigid, 121
 - instrument list, 117
 - intraoperative complications, 138–144
 - with obesity, 114–115
 - patient preparation, 117, 119
 - pediatric cystoscope, 83
 - postoperative complications, 144–147
 - renal calculi (*see* Renal calculi)
 - renal pelvis, 86
 - repair costs, 117
 - retrograde, 17–18
 - retrograde placement, 87
 - safety techniques, 137–138
 - stone composition and, 115–116
 - stone type and anatomic characteristics, 113
 - technique and perioperative setup, 119
 - ureteral stones
 - edema and polypoid reaction, 107
 - fluoroscopy, safety wire, 105, 106
 - hematuria, 103
 - infectious complications, 102
 - informed consent, 95–96
 - left ureteral stone, 104, 105
 - management pathway, urolithiasis, 90, 91
 - meta-analysis after ureteroscopy, 103
 - nuclear diuretic renal scan, 107
 - operative management, 96–102
 - postoperative imaging, 104
 - postoperative management, 103–104
 - preoperative counseling, 91–94
 - retained stent, 103
 - retrograde pyelogram, 105, 106
 - stone extraction, basket, 107
 - stone fragmentation, holmium laser, 106
 - SWL therapy, 104–105
 - ureteral injury, 102
 - ureteral stricture, 102
 - visualization, reactive polyps
 - and impaction, 106
 - well-organized back table, 24
 - Urinary calculi, 37
 - Urinary tract infection (UTI)
 - and sepsis, 183–184
 - SWL risk factors, 177
 - Urolithiasis
 - coagulopathy, 178
 - complex, 13, 14
 - conservative management, 179
 - ureteroscopy, 18
 - URS. *See* Ureteroscopy (URS)
 - UTI. *See* Urinary tract infection (UTI)
- V**
- Vesicoureteral reflux (VUR)
 - cystourethrogram, 145
 - oral pain medication, 145
 - ureteral orifice, 145
 - VUR. *See* Vesicoureteral reflux (VUR)
- W**
- Wireless approaches, 86, 119