

INDEX

- Accipiter gentilis* (goshawk), 153
 A. nisus (sparrowhawk), 156
Actitis macularia (spotted sandpiper), 19
Adaptation, 168
Adult predation hypothesis, 233–234
 and hatching, asynchronous, 233–234
 and survival rates of females, 234
Aethia pusilla (auklet), 163
Agelaius phoeniceus (red-winged blackbird), 5, 6, 11, 12, 16, 18, 19, 26, 30–32, 119
Aggression, 18–26
 and care, parental, 18–26
 investment
 posthatching, 22–23
 prehatching, 22
 male-male, 18–19, 22–23
 and testosterone, 18–26
 implant, experimental, 19–22
Alarm signal hypothesis, 159–160
 flash pattern, 160
Anabolism, induced, 90, 96–109
 index of condition, nutritional, 96–109
Anas platyrhynchos (mallard), 9
 A. sp. (duck), 155, 159
Anchovy, 150
Anthus pratensis (meadow pipit), 117, 118
Antihabituation hypothesis of song
 birds, 130
Aphelocoma c. coerulescens (scrub jay), 100–101
Aposomatism hypothesis, 151–152
Aptenodytes patagonicus (penguin), 162
Arenaria interpres (turnstone), 162
Ariana arbustorum (snail), 230
 and cannibalism, 230
Asio flammeus (short-eared owl), 227
Asynchrony of hatching, 191–270
 correlates, 210–211
 evolution hypotheses, 211–214
 paradox of, 192–194
 quantification, 203
Aves, phylogeny, 205–210
 orders, twenty-three, listed, 205
Beau geste hypothesis of bird song, 130
Birds
 breeding
 density and testosterone level, 17
 and endocrinology, 3
 opportunity, limited, hypothesis, 219
 success, 23–26
 mate attraction, 23–26
 condition, nutritional, 90
 anabolism, induced, 90
 body mass, 90
 fat supply, 90
 eyes
 clarity, visual, 150
 sighting lines, 148
 stripes, 148
 polygyny, 1–41
 song, 117–120
 adaptation, perceptual, 131
 signature adaptation, 131
 survey during migration, 43–48
 and testosterone, 1–41

- Birds (*cont.*)
 voice discrimination, 115–139
 wings, flash color, 150
- Blackbird, yellow-headed, 5, 6, 11–16,
 19–25
- Branta canadensis* (Canada goose), 92
- Breeding Bird Survey in North America,
 43, 66–73
 in Michigan, 73
 in Ontario, 70, 72
- Brood
 parasitism, 9–11
 hypothesis, 220–221
 reduction
 hypothesis, 193, 221–223
 strategy, 202
- Bubulcus ibis* (cattle egret), 229
- Buteo albonotatus* (zone-tailed hawk),
 154
B. jamaicensis (red-tailed hawk), 92
- Caching, 107–109
- Callipepla californica* (California quail),
 15
- Camouflage coloration, *see* Crypticity
- Cannibalism by nestlings, 230
- Care, parental, 18–26
 and male-male competition, 22
 posthatching, 22–23
 prehatching, 22
- Cathartes aura* (turkey vulture), 154
- Centrocercus urophasianus* (sage grouse),
 29
- Ceryle rudis* (pied kingfisher), 220
- Challenge hypothesis, 3–4
 and testosterone secretion, control of,
 3–4
- Chen caerulescens* (snow goose), 147
- Christmas bird counts, 44, 54
- Ciconia maguari* (Maguari stork), 251
- Cohesion of flock hypothesis, 160
- Colaptes auratus* (northern flicker), 159
- Colinus virginianus* (bobwhite), 202
- Coloration of birds, 141–190
 direction, future, for the study of, 169–
 178
 evolution, 141–190
 hypotheses
 functional, twenty-eight, 143–169
 nonfunctional, two, 144–145
 polymorphism, 154
- Columba leucocephala* (white-crowned
 pigeon), 236
C. livia (rock dove), 153
- Conflict, sexual, hypothesis, 231–232
 and hatching, synchronous, 231–232
- Correlations, genetic, hypothesis, 168–
 169
 and pleiotropy (linkage), 168
- Corvus frugileus* (rook), 226
- Coturnix coturnix* (Japanese quail), 26–
 29, 162
- Crypticity hypothesis, 151
 and birds, 151
 and insects, 151
- Cyanocitta stelleri* (jay), 159
- Delichon urbica* (house martin), 228
- Dendrocopos major* (great-spotted wood-
 pecker), 156
- Dendroica petechia* (yellow warbler),
 119, 215
- Dicrurus* sp. (drongo), 152
- Diomedea exulans* (wandering albatross),
 93
- Diversity, dietary, hypothesis, 227
- Dolphin, 150
- Dominance, social, 104–107
- Drift, genetic, hypothesis, 169
- Efficiency, aerodynamic, 145–146
 tail, functions of, 145
 wing shape, 145
- Effort, reproductive, 99–100
- Egg
 development, start of, 198
 zero, physiological, is 27–28 C., 198
 laying, energy constraint hypothesis,
 214
 protection hypothesis, 218–219
 size, 202
 temperature, 198
 viability hypothesis, 216–218
- Emberiza citrinella*, 118
- Empidonax minimus* (least flycatcher),
 233
E. trecillii (willow flycatcher), 148
- Energy constraint hypothesis of egg-lay-
 ing, 214
- Enhancement of displays hypothesis, 159
 plumage, distinctive, 159
- Eoaves, 204

- EPOQ program in Quebec, 54
Erithacus rubecula (European robin), 119, 121
Erythropygia zambesians (red-backed scrub robin), 150
Eudyptes chrysocome (penguin), 162
E. sp. (penguin), 201
 brood reduction strategy, 202
Euplectes progne (long-tailed widow-bird), 146
E. sp. (widow bird), 160
 Extra-pair copulation, 2
- Fagus americana* (beech), 106
Falco columbarius (merlin), 152, 155
F. peregrinus (peregrine), 109, 156
 in Wisconsin, 65
F. sparverius (kestrel), 109, 214
F. sp. (hawk), 47, 49, 54, 64
 Fault bar of feather, 92
 Feather
 fault bar, 92
 growth bar, daily, 89
 growth
 and temperature, ambient, 95
 and wind chill, 95
 and keratin, 91
 and melanin, 146
 pulled and growth of replacement, 90
 see Ptilochronology
 Feeding solicitation hypothesis, 158–159
Felis catus (cat), 152
Ficedula hypoleuca (pied flycatcher), 6, 7, 17, 19, 20, 119, 156, 214
 Fledging patterns and onset of incubation, 196
 Flight, nocturnal, 45, 47
 Floater and territory, 13–14
 Flock attraction hypothesis, 160
 Follicle history, 95–96
 Foraging, facilitation of, 148–151
 clarity, visual, 150
 flare reduction, 148
 flushing prey, 150–151
 sighting lines, 148–150
Fringilla coelebs (chaffinch), 118, 163
Fulica americana (coot), 167
- Gallinula chloropus (moorhen), 211
Geothlypis trichas, 118
 Glare reduction, 148
- Gloger's rule, 147
 Great tit, 124, 125
 Growth bar of feather
 causes, 91–92
 and condition, nutritional, see Ptilochronology
 definition, 90
 and keratin, 91
 one is equivalent to 24 hours of growth of feather, 93
 ubiquity, 92–93
 width, 93–96
- Habitat selection, 101–104
 Hatching, asynchronous, 191–270
 evolution hypotheses, 211–235
 tabulated, 213
 hypothesis, see separate hypothesis
 incubation, onset of, 191–270
 model, 239–250
 simple, 241
 stochastic, 244
 patterns in birds, phylogenetic, 203–210
 spread, 200–201
 tests, thirty-nine, experimental, 235–250
 designs, 235
 problems, 238–239
 results, 235–238
- Hawk Migration Association of North America, 54
Hirundo pyrrhonota (cliff swallow), 131, 162
H. rustica (barn swallow), 131, 146
 Homobatrachotoxin, 152
 Honest advertisement hypothesis of bird song, 130
 and male quality, 130
 Hormone hypothesis of hatching asynchrony, 215
 and onset of incubation, 197
 Hurry-up hypothesis of hatching, asynchronous, 228
Hylocichla fuscescens, 118
 Hypotheses for the evolution of coloration, thirty-nine
 facilitation of foraging, 148–151
 listed, 144–145
 nonadaptive, 167–169
 physiological, 145–148

- Hypotheses for the evolution of coloration, thirty-nine (*cont.*)
 signaling
 interspecific, 151–158
 intraspecific, 158–167
- Hypotheses, thirty, testing of, 169–173
 evaluations, 170, 173
 importance, relative 173
 predictions for the evolution of birds for coloration, 171–174
 ornamentation, 171–174
- Icterus virens*, 119
I. sp. (oriole), 157
- Identity badges, arbitrary, 162
- Image avoidance hypothesis, 154
 and color polymorphism, 154
- Incubation by birds
 and control of hatching, 192
 and fledging patterns, 196
- Indexing signal hypothesis, 158
- Insensitivity, social, hypothesis, 28
- Insurance hypothesis, 224–225
 and clutch size in birds, 224–225
 and eggs, last laid, 225
- Interaction, social, 26–28
 and testosterone level, 26–28
- Investment, parental, *see* Effort, reproductive
- Jaeger, 153
- Junco hyemalis* (dark-eyed junco), 29, 201
- Jynx sp.* (wryneck), 152
- Kin recognition hypothesis, 162
- Labidomera clivicollis* (beetle), 230
 and cannibalism, 230
- Lagopus lagopus lagopus* (willow ptarmigan), 28
L. lagopus scotius (red grouse), 16, 17
- Landbird
 population count in migration, 43–88
 species, seventy-six, listed, 75–76
- Lanius ludovicianus* (loggerhead shrike), 97, 102–103, 110
- Larder hypothesis, 230
 and hatching, asynchronous, 230
- Larus argentatus* (herring gull), 158, 218
L. atricilla (laughing gull), 228
L. delawarensis (gull), 162
- Larus argentatus* (herring gull) (*cont.*)
L. glaucescens (black-headed gull), 151
- Lizard, 29
- Long Point Bird Observatory, Ontario, 54, 57
- Male
 aggression, 18–26
 attraction, 23–26
 and breeding success, 23–26
 competition, 22–23
 migratory, 11–13
 nonmigratory, 11–13
- Mallard and testosterone level, 9
- Mate attraction, 23–26
 and breeding success, 23–26
- Mating systems, 1
- Melaenornis sp.* (black flycatcher), 152
- Melanerpes formicivorus* (acorn woodpecker), 220
- Melanin, 168
 and ultraviolet light, 147
- Melospiza melodia* (song sparrow), 5, 6, 17, 19, 26, 27, 29, 119, 122, 123
M. georgiana, 119
- Merops viridis* (nestling bee eater), 229
- Mettnau-Reit-Illnitz program of banding birds, 52
- Micrathene whitneyi* (elf owl), 219–220
- Migration of landbirds, 43–88
 count, 44–88
 by banding, 51–52
 choice of site, 49–50
 covariance analysis, 58–64
 daily, 47, 55–56
 data analysis, 56–63
 data collection, 48–56
 definition, 44–48
 evaluation for population monitoring, 64–74
 future in North America, 78–81
 index, annual, 56–61
 formula, 58
 nocturnal, 45, 47
 operation Baltic, 54
 population size, 64–65
 pros and cons, 74–78
 by radar, 48
 by Toronto Ornithological Club, 53
 by trapping, 50–52
 trends, 61–74

- Migration of landbirds (*cont.*)
 count (*cont.*)
 what is it? 44–48
- Mimicry
 aggressive, 154
 deceptive, 122
 female, 167
 hypothesis, 152–153
 social, 153
 types of, 152
- Mimus polyglottus* (northern mockingbird), 150
- Molothrus ater* (brown-headed cowbird), 7, 9, 27, 29
- Monogamy-polygamy boundary, 1–2
- Motacilla alba* (meadow pipit), 156
- Nectarinia johnstoni* (scarlet-tufted malachite sunbird), 146
- Neoaves, 204–205
- Neocossyphus* sp. (ant thrush), 152
- Neotoma floridana* (wood rat), 226
- Neotropical Migrant Bird Conservation Program (Partners in Flight), 44, 78–81
- Nest
 attentiveness, 198–199
 failure hypothesis, 232–233
 and hatching, asynchronous, 232–233
 and nest failure ratio, 233
- Nesting cycle
 timing of periods, 195–196
 factors, 231–235
- North American Migration Count, 54
- Oceanodroma leucorhoa* (storm petrel), 92
- Onset of incubation hypothesis, 191–270
 bases, physiological, 197
 constraint
 adaptive, 215–231
 nonadaptive, 214–215
 factors, external, 197–199
 and hormones, 197
 patterns, 199–203
- Operation Baltic, 54
- Oriolus* sp. (New Guinean orioles), 152
- Ornamentation of birds, 141–190
 evolution of, 141–190
- Ottenby Bird Observatory, 65
- Owner of territory, 13–14
- Parabuteo unicinctus* (hawk), 226
- Paradox of hatching asynchrony, 192–193
- Partners in Flight, *see* Neotropical Migrant Bird Conservation Program
- Parus atricapillus* 118
P. bicolor (tufted titmouse), 93, 94, 106, 119, 123
P. carolinensis (Carolina chickadee), 93, 94, 106, 110
P. major (great tit), 118, 124, 163
P. montanus (willow tit), 90, 104, 105, 111
P. palustris (march tit), 2, 4
- Passer domesticus* (house sparrow), 19, 20, 95, 216
 asynchrony modeling, 245–250
- Passerina cyanea*, 118
- Pandion haliaetus* (osprey), 222
- Peak load reduction hypothesis, 227
 and hatching, asynchronous, 227
- Pelecanus erythrorhynchos* (pelican), 202
- Penguin, 150, 162, 201, 2029
- Phainopepla nitens* (phainopepla), 147
- Phasianus colchicus* (ring-necked pheasant), 5, 109
- Philemon* sp. (friarbird), 152
- Phylloscopus* sp. (warbler), 159
- Picoides borealis* (red-cockaded woodpecker), 103, 1–4
P. pubescens (downy woodpecker), 94, 103
- Pinus australis* (longleaf pine), 103
P. echinata (shortleaf pine), 103
P. taeda (loblolly pine), 103
- Pipilo erythrophthalmus* (rufous-sided towhee), 119, 121
- Plectrophenax nivalis* (snow bunting), 233
- Ploceus taeniopterus* (masked weaver), 220
- Plumage
 carotenoid, 168
 color patterns, examples of, 149
 distinctive, 159
 efficiency, aerodynamic, 145–146
 elaboration, *see* Ornamentation
 protection
 physical, 146
 from ultraviolet light, 147–148
 red, 169
 thermoregulation, 146–147
 white, 169

- Podiceps gallardoi* (hooded grebe), 224
Poephila bichenovi (finch), 161
P. guttata (finch), 161, 226, 236
Point Reyes Bird Observatory, 55
Poison dart frog, 152
Polygamy, 1–2
- monogamy boundary, 2
and testosterone, 1–41
types of, 4–11
 sequential, 6–8
 simultaneous, 5–6
Pooectes gramineus (vesper sparrow), 8
Population of landbirds
 count in migration, 43–88
Prophyrio porphyrio (swampphen), 158
Predator deflection hypothesis, 156–157
 injury-feigning, 156
 and lizard, 156
Prey, unprofitable, hypothesis, 155
Promiscuity, 1, 8, 9
Pterodroma arminjoniana (petrel), 153
P. neglecta (petrel), 153
Ptilochronology, 89–114
 definition, 89
Ptilonorhynchus violaceus (satin bowerbird), 9, 14, 18, 29, 175
 and testosterone level, 9, 14,–15
Pursuit deterrence hypothesis, 158
Pursuit invitation hypothesis, 157–158

Quiscalus major (boat-tailed grackle), 233

Radar count of landbirds, 48
Ramphocelus sp. (tanager), 148
Recognition, individual, hypothesis, 162
Regulus calendula, 65
 R. satrapa, 65
 R. sp. (warbler), 159
Reproduction, synchronous, 191
Resource-defence polygyny, 2
Resource tracking hypothesis, 221
 and hatching, asynchronous, 221
Rissa tridactyla (kittiwake), 222

Seizurus aurocapillus, 118
Selection, sexual, hypothesis, 163–167
 choice of male by female, 163–166
 intersexual, 163
 intrasexual, 163
 plumage manipulation, 165

Setophaga ruticilla (redstart), 119, 123
Sex ratio manipulation hypothesis, 225–226
 and starvation, 226
Siblicide, 230
 facultative, 222
Sibling rivalry hypothesis, 228–230
 and hatching, asynchronous, 228
Signal
 background light, 175
 design, tactical, 175
 exploitation, sensory, 175–177
 hypothesis, interspecific, 151–158
 apostomatism, 151–152
 crypticity, 151
 image avoidance, 154
 mimicry, 152–154
 predator deflection, 156–157
 prey, unprofitable, 155–156
 pursuit deterrence, 158
 pursuit invitation, 157–156
 startle/confusion effects, 157
 hypothesis, intraspecific, 158–167
 alarm signal, 159–160
 display enhancement, 159
 feeding solicitation, 158–159
 flock attraction, 160
 flock solicitation, 160
 indexing signals, 158
 mimicry, female, 167
 recognition
 by individual, 162
 by kin, 162
 by species, 162
 selection, sexual, 163–167
 status signal, 163
 network, neural, 177–178
 selection, 174–175
 system, sensory, 175–177
Sitta carolinensis (white-breasted nuthatch), 93, 94
 S. europaea (nuthatch), 65, 108
Skua, 153
Social insensitivity hypothesis, 28
Song of bird, 117–120
 composition, 120–122
 hypotheses, three
 antihabituation, 130
 beau geste, 130
 honest advertisement, 130

- Song of bird (*cont.*)
 neighbor-stranger discrimination, 117–120
 for recognition by individual, 117–120
 rendition, 122
 repertoire of thirty-one species, 118–122
 voice, individual, 123–125
- Songbird flights, 54
- Species recognition hypothesis, 160–162
 and coloration, 161
 and hybridization, 161
- Spheniscus* (penguin), 150–151
- Spizella arborea* (tree sparrow), 96
S. pusilla (field sparrow), 117, 118
- Startle/confusion effect hypothesis, 157
- Status signal hypothesis, 163
- Sterna hirundo* (tern), 218
- Stizorhina* sp. (forest flycatcher), 152
- Storm petrel, 92, 99
- Streptopelia risoria* (ringed dove), 197
- Sturnella magna* (meadowlark), 119
S. neglecta, 7, 8, 119
- Sturnus vulgaris* (European starling), 17, 99, 129, 222
- Success, reproductive, and testosterone level, 17–18
- Sula dactylatra* (masked booby), 225
S. nebouxii (booby), 222
S. sula (red-footed booby), 225
- Sylvia communis* (white-throat sparrow), 57, 65
- Tail
 shape, 146
 types, 145
- Territory
 and breeding, 15
 floaters, 13–14
 owners, 13–14
 size, 97–99
- Testosterone
 and age, 14–15
 and breeding
 density, 17
 ecology, 11–14
 season, 4
 cost of high level, 28–32
 function in birds, 3
 implant, experimental, 30–32
- Testosterone (*cont.*)
 level
 and age, 14–15
 and breeding ecology, 11–14
 costly to male birds when high, 28–32
 and interaction, social, 26–28
 and reproductive success, 17–18
 manipulation, 15–16
 and mating system, 16–17
 plasma profile during mating season, 6
 and polygny, 1–41
 profile, 4–11
 secretion control, *see* Challenge hypothesis
 and success, reproductive, 17–18
- Tetrao urogallus* (capercaillie), 7, 8, 29
- Thermoregulation and bird coloration, 146
- Thunder Cape Bird Observatory, 54
- Trapping, 50–52
 and banding birds, 52
 of landbirds, 50–52
 standardizations, 51
- Troglodytes aedon* (house wren), 92, 109, 236
T. troglodytes (wren), 65
- Turdus merula* (blackbird), 156
- Turkey, wild, 5
- Tympanuchus phasianellus* (sharp-tailed grouse), 15
- Unprofitability of prey, 155
- Vespa orientalis* (hornet), 152
- Vidua* sp. (whydah), 154
- Voice discrimination in birds, 115–139
 neighbor-stranger, 117–120
 recognition, individual, 120–133
- Voice, individual, 123–125
 habituation, 127–129
 experimental, 128
 pitch, 126
 recognition in the field, 126–132
 experiments, 126–129
- Warning coloration, *see* Aposomatism
- Warning signal hypothesis, 159–160
 and flash pattern, 160
- Waterbirds, 45
- Wilsonia citrina* (hooded warbler), 98, 119, 122, 123
- Wisconsin checklist project, 54
- Wood warbler, 146, 147

Xanthocephalus xanthocephalus, see
blackbird, yellow-headed

Zonotrichia albicollis (white-throated
sparrow), 116, 118

Zonotrichia albicollis (white-throated
sparrow) (cont.)

Z. leucophrys (white-crowned spar-
row), 17, 19, 27, 92, 118

Zosterops lateralis chlorocephala, 119