

Index

A

- Abel and Becker Cognitions Scale (ABCS), 362
- Acute stress disorder
 - avoidance symptoms category, 187
 - categories, 186
 - depersonalization, 187
 - derealization, 187
 - diagnosis, 186
 - diagnostic features, 181–182, 186
 - dissociative responses, 187
 - dissociative symptoms category, 187
 - DSM-5, 187
 - duration, 186
 - duration of symptoms, 187
 - negative mood category, 187
 - prevalence, 188
 - symptoms, 186
- Adaptive and maladaptive coping, 494
- Addiction Severity Index (ASI), 332
- Adjusting assessment procedures, 230, 231
- Adjustment disorder
 - definition, 189
 - diagnostic features, 181–182
 - prevalence, 189–190
 - stressor, 189
 - symptoms, 189
 - time limit, 189
- Adlerian psychotherapy, 299
- Affective empathy, 46
- Afifi, T.O., 202
- Ageism, 470
- Age-related macular degeneration (AMD), 471
- Agoraphobia, 132, 140
- Akinesia, 82
- Aklin, W.M., 20
- Alcohol
 - Black and Latino users, 340
 - in moderation after treatment, 333
 - prevalence, 339
 - SIP-AD, 332
 - substance-use, 329
- Alcohol-induced persisting dementia, 113
- Alzheimer's disease (AD), 103, 104, 108, 116, 117, 123, 124, 226, 459
- American Academy of Pediatrics, 394
- American Psychiatric Association (APA), 9, 30, 216, 223
- Anderson, J.O., 361
- Andreski, P., 202
- Anorexia nervosa (AN)
 - ARFID, 378
 - BMI, 376
 - body weight, 376
 - characterization, 376
- Antisocial PD, 300
- Antisocial personality disorder, 62
- Anxiety
 - ADIS, 141
 - advantages and disadvantages, 136
 - agoraphobia, 132, 140
 - avoidance and escape behaviors, 135
 - BATs, 139
 - biased thinking and avoidance behaviors, 147
 - childhood and adolescent, 143
 - cognitions, behaviors and feelings, PD, 130
 - cognitive-behavioral model, 129
 - cognitive distortions, 147
 - cultural factors, 144
 - development and maintenance, 129, 130
 - diagnostic interviewing skills, 149

- Anxiety (*cont.*)
- differential diagnosis, 141, 145, 146
 - emotion-focused coping, 142
 - emotions, 147
 - evidence-based assessment, 134
 - and fear, 129
 - formal assessment
 - ADIS-5, 136
 - behavioral observation, 139
 - DIAMOND, 137
 - informants, 137
 - MINI, 136
 - SCID-5, 136
 - self-monitoring, 137
 - self-report, 137
 - functional impairment, 149
 - GAD, 131
 - heterogeneity, 146, 147
 - OCD, 141
 - older adults, 142, 143
 - panic attacks, 140
 - PD, 131
 - perinatal depression, 145
 - physical sensations, 135
 - race, 143, 144
 - SCID, 140
 - selective mutism, 133, 134
 - self-monitoring, 140
 - self-reports, 138, 140
 - semi-structured interviews, 134
 - separation, 133
 - social, 132
 - socioeconomic status, 144, 145
 - specific phobias, 132, 133
 - structured interviews, 134
 - symptoms, 135
 - transdiagnostic factors, 146
 - uncertainty and rumination, 149
 - unstructured interviews, 134, 135
 - worry-thoughts, 141
- Anxiety and Related Disorders Interview Schedule for DSM-5 (ADIS-5), 43, 136, 166, 458
- Anxiety Disorders Interview Schedule, 440
- Anxiety Disorders Interview Schedule for DSM-IV, Child and Parent version (ADIS-IV-C/P), 448
- Anxious distress, 232, 233
- Anxious distress specifier, 225
- Apgar score, 62
- Arthritis, 467
- Asmundson, G.J., 202
- Assessment
 - biological, 334
 - dimensional approach, 331
 - substance-use, 327
 - tools, 441
 - variation, in assessment environments, 327
- Auster, S., 46
- Austin, S.B., 200
- Autism Spectrum Disorder (ASD), 29
- Avoidant and restrictive food intake disorder (ARFID), 29, 378
- Avoidant PD, 302
- B**
- Bachem, R.C., 189
- Bailey, R.W., 46
- Bailey, T.D., 401–421
- Baker, F.M., 33
- Beck Depression Inventory-II (BDI-II), 220
- Beck Depression Inventory—Primary Care (BDI-PC), 221
- Beck, A.T., 33, 138, 305
- Behavioral avoidance tasks (BATs), 139
- Behavioral observations, 461
- Behel, J.M., 481–501
- Bell, C.C., 33
- Bemak, F.P., 18, 22
- Benjamin, L., 301
- Bergeron, L., 47
- Berthiaume, C., 47
- Billingsley, A.L., 129–149
- Binge eating disorder (BED), 29, 375
 - ICBs, 377
 - marked distress, 377
 - therapist, 383
- Biopsychosocial assessment, 230
- Bipolar disorders
 - age, 254
 - antidepressant medication, 253
 - assessment
 - screening measures, 250–251
 - semi-structured interview, 247–249
 - standardized format, 246
 - structured interviews, 249
 - symptoms, 246
 - bipolar I disorder, 240, 241
 - bipolar II disorder, 241, 242, 253
 - color blind, 254
 - community health center, 252
 - cyclothymic disorder, 242
 - description, 239, 240
 - diagnosis, 255
 - factors, 254
 - HCL-32, 253
 - hypomanic episodes, 243, 253

- low mood, 252
- major depressive episodes, 243, 252
- medical condition, 243
- medication, 253
- medication's side effects, 252
- mental health clinic, 251
- minority patients, 254
- monogamous relationship, 252
- multiple periods, 251
- patient's religious beliefs and practices, 254
- procedures, gathering information
 - comorbid conditions, 245
 - decision-making process, 245
 - depressive episodes vs. hypomanic/
manic episodes, 245
 - diagnostic impression, 246
 - in-person diagnostic interview, 245
 - interview, 245
 - interviewer, 245
 - patient's significant, 246
 - time constraints/situational factors, 246
- race, 254
- recommendations, 256
- recreational/prescription drug, 252
- religiosity and spirituality, 254, 255
- SCID for DSM-5, 253
- specifiers, 244–245
- substance/medication-induced bipolar and
related disorder, 242, 243
- symptoms of anxiety, 253
- unspecified, 244
- Bipolar I disorder, 240, 241
- Bipolar II disorder, 241, 242, 253
- Bipolar spectrum diagnostic scale (BSDS), 251
- Blacker, D.E., 192
- Blake, D.D., 193
- Bleuler, M., 263
- Blevins, C.A., 195
- Bluthenthal, R.N., 340
- Bockian, N., 293–321
- Body dysmorphic disorder (BDD), 162
- Body mass index (BMI), 376
- Bogaerts, S., 364
- Bohnert, A., 30
- Bohrer, B.K., 375–396
- Bondi, M.W., 112
- Borderline PD, 301
- Boyer, S.M., 401–421
- Boyes, A., 363
- Bradykinesia, 82
- Bradyphrenia, 83
- Brand, B.L., 401–421
- Breslau, N., 202
- Bresnick, M.G., 188
- Breton, J., 47
- Brief Negative Symptoms Scale (BNSS), 268
- Brief Psychiatric Rating Scale (BPRS), 268
- Broca's aphasia, 81
- Brown, L.M., 179–206
- Bruni, T., 378
- Bulimia nervosa (BN)
 - ICBs, 377
 - LOC, 377
- Burke, W.F., 308
- Burns, R.M., 201
- Bush, S.S., 115
- Byers, A.L., 455
- C**
- Caldarone, L.B., 201
- Camberwell Assessment of Need for the
Elderly (CANE), 230
- Cambridge Depersonalization Scale (CDS), 410
- Capgras syndrome, 86
- Carleton, R.N., 138
- Cartesian mind-body dualism, 481
- Cash, T.F., 161
- Center for Epidemiological Studies-
Depression Scale (CES-D), 220
- Cerebrovascular accident (CVA), 107
- Chambless, D.L., 138
- Chapa, D.A.N., 375–396
- Charles Bonnet syndrome, 88
- Charney, D.S., 193
- Cheng, J., 363
- Child and Adolescent Psychiatric
Assessment, 446
- Child Behavior Checklist, 442
- Children
 - avoidance behaviors, 439
 - behavioral concerns, 431
 - behavioral observations, 441
 - caregivers, 434, 452
 - clinicians, 431
 - cognitive and emotional development, 427
 - contextual factors, 428
 - cost of services, 432
 - diagnostic assessment, 435
 - diagnostic interview, 433, 434
 - direct observation, 435, 436
 - environment, 435
 - generalized anxiety disorder, 438
 - healthcare or mental health, 436
 - interview, 430, 437
 - interview process, 428
 - mental health concerns, 427
 - mental health disorders, 428

- Children (*cont.*)
 motivation, 428
 observations, 436
 observing behavior, 435
 open-ended conversation, 439
 oppositional behavior, 430
 parent behaviors, 435
 parent/guardian, 431
 reference materials, 450
 referral question, 430
 semi-structured interview, 441
 service providers, 432
 speaking, 436
 structured diagnostic interview, 437
 symptom presentation, 428
 teachers, 431
 treatment process, 434
 young, 430
- Children's Yale-Brown Obsessive-Compulsive Scale (CYBOCS), 166
- Chiu, W.T., 185
- Chorea, 82
- Christine's symptoms, 464
- Chronic alcohol abuse, 113
- Chronic obstructive pulmonary disease (COPD), 142
- Chung, R.C., 18, 22
- Circadian rhythms, 473
- Client symptomatology, 32
- Clinical Assessment Interview for Negative Symptoms (CAINS), 268
- Clinical interview
 assessment procedure, 30
 client diversity, 4
 emergency and crisis settings, 4, 5
 emotional reactions, 26
 empathy, 14
 establishing rapport, 13, 14
 functioning, 4
 goals, 24
 interpersonal (*see* Interpersonal skills)
 interview formats, 30
 jail, prison, corrections and courthouse settings, 7–9
 medical settings, 6, 7
 mental disorders, 25
 multiple hypotheses, 25
 opinions/values, 26
 opportunity, 25
 outpatient mental health settings, 5, 6
 panic attacks, 25
 paying attention to language and avoiding jargon, 15
 psychotherapeutic process, 3
 reflection statement, 14, 15
 self-disclosure, 16, 17
 structured vs. non-structured, 25
 technical (*see* Technical skills)
 using humor, 16
- Clinical severity rating (CSR), 448
- Clinician-Administered PTSD scale (CAPS), 193, 194
- Clock drawing test, 97
- Cocaine use, 333
- Cognition and cognitive testing, 489
- Cognitive assessments, 441
- Cognitive-behavioral models, 129
- Cognitive-behavioral therapy, 441
- Cognitive empathy, 46
- Cognitive processing therapy (CPT), 206
- Cohen, M., 188
- Color blind, 254
- Comorbid anxiety disorder, 225
- Comorbid disorders, 202
- Complex PTSD, 203
- Complex trauma (CT), 411
- Complicated grief, PTSD, 203, 204
- Compo, N.S., 44
- Composite International Diagnostic Interview Version 3.0 (CIDI), 249
- Concussion, 106
- Conelea, C.A., 161
- Confidentiality
 age, 10
 APA ethical principle, 9
 clinician-client relationship, 10
 duty to warn and protect, 11, 12
 informed consent, 9
 limitations, 9
 temptation to discuss cases, 12, 13
 written records, 10, 11
- Connolly, S.L., 239–257
- Consultation and liaison service, 482
- Corliss, H.L., 200
- Cornell Scale for Depression in Dementia (CSDD), 221
- Countertransference, 306–309
- Cox, B.J., 202
- Criminal justice system, 328
- Criminal Sentiments Scale (CSC), 362
- Cukor, J., 185
- Cultural formulation, 21
- Cultural formulation interview (CFI), 20, 229
- Current procedural terminology (CPT), 497
- Cushing's Syndrome, 243, 463
- Cyclothymic disorder, 217, 242

D

- Dana, R.H., 18
- Dancu, C.V., 194
- Daughters, S.B., 325–342
- Davis, G.C., 202
- Davis, M.L., 155–175
- Davis, M.T., 195
- de Vogel, V., 364
- de Vries, M., 364
- Degenerative joint disease (DJD), 490
- Delusions, 264, 269–272
- Dementia, 103–105, 107–109, 113, 122–125
- Demler, O., 185
- Department of Defense (DoD), 201
- Dependent PD, 302–303
- Depression, 190
- Depressive and bipolar related disorders, 279
- Depressive disorders
 - adjusting assessment procedures, 230, 231
 - age, 226
 - assessment, 223, 224, 226
 - assessment and treatment, 233, 234
 - biopsychosocial assessment, 230
 - cultural formulation
 - CFI, 229
 - client-clinician relationship, 228
 - cultural identity, 228
 - DSM-IV*, 228
 - immigration status, 228
 - mood symptoms, 228
 - psychosocial environment and levels of functioning, 228
 - standardized tools, 228
 - zero-sum model, 228
 - description, 213
 - differential diagnosis, 215, 231–232
 - DMDD, 214
 - DSM-IV-TR*, 213
 - dysthymia, 217
 - limited assessment, 219
 - MDD, 215–217
 - medical condition, 219
 - minority older adult with MDD, 224, 225
 - pathomechanisms, 214
 - PDD, 217
 - PMDD, 217–218
 - primary care patient, 225, 226
 - procedures, gathering information
 - clinician/mental health practitioner, 220
 - clinician-administered measures, 223
 - direct observation, 223
 - evidence-based perspective, 220
 - informal assessment, 220
 - interview, 220
 - mental health organizations, 220
 - outcomes/clinical change, 223
 - screening measures, 220, 221
 - self-monitor depressive symptoms, 223
 - self-report measures, 220
 - self-report screening measures, 221, 222
 - semi-structured interviews, 222, 223
 - structured interviews, 222, 223
 - vs. psychiatric conditions, 231–232
 - race and ethnicity, 226
 - rapport building, 226, 227
 - socioeconomic status, 226
 - specifiers
 - anxious distress, 232, 233
 - atypical features, 232
 - clinical status, 232
 - melancholic features, 232
 - mood episode, 232
 - psychotic features, 233
 - severity, 233
 - symptom, 233
 - standardized diagnostic and screening tools, 227, 228
 - substance/medication-induced depressive disorder, 218
 - unspecified, 219
- Depressive episode with insufficient symptoms, 219
- Depressive PD, 305
- Depressive symptoms, 225, 469
- Desire disorders, 351
- Detailed Assessment of Posttraumatic Stress (DAPS), 410
- Diagnostic and Statistical Manual of Mental Disorders (DSM-5), 29, 458
- Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV), 161
- Diagnostic interview, 427, 451
- Diagnostic Interview for Anxiety, Mood, and OCD and Related Neuropsychiatric Disorders (DIAMOND), 42, 137
- Diagnostic Interview for Children and Adolescents, 447
- Diagnostic Interview Schedule (DIS), 36, 37
- Diagnostic Interview Schedule for Children – Fourth Edition, 444
- Diagnostic interviewing, 434
- Dieting behaviors, 393
- Difede, J., 185
- Diffuse axonal injury (DAI), 107
- Dimensional approach, 329, 331, 332, 334
- Dimensional Obsessive-Compulsive Scale (DOCS), 167

- Disorganized speech, 272
- Display paranoia-like symptoms, 254
- Disruptive mood dysregulation disorder (DMDD), 29, 214
- Dissociative disorders (DDs)
- African American male, 412–413
 - assessment, 404
 - behavioral indicators, 404
 - CT-based problems, 411
 - DDIS, 410
 - depersonalization and derealization, 408
 - development, 401
 - diagnosis, 405, 415
 - dissociative symptoms, 402, 416
 - DSM-5, 403
 - emotion regulation, 413
 - epidemiological research, 403
 - fluctuations, skills, 409
 - 4-D model, 402
 - memory and amnesic gaps, 415
 - memory disruption, 407
 - MID, 410
 - OSDD, 408
 - passive influence, 408
 - prevalence rate, 403
 - PTSD, 403, 409, 416
 - SCID-D, 404, 405
 - SCID-D-R, 405
 - self-report measures, 409
 - semi-structured interviews, 404
 - somatoform symptoms, 408
 - symptoms, 404
 - therapeutic relationship, 407
 - TOMM, 419
 - trauma, 419
 - trauma processing, 412
- Dissociative Experiences Scale (DES), 410
- Dissociative identity disorder (DID), 403
- Ditlevsen, D.N., 199
- Diversity
- clinical diagnosis, 20–22
 - cultural identity, 17
 - culture, 17
 - multicultural competence, 18
 - therapeutic relationship, 18, 19
- Dolan, K., 334
- Domino, J.L., 195
- Donders, J., 116, 117, 125
- Dorahy, M.J., 409
- Draguns, J.G., 22
- Drayton, A., 378
- Drugs
- classes, 329
 - Drug Abuse website, 329
 - drug cravings, 336
 - drug-related crimes, 328
 - medical care and drug treatment, 335
 - screens, 336
 - with SUDs (*see* Substance use disorders (SUDs))
- Dunn, E.J., 46
- Dwyer, L., 468
- Dysarthria, 81
- Dysthymia, 217
- Dystonia, 82
- E**
- Eating disorder examination (EDE), 386, 388
- concepts and diagnostic criteria, 388
 - EPSI-CRV, 389
 - factor structure, 388
 - sub-scales, 388
- Eating Disorder Inventory 3 (EDI-3), 387
- Eating disorders (EDs)
- African American persons, 391
 - AN, 376
 - ARFID, 378
 - assessment, 390
 - athletes, 394
 - BED, 375, 377
 - BN, 376
 - diagnoses, 375, 393
 - DSM-5, 375
 - EDI-3, 387
 - EPSI, 387
 - ethnicity and culture, 390
 - ethnic-racial minority patients, 391
 - exercise, 385
 - gender, 392
 - health risks and medical complications, 394
 - ICBs, 381
 - medical complications, 390, 394
 - mental health condition, 386
 - non-athletes, 394
 - OSFED, 377
 - physical examination, 389
 - prevalence, 390, 392
 - psychiatric, 386
 - psychiatric disorders, 375, 378
 - psychological assessment, 389
 - psychopathology, 386
 - SCOFF, 387
 - sexual-minority populations, 392, 393
 - sexual-minority women, 393
 - transgender experiences, 392
 - women, 391

- Eating pathology symptoms inventory (EPSI), 387
- Eating Pathology Symptoms Inventory—Clinician Rated Version (EPSI-CRV), 388
- Echolalia, 85
- Ecologic adaptation, 296
- Ecologic modification, 296
- Edelbrock, C., 30
- Edelstein, B., 455–459, 461–473
- Edwards, C.B., 129–149
- Ego-dystonic vs. ego-syntonic thoughts, 158, 159
- Ehrenreich-May, J., 427–453
- Elklit, A., 199
- Emerging professionals
 - client's ambivalence, 23
 - clinical interview, 22
 - clinical supervision and personal psychotherapy, 24
 - clinician-client relationships, 24
 - clinician's anxiety/discomfort, 22
 - consultations, 24
 - content, 23
 - diagnostic interviews, 22
 - financial assistance, 23
 - information-gathering process, 22
 - personal questions, 24
 - psychotherapy, 23
 - young clinicians, 23
- Emotional functioning, 483
- Empathic accuracy, 46
- Erbaugh, J.K., 33
- Erectile disorder (ED), 351
- Evidence-based treatment, 206
- Excoriation disorder, 163
- Executive functioning, 123
- Exum, A., 375–396
- Eye diseases, 471
- Eye movement desensitization and reprocessing (EMDR), 206
- F**
- Family Accommodation Scale (FAS), 167
- Farias, S.T., 123
- Fauerbach, J.A., 188
- Fear
 - anxiety, 129, 133, 148
 - and avoidance, 140
 - bodily sensations, 130
 - phobia, 133
 - social situations, 132
- Fear of fear model, 131
- Feliciano, L., 213–234
- Female orgasmic disorder (FOD), 351
- Female sexual arousal disorder (FSAD), 351
- Fetal alcohol syndrome, 122
- Fingerhut, R., 13
- First, M.B., 37, 38, 192
- Fleet, C., 202
- Fleisher, C.L., 188
- Flessner, C.A., 161
- Flint, M., 55–75
- Florida Obsessive-Compulsive Interview (FOCI), 167
- Foa, E.B., 186, 194
- Forbush, K.T., 375–396
- Formal assessment
 - cognitive and functional impairment, 267
 - standardized interview formats, 267–269
 - thorough clinical evaluation, 267
 - tools, 443
- Freud, A., 298
- Frost, R.O., 161
- Functional assessment, 459, 460
- G**
- Gaher, R., 138
- Gara, M.A., 274
- Gastrointestinal (GI) complications, 389
- Gauvin, S., 350–369
- Gender dysphoria (GD), 352
 - in children and adolescents/adults, 353
 - comprehensive psychosocial assessment, 356
 - diagnosis, 352
 - and paraphilic disorders, 367
- General behavior inventory (GBI), 251
- General clinical health psychology interview, 489
- Generalized anxiety disorder (GAD), 131
- Generalized Anxiety Disorder-7 items (GAD-7), 225
- Genetic malformations, 122
- Genitopelvic pain/penetration disorder (GPPPD), 352
- Geriatric Anxiety Inventory, 142
- Geriatric Anxiety Scale, 142
- Geriatric Depression Scale (GDS), 221, 464
- Geriatric Depression Scale-15 (GDS-15), 491
- Geriatric Suicide Ideation Scale, 459
- Glasgow Coma Scale (GCS), 121
- Goodman, W.K., 155–175
- Gore, W.L., 305
- Grau, K., 214
- Greenbaum, P.E., 360

Gress, C.L.Z., 361
 Grisham, J., 161
 Grossman, R.A., 427, 428, 430, 431, 433–448,
 450–452
 Groth-Marnat, G., 47
 Gum, A.M., 213–234
 Gusman, F.D., 193

H

Haag, A.M., 363
 Hair-pulling disorder, 162
 Haley, W.E., 471
 Hallucinations, 264, 269–272
 Halpern, J., 46
 Hamilton Rating Scale for Depression
 (HRSD), 221, 223
 Hanson, R.K., 360, 362, 363
 Haralambous, B., 466
 Hardy, K.V., 367
 Hare, R.D., 362
 Harm reduction programs, 328
 Harvard Trauma Questionnaire-Part IV
 (HTQ-IV), 199
 Hays, P.A., 413
 Health care setting, 482
 Health psychology
 biopsychosocial approach, 495
 cancer, 486
 cultural factors, 495
 evaluation process, 482
 gross cognitive impairments, 494
 health care settings and situations, 482
 integrated primary care, 483
 LGBTQ, 496, 497
 medical diagnoses, 485
 population-based approach, 484
 procedure, 487
 psychological evaluations, 482
 referral question, 483
 rehabilitation program, 484
 religion, 495
 settings and populations, 482
 sexuality, 496
 training, 481
 Hemiparesis, 82
 Herbs and dietary supplements (HDS), 71
 HIPAA-compliant databases, 248
 Histrionic PD, 301
 Hoarding disorder, 162
 Homicidal thoughts, 87
 Hopkins verbal learning test (HVLT), 465
 Hormone therapy, 357
 Horowitz, M.J., 203
 Hospital bound behaviors, 499

Houston, W.S., 112
 Hrabosky, J.I., 161
 Huntington's disease, 82, 105
 Hypoactive sexual desire disorder (HSDD), 351
 Hypochondriasis, 86
 Hypomania checklist (HCL-32), 250
 Hypomanic episodes, 214, 241–246, 248, 250,
 252–255
 Hypotheses, 470
 Hysteria disorders, 183

I

Inappropriate compensatory behaviors
 (ICBs), 377
 Informants, 313
 Instrumental activities of daily living (IADLs),
 92, 103
 Integrated primary care, 483
 Integumentary system, 389
 International Classification of Diseases
 (ICD-10), 458
 International Classification of Diseases,
 Eleventh Revision (ICD-11), 331
 Interpersonal skills
 empathy, 46
 rapport, 44–46
 Intersubjectivity, 413, 414
 Interviewing, schizophrenia
 clinical expertise, 287
 diagnosis, 287
 don't apologize for questions, 286
 empathic and patient approach, 286
 primary care setting, 327
 PRISM-5, 334
 specific time periods/events, 287
 veracity of, psychotic experience, 286
 Intra-interview behaviors, 407
 Ivey, D.C., 471

J

Jacobson, J.O., 340
 James, J.W., 471
 Jimenez, D.E., 228
 Johanson, K.A., 213–234
 John, S.E., 77–101
 Johnston, F.E., 44
 Julien, R.M., 333
 June, A., 3–26

K

Kaloupek, D.G., 193
 Karney, B.R., 201

Kassam-Adams, N., 188
 Katz, E., 455–459, 461–473
 Kaufman Assessment Battery for Children, 442
 Keane, T.M., 193, 195, 203
 Keith-Spiegel, P., 13
 Kelley, J.M., 46
 Kernberg, O., 308
 Kessler, R.C., 185
 Keuthen, N.J., 161
 Kiddie Schedule for Affective Disorders and Schizophrenia for DSM-5 (K-SADS-5), 40, 41
 Kimber, J., 334
 Knapp, S.J., 13
 Knight, R.M., 378
 Kobak, K.A., 41
 Koenen, K.C., 200
 Koocher, G.P., 13
 Korsakoff's syndrome, 113
 Koster, K., 364
 Kraepelin, E., 263
 K-SADS-PL (Present and Lifetime version), 40

L

Laboratory tests, 462
 and medical records, 462
 Lang, P.J., 138
 Language functioning, 123
 Larrabee, G.J., 115
 Laszloffy, T.A., 367
 Laws, D.R., 360, 361
 Lesbian, gay, and bisexual (LGB), 392
 Lesbian, gay, bisexual, transgender, and queer (LGBTQ), 200, 496
 Liebowitz, 138
 Life event checklist (LEC)
 trauma-related self-report measures, 194, 195
 Likert-type scale, 33
 Litz, B.T., 195
 Loewenstein, R.J., 404, 405
 Lonner, W.J., 22
 Lorenz, L., 189
 Loss-of-control (LOC), 376
 Lovibond, P.F., 138
 Lovibond, S.H., 138
 Lupesko-Persky, O., 179–206

M

MacNeil, A., 363
 Maercker, A., 189
 Magidson, J.F., 325–342
 Major depressive disorder (MDD), 189

APA, 216
 classification, 216
 emotional/physical functioning, 215
 grief and bereavement, 216
 manic/hypomanic episodes, 215
 medical settings, 216
 minority older adult, 224, 225
 palliative care recipients, 216
 prevalence, 216
 psychosis, 216
 specifiers, 216
 symptoms, 215, 216
 Major depressive episodes (MDEs), 215–217, 219, 232, 243, 252
 Major/mild neurocognitive disorder, 104
 Malingering, 204
 Mania, 240, 242–246, 248, 255
 Manic episodes, 214, 240–246, 248–250, 254, 255
 Marx, B.P., 195
 Mattia, J.I., 32
 McGoldrick, M., 313
 McIngvale, E., 155–175
 McKibben, J.B., 188
 McPhail, I.V., 361
 Medication-assisted treatment (MAT), 327
 Mendelson, M., 33
 Mental disorders, 29
 Mental health literacy, 227
 Mental Health Parity and Addiction Equity Act (MHPAEA), 29
 Mental health professionals, 3, 356
 Mental health vs. primary care behavioral health, 485
 Mental status examination (MSE), 458, 459
 appearance, 79, 80
 arrival, 79
 assessment
 procedures, gathering information, 96, 98
 behavior and psychomotor activity, 81, 82
 client clock drawing, 97
 client's symptoms, 78
 cognition
 abstract reasoning, 94
 adaptive and social functioning, 92
 attention, 92
 intellectual functioning, 91, 92
 memory, 93, 94
 orientation, 91
 components, 78
 comprehensive examination, 78
 descriptions, 77
 emotion
 affect, 89, 90
 mood, 88, 89

- Mental status examination (MSE) (*cont.*)
 evaluation, 79, 100
 factors
 insight and judgment, 95, 96
 reliability, 95
 impression, 78
 integrative biopsychosocial approach, 100
 manner of relating and attitude, examiner,
 82, 83
 observations, 78, 79, 82, 83
 patient-centered approach, 100
 populations
 aging, 98, 99
 cultural considerations, 99
 settings, 98
 speech and language
 anomia, 81
 Broca's aphasia, 81
 content, 80
 dysarthria, 81
 hypophonia, 80
 monotone, 80
 premorbid functioning and educational/
 occupational attainment, 81
 prosody, 80
 rate, 80
 tone, 80
 volume, 80
 structured, 100–101
 thinking
 perception, 87, 88
 suicidal and homicidal ideation, 87
 thought content, 85, 86
 thought process, 83–85
- Metaphor, 3
 Meyer, T.J., 138
 Michigan alcohol screening test (MAST), 497
 Mild cognitive impairment (MCI), 103, 104, 498
 Miller, C., 29–48
 Miller, C.J., 239–257
 Millman, Z.B., 261–288
 Millon, T., 295, 297, 314, 315
 Mindfulness, 308, 309, 320
 Mini International Neuropsychiatric Interview
 for Children and Adolescents, 444
 Mini International Neuropsychiatric Interview
 for DSM-5 (MINI-5), 41, 42
 Mini-International Neuropsychiatric Interview
 (MINI), 136, 249, 388
 PTSD, 192, 193
 trauma, 192, 193
 Mini-International Neuropsychiatric Interview
 7.0 (M.I.N.I.), 458
 MINI-Kid, 42
- Minimal clinical training, 43
 Mini-Mental State Exam (MMSE), 94, 97
 Minnesota Multiphasic Personality Inventory
 (MMPI), 312, 442
 Minnesota Sex Offender Screening Tool
 (MnSOST), 360, 363
 Minority older adult, MDD, 224, 225
 Mock, J.E., 33
 Montgomery Asberg Depression Rating Scale
 (MADRS), 221
 Montgomery Asberg Depression Rating
 Scale-Short form (MADRS-S), 220
 Montreal Cognitive Assessment (MoCA), 94, 97
 Mood disorder questionnaire (MDQ), 250
 Mood disorders, 213, 226, 229, 233
 Mood effects, 243
 Moral injury, PTSD, 204
 Morris, L.B., 301, 305
 Morris, N., 378
 Morrison, J., 45
 Motivational Interviewing approach, 297
 Multiaxial system, 29
 The Multidimensional Inventory of
 Development, Sex, and Aggression
 (MIDSA), 360
 Multidimensional Inventory of Dissociation
 (MID), 410
 Multidimensional Pain Inventory (MPI), 485
 Multiphasic Sex Inventory (MSI), 360
 Multiple sclerosis, 243
 Multiscale Dissociation Inventory (MDI), 410
 Murdock, T., 186
 Muscle dysmorphia, 162
- N**
 Nagy, L.M., 193
 Narcissistic PD, 301–302
 Narrow assessment tools, 442
 National Stressful Events Survey Acute Stress
 Disorder Short Scale (NSESSS)
 trauma-related self-report measures,
 195, 196
 Natural curiosity, 3
 Negative symptoms, 265
 NetSCID-5, 40
 Neurocognitive disorder, 103, 108, 122–125
 Neuropsychological assessment interview
 advantages, 110, 111
 client and caregiver, 117
 cognitive impairments/motivations, 117
 confirmatory bias, 116
 daily functioning, 115
 dementia, 122–125

- developmental history, 113, 114
- factors, 110
- health care system, 117
- hypothesis-testing approach, 116
- interruption, 118
- legal history, 114, 115
- medical history, 111, 112
- mental health, 112, 113
- model, 110
- neurocognitive disorder, 122–125
- occupational history, 114
- rambling, 118
- rapport, 115–118
- referrals, 118–120
- sources, 109
- styles, 110
- substance use, 113, 117
- TBIs, 120–122
- test hypotheses, 109
- valuable behavioral observations, 116
- Neuropsychology
 - assessment (*see* Neuropsychological assessment interview)
 - behavioral observations, 104, 105
 - cognitive/brain dysfunction, 103
 - cognitive symptoms, 107, 108
 - factors, 108
 - goal, 103
 - medical/neurological history, 105–107
 - patients and clinicians, 103
 - referrals, 108
- Newman, M.G., 138
- Nijenhuis, E.R., 183
- Ninan, P.T., 161
- Nine-Item Patient Health Questionnaire (PHQ-9), 221
- Non-interview-based assessment methods, 456
- Non-systematic/idiosyncratic approach, 32
- North, C.S., 32
- Nosological systems, 32, 35
- O**
- O'Donohue, W., 138
- O'Toole, G., 44
- Obsessive-compulsive disorder (OCD)
 - age, 171
 - anxiety disorder, 141
 - assessment tools, 155, 161
 - BDD, 162
 - body dysmorphic-like disorders, 164
 - body-focused repetitive behavior disorders, 164
 - contamination, 156
 - content themes, 156, 157
 - description, 156
 - diagnostic criteria, 171–173
 - diagnostic interviewing, 155, 164, 174
 - differential diagnoses, 172
 - evidence-based assessment, 164–166, 175
 - excoriation disorder, 163
 - factors, 155
 - formal assessment
 - child/parent-specific measures, 167
 - clinician-rated instrument, 166
 - family assessment and gathering information, 167
 - self-report measures, 167
 - semi-structured clinical interviews, 164
 - gender, 170
 - hoarding disorder, 162
 - mind elements, 174
 - patient's level, 173
 - race/culture, 170, 171
 - responsibility for harm, 156, 158
 - streptococcal infections, 164
 - structured assessment, 169
 - substance/medication-induced disorder, 163
 - symmetry/just right-ness, 160
 - symptoms, 163, 173
 - thought content, 169
 - trichotillomania, 162, 163
 - unacceptable, taboo thoughts
 - ego-dystonic and ego-syntonic, 158
 - immoral thoughts, 160
 - neutralize, 158
 - sexual thoughts, 159, 160
 - stigma and shame, 158
 - violent thoughts, 160
 - unstructured clinical interview, 168, 169
- Obsessive-Compulsive Inventory-Revised (OCI-R), 167
- Obsessive-compulsive PD, 303
- Obstructive sleep apnea (OSA), 498
- Office Mental Status Exam for Dissociation (OMSE), 405
- Olden, M., 185
- Older adults
 - ADIS-V, 458
 - adverse drug effects, 468
 - age cohort, 473
 - anxiety disorders, 455
 - assessment method, 456
 - cancer diagnoses, 467
 - clinical interview, 456–457, 471
 - cognitive effort, 461
 - cognitive impairment, 459
 - cognitive tasks, 473

- Older adults (*cont.*)
 depressive symptoms, 463
 DSM, 469
 DSM-5 and ICD-10, 470
 factors, 456
 functional assessment, 459
 GDS, 464, 465
 hearing, 471, 472
 hearing loss, 472
 instructions, 472
 late-life depression, 469
 LGBT, 466
 medical conditions, 467
 medical history, 462
 medical records, 457
 memory and cognitive changes, 472
 mental disorders, 456
 mental illness, 464
 mental status examination, 458
 mood disorders, 455
 SCID, 458
 structured interviews, 457
 unstructured interviews, 456
 US population, 455
 vitamin D deficiency, 462
- Oldham, J.M., 301, 305
 Olfactory reference syndrome, 164
 Oltmanns, T.F., 469
 Operation Enduring Freedom (OEF), 190, 201
 Operation Iraqi Freedom (OIF), 190, 201
 Oppositional defiant disorder, 214
 Orgasmic disorders, 351
 Osborn, C.A., 360
 Osilla, K.C., 201
 Outline for Cultural Formulation (OCF), 202
- P**
 Palmieri, P.A., 195
 Paniagua, F.A., 19, 22
 Panic disorder (PD), 130, 131
 Panic Disorder Severity Scale, 140
 Paquette, C.E., 325–342
 Paranoid PD, 299–300
 Paraphilia, 353
 Paraphilic disorders, 353, 355
 forensic setting, 358
 prevalence, 355
 questionnaires, 359
 specifiers, 355
 Paresthesia, 82
 Parkinson's disease, 83, 123, 467
 Passive-aggressive (negativistic) PD, 303–304
 Pathomechanisms, 214
 Patient health questionnaire-9 (PHQ-9), 225–227
 Pedersen, P.B., 22
 Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcal Infections (PANDAS), 164
 Pediatric psychiatric disorder, 214
 Peer, J.E., 261–288
 Peltier, M.R., 274
 Period limb movement disorder (PLMD), 498
 Perkonig, A., 185
 Persistent depressive disorder (PDD), 217
 Personality assessment, 442
 Personality Assessment Inventory (PAI), 486
 Personality disorders (PD)
 adaptation, 296
 antisocial, 296, 300, 314
 architectural structures, 299
 assessment of, 294
 avoidant, 302
 borderline, 297, 301
 clinicians, 295, 313
 compliant-compulsive personality spectrum, 297
 dependent, 302–303
 depressive, 305
 description, 293
 diagnosis, 319
 domains, 298
 domestic violence against women, 314
 DSM principles, 313
 ego-syntonic perspective, 294
 evolutionary theory, 297
 formal assessment, 311–313
 gathering information, 306–309
 genetic basis and experience, 293
 histrionic PD, 301
 identity disturbance, 315
 inflexibility, 293
 initial intake, 294
 law enforcement professions, 294
 mild, moderate and severe psychopathology, 295
 Millon's theory, 295
 mind-altering drugs, 315
 Motivational Interviewing approach, 297
 narcissistic, 301–302
 obsessive-compulsive, 303, 314
 paranoid, 294, 299–300
 passive-aggressive (negativistic), 303–304
 pathological narcissistic personality disorder, 299
 planning and reasoning, 314
 replication, 296

- sadistic, 304
- sadistic and masochistic, 295
- schizoid, 300
- schizoid and/or avoidant, 297
- schizotypal, 300
- self-defeating, 304–305, 317
- self-defeating/masochistic spectrum, 297
- social conditions, 315
- Peterson, E., 202
- Phallometry, 361, 366
- Phillips, K.A., 161
- Phillips, M., 46
- Phobias, 132, 133
- Physical and occupational therapy, 492
- Physical symptoms, 494
- Pifer, M., 3–26
- Positive and Negative Syndrome Scale (PANSS), 268
- Positive symptoms, 264
- Posttraumatic amnesia, 121
- Posttraumatic stress disorder (PTSD), 281, 403
 - and anxiety, 182
 - armed forces, 201
 - assessment
 - CAPS, 193, 194
 - MINI, 192, 193
 - PSS-I, 194
 - SCID, 191, 192
 - trauma-related specific interviews (*see* Trauma-related specific interviews)
 - avoidance of stimuli symptom, 183
 - car accident survivor, 197
 - clinician compares, 184
 - cultural awareness, 201–202
 - cultural competency, 198
 - culture, 198
 - depersonalization, 185
 - derealization, 185
 - description, 180
 - diagnosis
 - comorbid disorders, 202
 - complex PTSD, 203
 - complicated grief, 203, 204
 - malingering, 204
 - moral injury, 204
 - symptomatology, 203
 - diagnostic criteria, 179
 - diagnostic features, 181–182
 - Do's and Don'ts, 205–206
 - DSM-IV-TR, 180
 - exposure to traumatic event, 182
 - gender and age, 199
 - intrusion symptom, 183
 - intrusive symptoms, 183
 - Marine combat veteran, 196, 197
 - marked alterations in arousal and reactivity, 184
 - mental health professionals, 198
 - negative alterations in mood and cognition, 183, 184
 - PDSQ, 193
 - pleasurable activities, 184
 - prevalence, 185, 186
 - racial and ethnic minorities, 199–200
 - sexual minority adults, 200
 - substance use/medical condition, 184
 - subsyndromal/subthreshold, 179
- Potts, S., 103–125
- Pre-interview informational handout, 429
- Premenstrual dysphoric disorder (PMDD), 217–218
- Premenstrual syndrome (PMS), 218
- Presenting problem
 - alternative and nonprofessional treatment methods, 61
 - assessment, 59, 60
 - cultural and contextual framework, 57
 - depression, 58, 59
 - diagnostic hypotheses, 56, 58
 - frequency, 58
 - inquiry, 55
 - insight, 60
 - intensity, 58
 - interviewer, 56–58, 61
 - interviewing process, 56
 - medical and psychotherapeutic techniques, 61
 - medically oriented practices, 60
 - mental disorders, 56, 60
 - panic attacks, 57
 - psychoactive substance, 56
 - psychopathology, 56
 - sleep disturbance, 60
 - stressors, 58
 - time-consuming, 61
- Primary Care Behavioral Health Model, 484
- Primary care patient, depressive symptoms, 225, 226
- Procedures, gathering information
 - assessment approaches, 490
 - clinical, 487, 490
 - DJD, 490
 - health record, 493
 - mental health services, 491
 - open ended approach, 488
 - patient's health record, 487
 - psychiatric history, 491
 - stereotypic psychologist behaviors, 488

- Profile of Mood States (POMS), 220
 Prolonged exposure therapy (PET), 206
 Prospective memory, 94
 Psychiatric diagnosis, 497
 Psychiatric diagnostic screening questionnaire (PDSQ)
 PTSD, 193
 trauma, 193
 Psychiatric disorders, 213
 Psychiatric history, 491
 The Psychiatric Research Interview for Substance and Mental Disorders (PRISM-5), 334
 Psychoactive substances, 333
 Psychoeducation, 190
 Psychological evaluations, 483
 Psychological trauma, 180
 Psychopathology, 386, 459
 and BPRS, 268
 positive, negative and general scales, 268
 Psychopathy Checklist-Revised (PCL-R), 360, 362, 363, 366
 Psychosis
 normative development, children and adolescents, 275
 risk factors, 273
 Psychosocial histories, 461
 Psychotherapy services, 253
 PTSD checklist for diagnostic and statistical manual of mental disorders (PCL)
 trauma-related self-report measures, 195
 PTSD symptoms scale interview (PSS-I)
 trauma-related specific interviews, 194
 Pukall, C.F., 350–369
- R**
- Racial and cultural factors, 466
 Ramchand, R., 201
 Reasons for Living – Older Adults scale, 459
 Recurrent brief depression, 219
 Renn, B.N., 77–101
 Repeatable Battery for the Assessment of Neuropsychological Status (RBANS), 490
 Reproductive individuation, 296
 Reproductive nurturance, 296
 Research diagnostic criteria (RDC), 35, 248
 Restless leg syndrome (RLS), 498
 Revised Children's Anxiety and Depression Scale, 442
 Richson, B., 375–396
 Riess, H., 46
 Riggs, D.S., 186, 194
 Roberts, A.L., 200
 Robinson, P.L., 340
 Rogers, R., 32, 36, 45, 47
 Rosenthal, R., 44
 Ross, L., 45
 Rothbaum, B.O., 161, 186, 194
 Rouen, D., 334
 Rush, A. Jr., 192
 Rybarczyk, B., 481–501
- S**
- Sadistic PD, 304
 Safety, 8
 Saint Louis University Mental Status exam (SLUMS), 97
 Santos, M.T., 161
 Sareen, J., 202
 Sattler, J.M., 45, 47, 103
 Sayers, L., 378
 Scharron-del Rio, M.R., 22
 Schedule for Affective Disorders and Schizophrenia (SADS), 247, 248
 Schizoaffective disorder, 280
 Schizoid PD, 300
 Schizophrenia, 82, 112, 254
 causes of disability, 262
 characteristic symptoms, 277
 clinical interview, 266
 clinical vignettes, 269
 cognitive impairment, 284
 complex, heterogeneous disorder, 261
 conceptualizations and diagnostic criteria, 263
 constellation of symptoms, 262
 depressive and bipolar related disorders, 279
 disorganized speech, 272
 duration, 278
 environmental factors, risks of, 262 (*see also* Formal assessment)
 gender, race, culture, age and diversity, 272–277
 general medical conditions, 281
 genetic factors, 262
 guarded and suspicious presentation, 284, 285
 hallucinations and delusions, 269–272
 negative symptoms, 265, 284
 patient self-report, 261
 poor insight, 282, 283
 poor social, educational and vocational functioning, 265
 positive symptoms, 264
 progressive disorder, 262

- PTSD, 281
- risk of suicide, 265
- Schizoaffective disorder, 280
- SCID-5 (*see* Structured Clinical Interview for DSM-5 (SCID-5))
 - social/occupational dysfunction, 277–278
 - substance-related disorders, 280
 - treatments, 265
- Schizotypal PD, 300
- Schneider, S.C., 155–175
- Schnurr, P.P., 195
- SCID-5-AMPD (Alternative Model for Personality Disorders), 40
- SCID-5-CT (Clinical Trials Version), 39
- SCID-5-CV (Clinician Version), 40
- SCID-5-RV (Research Version), 39
- Scientist-practitioner model, 47
- Screening measures
 - bipolar disorders
 - advantages and disadvantages, 251
 - antidepressant prescriptions, 250
 - BSDS, 251
 - CIDI, 250
 - depression diagnoses, 250
 - diagnosis, 251
 - GBI, 251
 - HCL-32, 250
 - MDQ, 250
 - psychometric properties, 251
 - SCID, 250
 - time-constrained situations, 250
 - depressive disorders, 220, 221
- Screening, Brief Intervention, and Referral to Treatment (SBIRT) model, 327–328
- Scrupulosity, 160
- Segal, D.L., 3–26, 34, 36, 47
- Selective mutism, 133, 134
- Self-defeating PD, 304–305
- Self-injurious behaviors, 409
- Self-monitor depressive symptoms, 223
- Self-report inventories, 311
- Self-report screening measures
 - depressive disorders, 221, 222
- Semi-structured interviews, 433, 446, 457
 - ADIS-5, 43
 - bipolar disorders
 - administration, 247
 - comprehensive depression module, 247
 - diagnostic criteria, 247
 - SADS, 247, 248
 - SCID, 247, 248
 - standardized, 247
 - depressive disorders, 222, 223
- DIAMOND, 42
- DIS, 36, 37
- K-SADS, 40, 41
- MINI-5, 41, 42
- SCID-5-, 37–40
- Sensory modality, 471
- Separation anxiety disorder, 133
- Severity specifiers, 233
- Sex Offender Risk Appraisal Guide (SORAG), 363
- Sex therapy, 364, 365
- Sexual desire and arousal disorders, 351
- Sexual dysfunctions
 - to biological sex, DSM-5, 350
 - clinical interview, 355
 - clinically significant issues, 350
 - comorbid nonsexual mental disorders, 350
 - four-phase sexual response cycle, 351
 - general biological factors, 355
 - “generalized”, 350
 - “lifelong”, 350
 - psychological factors, 356
 - psychometric measures, 364
 - questionnaires, 356
- Sexual functioning, 356
- Sexual History Questionnaire-Revised (SHQ-R), 360
- Sexual interest/arousal disorder (SIAD), 351
- Sexual violence, 182
- Shankman, S.A., 37
- Shay, J., 204
- Shear, M.K., 138
- Short-duration depressive episode, 219
- Shorter two-item version of the Patient Health Questionnaire (PHQ-2), 221
- Shuman, D.W., 32
- Simons, J., 138
- Single assessment methods, 456
- Skepticism, 416
- Social anxiety disorder, 132
- Social functioning, 92
- Social history, 6
 - and sexual functioning, 65, 66
 - biographical method, 62
 - chronological method, 62
 - developmental milestones, 62, 63
 - diversity, 67, 68
 - educational, 65
 - family, 63–65
 - interviewers, 62
 - legal, 68, 69
 - medical, 72, 73
 - substance use and misuse, 70–72
 - technique, 73, 74
 - transitions, 62

- Social history (*cont.*)
 trauma and abuse, 69, 70
 vocational, 66, 67
- Social justice, 18
- Social/occupational dysfunction, 277–278
- Somatization, 495
- Somatoform Dissociation Questionnaire (SDQ), 410
- Spalding, R., 455–459, 461–473
- Spielberger, C.D., 138
- St. Georges, M., 47
- Stages of Change Readiness and Treatment Eagerness Scale (SOCRATES), 328
- Standardized diagnostic interviews, 329, 330, 334
- Standardized interview formats, 449
- Stanley, M.A., 161
- Static-99, 363
- Stein, M.B., 202
- Steinberg, M., 402
- Steinman, S.A., 129–149
- Steketee, G., 161
- Stem cell/bone marrow transplants, 486
- Storch, E.A., 155–175
- Storz, S., 185
- Stress
 acute stress disorder (*see* Acute stress disorder)
 definition, 180
 PTSD (*see* Posttraumatic stress disorder (PTSD))
- Stroke, 243
- Structure Interview of Reported Symptoms (SIRS), 419
- Structured and semi-structured interviews, 311
- Structured clinical interview for DSM-5 (SCID-5), 37–40, 136, 166, 388, 458
 bipolar disorders, 247, 248
 diagnostic interview, 266
 modules, 266
 overview, 266
- Structured Clinical Interview for DSM-5 Personality Disorders (SCID-5-PD), 311
- Structured clinical interview for the DSM (SCID)
 PTSD, 191, 192
 trauma, 191, 192
- Structured interviews, 33–36, 433, 444, 457
 bipolar disorders
 CIDI, 249
 epidemiological research, 249
 MINI, 249
 depressive disorders, 222, 223
- Substance Abuse and Mental Health Services Administration's (SAMHSA), 328
- Substance use, 190
- Substance use disorders (SUDs)
 assessment, 327
 case studies, 335–337
 categorical classification, 332
 comorbidity, 326, 334
 criminal justice system, 328
 description, 325
 diagnostic assessment, 329
 dimensional approach, 331
 DSM-5, 329
 epidemiological data, 326
 harm reduction, 328
 primary care, 327
 SBIRT model, 328
 symptoms, 331
 treatment goals, 333
- Substance use symptoms, 331
- Substance/medication-induced bipolar and related disorder, 242, 243
- Substance/medication-induced depressive disorder, 218
- Substance-related disorders, 280
- Substance-use
 assessment, 333
 public health and economic consequences, 325
 rate of treatment, 326
 screening, 327
 self-report measures and biological measures, 333
 stigma, 335
- Sue, D.W., 19, 22
- Suicidality, 87
- Summerfeldt, L.J., 43
- Swaminath, S., 325–342
- Symptom clustering, 35
- Szymanski, J., 138
- T**
- Tansey, M.J., 308
- Tardive dyskinesia, 82
- Taylor, S., 138
- Technical skills
 semi-structured interviews, 31, 36–43
 structured interviews, 30, 31, 33–36
 unstructured interviews, 30–33
- Test of Memory Malingering (TOMM), 419
- Thought content
 characteristics, 84
 erotomanic delusions, 86
 grandiose delusions, 86

- nihilistic delusions, 86
 - paranoid delusions, 86
 - persecutory delusions, 85
 - referential delusions, 86
 - somatic delusions, 86
 - Thought process
 - alogia, 85
 - bradyphrenia, 83
 - characteristics, 84
 - circumstantiality, 84
 - clang associations, 85
 - echolalia, 85
 - flight of ideas, 83
 - neologisms, 85
 - perseveration, 85
 - poverty of thought, 84
 - tangentiality, 84
 - thought blocking, 84
 - word salad, 85
 - Thoughts/behavior disorganization, 264
 - Thyroid function, 463
 - Thyroid stimulating hormone (TSH), 112
 - Tickle-Degnen, L., 44
 - Tolin, D.F., 42, 161
 - Tonarely, N.A., 427–453
 - Total hip arthroplasty (THA), 490
 - Trauma, 409
 - assessment
 - MINI, 192, 193
 - PDSQ, 193
 - SCID, 191, 192
 - self-report measures (*see* Trauma-related self-report measures)
 - specific interviews (*see* Trauma-related specific interviews)
 - procedures, gathering information
 - biological factors, 190
 - client's collaboration, 190
 - cognitive and mood assessments, 191
 - collateral sources, 191
 - evaluation process, 191
 - identify, 190
 - psychological considerations, 190
 - ROI, 191
 - self-report measures, 191
 - social elements, 190
 - structured/semi-structured
 - interviews, 191
 - treatment goals, 190
 - unstructured interview, 191
 - psychological trauma, 180
 - Trauma-related self-report measures
 - LEC, 194, 195
 - NSESS, 195, 196
 - PCL, 195
 - Trauma-related specific interviews
 - CAPS, 193, 194
 - PSS-I, 194
 - Traumatic brain injuries (TBIs), 106, 120–122, 190, 243
 - Traumatic event, 182
 - Traumatic Life Events Questionnaire (TLEQ) scale, 195
 - Trichotillomania, 162, 163
 - Trimble, J.E., 22
 - Turner, S.M., 20, 138
- U**
- Uniform recording system, 34
 - Unspecified depressive disorder, 219
 - Unstructured interviews, 31–33, 433, 457
 - Urinalysis, 334
- V**
- Valla, J., 47
 - Vallano, J.P., 44
 - Van der Hart, O., 183
 - Vancouver Obsessional Compulsive Inventory (VOCI), 167
 - VandeCreek, L.D., 13
 - Vandeleur, C.L., 216, 217
 - Vandermorris, A.K., 200
 - Vanderploeg, R.D., 116, 119
 - Veale, D., 161
 - Veterans Health Care Administration (VHA) services, 200
 - Violence Risk Appraisal Guide (VRAG), 360, 363, 366
 - Violence Risk Scale: Sex Offender Version (VRS:SO), 364
 - Virden III, T.B., 55–75
 - Vitamin D insufficiency, 462
 - Voss, M.M., 375–378, 381–395
- W**
- Walsh, W., 186
 - Walters, E.E., 185
 - Walther, M.R., 161
 - Ward, C.H., 33
 - Weathers, F.W., 193, 195
 - Wechsler Individual Achievement Test, 442
 - Wechsler Intelligence Scale for Children, 441
 - Weiner, S.J., 46
 - Werth, J.L., 12
 - Widiger, T.A., 305
 - Wiechman Askay, S.A., 188
 - Williams, K.N., 34, 36, 47

- Wilson, J.P., 203
Winchel, R.M., 161
Winston, F.K., 188
Wirove, R., 363
Wittchen, H.U., 185
Witte, T.K., 195
Wolpe, J., 138
Woodcock-Johnson Tests of Achievement, 442
Woods, D.W., 161
Working Alliance Inventory (WAI), 44
World Health Organization Five-Item
Well-Being Index (WHO-5), 220
Wright, A.J., 47
- Y**
Yahav, R., 188
Yale-Brown Obsessive-Compulsive Scale
(YBOCS), 166
Yochim, B., 103–125
- Z**
Zanarini, M.C., 192
Zero-sum model, 228, 234
Zimmerman, M., 32