

Index

A

AACAP Workgroup on Infant and Preschool Psychiatry, 143

Aberrant Behavior Checklist (ABC), 303

Academy of Child and Adolescent Psychiatry (AACAP), 143

ADOS-2 administration, 165

ADOS-2 coding, 166–170

Adult attachment interview (AAI), 83

analysis, 85

certification and trainings, 85

childhood attachment experiences, 83

childhood mental health, 85

classification, 84

interview questions, 84

questions, 83

scoring and classification, 84, 85

SSP, 85, 86

Ages and Stages Questionnaire (ASQ)

ADOS-2, 163, 164, 171

DB-DOS, 171–173

parent-report developmental screening, 163

social-emotional development and behaviors, 163

American Academy of Pediatrics (AAP), 14

American Academy of Pediatrics Screening Recommendations, 14

Anxiety disorders, 287, 305

Assessment process, 22–24, 39

Attachment and Biobehavioral Catch-up (ABC), 341, 343

Attention deficit/hyperactivity disorder (ADHD), 277, 330, 342

amphetamine salts, 301, 302

medical explanatory model, 292

methylphenidate, 300, 301

non-stimulant medications, 302

psychopharmacological interventions on brain development, 299

psychopharmacological treatments for preschoolers, 297, 299, 302

psychopharmacology trials, 300

risk, 293

symptom tracking for preschoolers, 299

Atypical antipsychotics, 304

Atypical maternal behavior instrument for assessment and classification (AMBIANCE), 88

Autism Diagnostic Observation Schedule (ADOS), 144

Autism Diagnostic Observation Schedule-2 (ADOS-2), 163, 164, 171

Autism Spectrum Disorders (ASD), 13, 196, 231, 276

atypical antipsychotics, 304

behavioral interventions, 305

core symptoms, 303

evidence-based therapies, 305

psychotropic medications, 305

RCT, medications for preschoolers, 304

treatment for children, 303

Autobiographical memory, 199

B

Bayley Scales of Infant and Toddler Development, 229

Behavioral interventions, 310

Beliefs, 56

Berkeley Puppet Interview, 206

- Bowlby's attachment theory, 5
 Brazelton Neonatal Assessment Scale (BNAS), 190
- C**
- Caregiver and child behavior, 114
 Caregiver-child interaction, 109, 119, 288
 descriptions, 114
 emotion, situations, 107
 factors, 114
 sensitivity, 107
 Caregiver-child relationships, 5, 21, 45, 81, 109, 185, 191, 197, 203, 205, 288
 Caregiver-infant interaction
 assessment, 109, 112
 caregiver's mood, 111
 clinical questions, 117
 clinic-based assessment, 116
 components, 111
 depression and anxiety, 108, 109
 developmental level, 117
 evaluation, 114
 functioning, 110-111
 Japanese families, 113
 maternal-infant attachment, 108
 medical conditions, 132
 NCAST and PCI, 120
 psychological and personal factors, 108
 separation and reunion, 117
 traumatic events, 131
 treatment process, 115
 warmth and responsiveness, 107
 young child assessment, 133
 Caregiver management training approaches, 287
 Caregiver perceptions, young child
 AAI, 83, 84
 anxious, 79
 breed toughness, 80
 caregiver, 80
 caregiver-infant interactions, 82
 child and parenting, 78
 clinical assessments, 77
 clinical environment, 81
 culturally guided knowledge, 79
 culture, 79
 depressed parent, 82
 emotional states, 81
 environmental influences and stressors, 79
 internal working models, 78, 82
 language, 80
 memories, 78, 80
 psychotic caregiver, 82
 PTSD, 82
 reflective functioning, 77
 sensitivity, 77
 stressors, 78
 toddlers, 81
 trauma responses, 81
 Caregiver transference, 335
 Caregiver traumas, 102
 Caregiving environment and infant/young child adaptation, 269, 270
 Case conceptualization, 328
 Center for Autism and the Developing Brain (CADB), 163
 Child Behavior Checklist (CBCL), 225, 279
 Child developmental and sensorimotor assessments, 22
 Child directed interaction (CDI), 340
 Child Parent Psychotherapy (CPP), 287, 339, 343
 Child Psychiatric Rating Scale (CPRS), 303
 Child's active engagement, 198
 Child's and family's sociocultural background, 42
 Child's functioning, 185
 Child's perception, 189
 Childhood Autism Rating Scale (CARS), 303
 Circle of Security (COS), 342, 343
 Circle of Security Interview (COSI), 90
 AAI, 91
 caregiver-child interactional procedure, 91
 child's behaviors, 92
 clinical interview, 90
 intervention, 90, 91
 protocol, 90
 Clinical case formulation
 age-related considerations, 330, 331
 aggressive behaviors, 346
 behavioral formulation, 329
 behavioral observations, 348, 349
 CFI, 329
 cultural considerations, 331, 336
 cultural features, vulnerability and resilience, 350
 cultural formulations, 329
 definitions, 328
 developmental history, 348
 explicit and implicit formulations, 328
 five Ps (*see* Five P's approach)
 idiographic and nomothetic, 328
 joint treatment plan (*see* Joint treatment plan)
 medical history, 348
 mental health treatment, 349, 350

product vs. process, 329
 psychodynamic and cognitive, 329
 psychosocial stressors, 350
 social history, 347
 Clinician assisted video-feedback exposure
 sessions (CAVES), 95
 protocol, 96
 video-feedback, 95
 video review, 96
 Clinician-child relationship, 149, 150
 Clonidine, 310
 Cognitive behavioral therapies (CBT), 287,
 305, 307
 Cognitive functioning, 187
 Collaborative plan for assessment, 28
 Committee on Preschool Special Education
 (CPSE), 278
 Committee on Quality Issues (CQI), 143
 Comprehensive assessment, 315, 316
 Comprehensive joint clinical formulation,
 co-construction, 333, 334
 Comprehensive treatment plan, 336
 Conduct Disorder (CD), 274
 Conducting assessment, 27
 Countertransference, 335
 Crowell procedure, 115, 116, 133, 134
 Cultural and religious beliefs, 56
 Cultural formulation, 329
 Cultural Formulation Interview (CFI), 329
 Culture, 56, 331, 332
 Culture-based resilience, 350
 Culture-based stigma, 350, 352
 Culture-specific socialization processes, 191

D

Deferred imitation, 192
 Department of Children and Families
 (DCF), 59
 Depression, 287, 330
 Depressive Disorder
 pharmacotherapy, 312
 preschool, 312, 313
 symptoms, 312
 treatment effects, 312
 Developmental competence, 265
 Developmental Coordination Disorder, 280
 Developmentally based intervention, 71
 Diagnostic and Statistical Manual of Mental
 Health Disorders fifth edition
 (DSM-5), 228, 254
 Diagnostic Classification Revision Task
 Force, 259
 Diphenhydramine, 310

Disorder of Dysregulated Anger and
 Aggression (DDAA)
 children's aggressions, 274
 data-driven approach, 275
 early childhood, 274
 emotional and behavioral dysregulations, 274
 emotional and behavioral regulation, 274
 emotions and behaviors, 274
 family-focused therapy, 312
 impulsive dysregulation in preschoolers, 312
 irritability and aggression, 274
 mood and behavioral dysregulation, 275
 pharmacotherapy, 311
 preschoolers, 274
 problematic regulation of mood, 311
 symptom tracking, 311
 Disruptive Behavior Diagnostic Observation
 Schedule (DB-DOS), 171–173
 Diversity-Informed Tenets, 113
 Domestic violence (DV), 150
 DSM-5 Dysregulated Mood Disorder with
 Dysphoria (DMDD), 313
 Dyad's affective tone, 112
 Dyadic assessment, 133–134
 Dyadic Parent Child Interaction Coding
 System (DPICS), 280
 Dyad's behavior, 122

E

Early childhood assessment
 assessment procedures, 12
 behavioral evaluations, 13
 behaviors, 9
 bilingual homes, 12
 biology, 11
 caregiving relationship, 12
 communication skills, 17
 communicative behaviors, 9
 complication, 9
 cultural researchers, 12
 ecological and environmental factors, 12
 educational assessments, 13
 environment, 11
 evaluation, 11
 evaluator qualities, 18
 language abilities, 9
 language and communication skills, 9
 medical setting, 12
 motor movement, 10
 physical and cognitive development, 13
 scores, 14
 screening instrument, 14
 social emotional development, 11

Early childhood assessment (*cont.*)
 social-emotional needs, 9
 social-emotional skills, 10
 standardized test scores, 15
 structural recommendations, 19
 temperament and personality, 11
 test scores and performance, 15
 verbal and nonverbal communication, 11

Early Intervention (EI), 224

Early Intervention Program (EIP), 66

Ecological theory, 7

Emotional dysregulation, 312

Empathy, 193

Evidence-based interventions, 317
 ABC, 341, 343
 COS, 342, 343
 CPP, 339
 IY, 341
 PCIT, 340, 344
 TF-CBT, 340
 theraplay, 342, 344

Evidence-based psychotherapeutic
 treatments, 303

Evidence-based therapies, 285, 305

Evidence-based treatments, 297

Evidence-informed treatment approaches, 287

Explicit comprehensive clinical formulation, 331

Expressive language, 160

Eyberg Child Behavior Inventory (ECBI),
 244, 279

F

Family's support system, 52

Federally Qualified Health Center
 (FQHC), 62

Feeding situation, 124

Five P's approach
 perpetuating factors, 330, 352
 precipitating factors, 330, 352
 predisposing factors, 330, 351, 352
 presenting concerns, 351
 presenting problem, 330
 protective/positive factors, 330, 352

Food and Drug Administration (FDA), 295

Foster care system, 53

Free play situation, 124

G

Gastroesophageal reflux disorder
 (GERD), 242

Gastroschisis, 314, 315

Generalized anxiety disorder (GAD), 287, 309

H

Health-care, 53

Historical trauma, 42

History-taking
 artificial distinction, 27
 child medical and developmental history, 62
 domains, 50, 51
 family's socio-cultural and historical
 context, 41
 goal, 24
 guiding principles
 adaptation and strengths-based
 framework, 48
 affect, gestures, movement, pacing and
 vocalizing, 47
 calm and self-regulation, 49
 context of relationship, 46, 47
 cultivate self-awareness, introspection
 and reflectivity, 47
 limitations, 49
 observation, wondering and tolerance
 of discomfort, 48
 organic process and family's lead, 49
 helping systems, 59
 identifying information and description of
 presenting problem, 59
 infant, child and family, 50
 infant mental health, 24
 initial interviews, 26 (*see also* Interview
 process)
 narrative process, 23
 parenting as relationship, 50
 practitioner's stance and interpersonal
 processes, 46
 presenting concerns, 60
 relational history of family members, 42, 45
 setting up, 28
 social and family history, 60, 61
 strengths-based approach, 25
 ten core skills, 25, 26
 therapeutic alliance, 25
 trauma, consideration, 27, 28

I

Impulsivity and irritability, 314

Incredible Years (IY), 341, 343

Individuals with Disabilities Education Act
 (IDEA), 226

Infant and early childhood mental health
 (IECMH) assessment, 1, 24, 25, 51,
 253, 255, 256, 329
 assessment, 2
 attachment theory, 5

- childhood cognitive processes, 4
- definition, 1
- disciplines, 5
- dynamic process, 5
- field, 3
- historical context, 2
- multidisciplinary approach, 6
- teachers and parents, 4
- WAIMH, 2
- ZTT, 3
- Infant and Toddler Mental Status Exam (ITMSE), 189
 - affect and mood, 161
 - appearance, 159, 176
 - behavioral observations, 176
 - caregiver context, 159
 - child medical and developmental history, 175, 176
 - cognition, 162
 - formulation and plan, 179–181
 - mood/affect, 179
 - motor, 160
 - motor skills, 177
 - observational skills, 159
 - play, 161, 162, 177
 - presenting concerns, 174
 - rating scale information, 179
 - reaction to situation, 159, 176
 - relatedness, 162
 - relatedness (to grandmother), 178
 - relatedness (to mother), 178
 - self-regulation, 159, 160, 177
 - social and family history, 174, 175
 - speech, 177
 - speech and language
 - expressive language, 160
 - phonologic speech, 160
 - pragmatic language, 161
 - receptive language, 160
 - thought, 161
- Infant-child-family profile, 51
- Infant mental health, 288
- Infant parent psychotherapy, 287
- Infant–Toddler Social and Emotional Assessment (ITSEA), 224
- Inner world of young child
 - age grouping, 186
 - ages 2 to 5 years
 - autobiographical memory, 199
 - caregivers, 201
 - child’s ability, 202
 - child’s daily life and home environment, 202
 - child’s participation, 202
 - child’s play, 201
 - children’s guidance, 204
 - encouragement and scaffolding, 204
 - expressions, 203
 - expressive modalities, 199
 - gender and cultural differences, 200
 - gender identity, 199
 - internal representations, 198
 - interviewing, 203
 - language fluency, 198
 - metaphorical and fantasy-based, 200
 - modalities, 202
 - physical reenactments, 203
 - play materials, 205
 - playhouses, 206
 - secure base, 200
 - sensory-motor and emotional pathways, 198
 - sexual abuse, 206
 - structured play, 205
 - symbolic play, 199, 200
 - symptoms and environmental triggers, 201
 - TOM, 199
 - unguided play, 204
 - verbal communication, 198
- ages birth to 24 months
 - aggression, 197
 - attachment quality, 192
 - behavior and emotional responses, 190
 - caregivers, 190, 191
 - child’s internal schemata, 194
 - cognitive and memory standpoint, 192
 - cognitive and motor development, 195
 - cognitive and social capacities, 190
 - compelling structured paradigm, 196
 - crawling, 196
 - custom’s inspection, 191–192
 - developmental and clinical research, 190
 - developmental delay, 196
 - developmental testing, 193
 - distress, 194, 195
 - empathy, 193
 - facial expressions, 190
 - feeding materials, 197
 - growing capacity, 193
 - infants, 191, 196
 - mother and child’s play, 195
 - physiological regulation, 191
 - reciprocal play, 195
 - secure behaviors, 196
 - sensory-motor based patterning, 191
 - separation/reunion sequences, 196, 197
 - sequential approach, 195

- Inner world of young child (*cont.*)
- small structuring and scaffolding, 197
 - social relatedness, 190
 - traumas, 193
 - biological/temperamental factors, 188
 - caregiver–infant interactions, 186
 - caregivers, 188
 - child sessions, 210, 211
 - child’s internal representations, 185
 - child’s play, 208
 - child’s play skills, 190
 - clinician’s conceptualization, 186
 - clinicians, 187
 - cognitive functioning, 187
 - data gathering, 187
 - developmental history, 209
 - diagnosis and treatment, 211, 212
 - evaluation process, 187
 - identifying information and description,
 - presenting concerns, 207
 - infantile amnesia, 185
 - infants, 188
 - play and expressive communication, 188
 - presenting concerns and background
 - information, 209
 - psychiatric assessment process, 185, 186
 - reciprocity and social relationships, 189
 - relational scripts, 185
 - self-expression, 189
 - self-reflection, 187
 - sessions, 189
 - time frames, 186
 - toddlers, 188
- Insightfulness Assessment (IA), 92
- assessment, 92
 - child’s behavior, 92
 - classifications, 93
- Intergenerational transmission, 42, 45, 95
- Internal working models, 78, 83
- Interview process
- adequacy of parent, 24
 - caregiver–family, 24–26
 - case studies (Ana L and Paquito)
 - access to services, 66
 - behavioral observations, 62, 63
 - collateral agencies, involvement of, 66
 - concrete needs, housing, safety, food,
 - financial needs, 67
 - current medical/dental and mental
 - health, 66, 67
 - current members and household, 64
 - diagnosis and formulation/plan, 70, 71
 - educational skills and employment
 - history, 65
 - family, cultural and religious values
 - and socio-cultural context, 68
 - immigration status and concerns, 67
 - interviewer’s impressions based on
 - history, 70
 - legal involvement and concerns, 67
 - parent’s internal representation
 - of child, 68
 - pregnancy, birth and early life, 64
 - protective and risk factors, 67
 - questions and responses, 68, 69
 - significant caregivers, moves,
 - separations and trauma, 65
 - developmental status of child, 24
 - domains of inquiry, 51
 - access to services, 53
 - educational and social service
 - agencies, 53
 - educational, skill and employment
 - history, 52
 - family, cultural and religious beliefs, 56
 - health/dental/mental health care, 53, 54
 - immigration status/concerns, 54
 - legal involvement and concerns, 54
 - members of family and family’s
 - support system, 52
 - parent’s internal representation of
 - infant/child, 57
 - pregnancy, birth and/infant/child
 - information, 51, 52
 - protective and risk factors, 55, 56
 - safety, housing, food, clothing and
 - financial concerns, 55
 - significant caregivers, losses and
 - traumas, 52
 - WMCI questions, 57
 - practitioner reflects, 58
 - practitioner’s stance and interpersonal
 - processes, 46
 - presenting problem, 39, 40
 - proper goodbye, 58
 - question-answer process, 51
 - reasons for referral, 39
 - reflections of interviewer’s experience, 58, 59
 - setting up, 28, 45, 46
 - therapeutic frame, setting, 38, 39
 - validating experiences and meaning-
 - making, 49, 50
- J**
- Joint clinical case formulation, construction
- caregiver’s emotional responses, 335
 - caregiver’s perspective, 333

- clinicians, 334
 - comprehensive joint clinical formulation, 333, 334
 - conceptualization to treatment plan, 333
 - consistent and routine invitation, caregiver, 336
 - depression/substance abuse, 335
 - development, 333
 - empathic and inclusive approach, 333
 - failure to construct, 334
 - infant and early childhood mental health, 332
 - nonjudgmental language, 353
 - patience, thoughtfulness, time and commitment, 334
 - provider's management, 335
 - psychoeducation, 334
 - soft start up, 334–336
 - subsequent soft formulation, 335
 - Joint treatment plan
 - assessment and mental health service provision, 336
 - buy-in, 337
 - caregiver strengths and challenges, 337
 - chief complaint, 336, 337
 - clinical case formulation and cultural considerations, 336
 - clinician and family, 336
 - components, 338
 - construction, 338
 - development, 337
 - infant and early childhood mental health clinicians, 338
 - interventions, 337, 338
 - multitiered behavior, 337
 - production, 338
 - SMART goals, 338
- L**
- Level of Caregiving Environment and Infant/Young Child Adaptation, 269
 - Levels of Adaptive Functioning–Caregiving Dimension, 268
- M**
- MacArthur Story Stem Battery (MSSB), 205
 - McMaster Family Assessment Device, 269
 - Medicaid services, 53
 - Medications
 - anxiety disorders, 306, 307
 - ASD, 305
 - caregivers of preschoolers, 290
 - case studies, preschoolers with anxiety, 306
 - decisions and informed consent, 288
 - mental health concerns, 291
 - non-stimulant, 301, 302
 - OTC, 309
 - perceptions, 290
 - psychotropic, 305
 - PTSD, 308
 - rapid development, 293
 - RCT, 295
 - treatment decisions, 291
 - Melatonin, 310
 - Mental health disorders
 - approaches and assessment, 254
 - caseness, 255
 - classification, 255
 - comprehensive assessment, 266
 - data collection, 266
 - DC
 - assessing symptoms, 260
 - classification system, 258
 - comprehensive assessment, 260
 - cultural context, 261
 - cultural domains, 259
 - developmental competence, 265
 - diagnostic assessments, 260
 - diagnostic classification system, 259
 - infant and early childhood psychopathology, 258
 - multiaxial approach, 259
 - physical conditions & considerations, 263
 - psychosocial stressors, 264, 265
 - reflective practice, 260
 - risks and benefits, 259
 - self-awareness, 260
 - self-reflection, 261
 - Task Force recommends, 259
 - developmental history, 277
 - diagnosing mental health and developmental disorders
 - child's symptoms, 270
 - clinical disorders, 271–273
 - contextual factors, 274
 - cultural and contextual dimensions, 271
 - DC:0-5, 273
 - DDAA, 274
 - diagnostic algorithms, 274
 - Relationship-Specific Disorder, 271
 - distractibility and self-regulation, 277
 - education history, 278
 - emotional well-being, 254
 - family and social history, 278
 - infancy and early childhood, 255
 - infant-family system, 253

- Mental health disorders (*cont.*)
- multiaxial approach, 255
 - nosology, 255
 - relational context
 - assessment tools, 268
 - caregiving characteristics, 267
 - caregiving environment, 266, 269
 - caregiving–infant /young child relationship, 268
 - child–caregiver relationship, 269
 - child-caregiving environment, 267
 - child’s social and emotional development, 270
 - infant/young child, 267
 - relational context, 267
 - stability, predictability and emotional regulation, 269
 - school observation, 279
 - self-regulation and aggression, 280
 - symptom checklists, 279, 280
- Mental health practitioners, 50
- Mental health treatment, 349–351
- Mental representations, 208
- Methylphenidate, 300, 301
- Minnesota Parent Rating Scale, 179
- Monotherapy, 289
- Mood problems, 313
- Motherese/infant-directed speech, 194
- Motor movements, 10
- Mullen Scales of Early Learning, 229
- Multidimensional Assessment Profile of Disruptive Behavior, 224
- N**
- National Institute of Mental Health, 255
- National Scientific Council on the Developing Child, 8
- Negative predictive value (NPV), 235
- Nonproprietary preschool measures, 289
- Non-standardized assessments, 15
- Non-standardized/authentic assessment, 15
- Nursing Child Assessment Satellite Training (NCAST), 115, 119
- assessment process, 125
 - clinical practice settings, 120
 - clinicians, 121
 - health care professional flexibility, 120
 - infant/child’s experience, 122
 - interaction procedure, 123
 - national trainer, 120
 - parent–child relationships, 122, 123 and PCI, 120
 - protocols, 121
 - training package, 123
- O**
- Observation tool, 121
- Observational assessment, 109, 115, 137
- Observational methods, 113
- Obsessive Compulsive Disorder (OCD), 277, 278, 280, 308, 309
- Obstructive Sleep Apnea (OSA), 309
- Oppositional Defiant Disorder (ODD), 274, 277
- P**
- Parent Child Interaction Therapy (PCIT), 242, 287, 299, 312
- Parent Directed Interaction (PDI), 340
- Parent Management Training (PMT) approaches, 311, 316
- Parent Stress Index, 179
- Parental reflective functioning, 50
- Parental trauma history, 46
- Parent–Child Early Relationship Assessment, 114
- Parent-child interaction assessment, 134
- Parent-Child Interaction Therapy (PCIT), 149, 340, 344
- Parent-child relationship, 41, 98
- Patient Centered Medical Home (PC-MH), 54
- Phonologic speech, 160
- Piaget’s study, 4
- Play therapy, 288, 308
- Play-based assessment, 187, 207
- Play-based scaffolding techniques, 206
- Positive predictive value (PPV), 235
- Posttraumatic Stress Disorder (PTSD), 82, 245, 276
- CBT, 287
- Child Parent Psychotherapy and Infant Parent Psychotherapy, 287
- play therapy, 308
 - preschool, 307, 308
- Practice parameters
- AACAP, 145
 - caregivers, 145
 - child participation, 146
 - child room environment, 146
 - clinician’s roles
 - clinician-child relationship, 149, 150
 - documenting observations, 152
 - interacting with caregivers, 151, 152
 - play technique and establishing rapport, 148, 149
 - description, 143
 - developmental milestone table, 156–157
 - disruptive behavior, 145, 147
 - evidence base, 144
 - factors, 144
 - ideal minimalist office set up, 147

- ITMSE (*see* Infant Toddler Mental Status Exam (ITMSE))
- mental health assessment, 146
- miscellaneous toys for developmental assessment
- balls, 158
 - behavioral disorders, 155
 - books, 158
 - garage with cars and keys, 155, 158
 - paper and washable thick markers, 158
 - puzzles, 158
- observational play, 145
- observational skills, 144
- parent interview room, 147
- semi-structured tools, 144
- structured and semi-structured assessment tools, 158
- toys and materials
- aggressive toys, 154
 - categories, 153
 - clinical setting, 153
 - ethnic and cultural factors, 153
 - expressive toys, 154
 - family/nurturing toys, 153, 154
 - pretend/fantasy toys, 154
 - rules, room environment, 153
 - scary toys, 154
 - sensory toys, 155
- video-conferencing applications, 147
- young child assessment, 144
- young children's behavior, 145
- Pragmatic language, 161
- Praise, reflection, imitation, description, and enthusiasm (P.R.I.D.E.), 149
- Preschool ADHD Treatment Study (PATS), 294
- Preschool Anxiety Rating Scale (PARS), 306
- Preschool psychopharmacology
- child medical and developmental history, 314, 315
 - diagnosis, formulation and plan, 315–317
 - principles, 317
 - treatment (*see* Psychopharmacologic treatment, preschool children)
- Preschoolers, 144, 151, 172, 181
- Prescribing practices
- caregiving status and demographics, 297, 298
 - child factors, 297, 298
 - insurance, 297, 298
 - trends, 296
 - workforce, 296
- Primary care providers (PCPs), 224
- Probable experience scales, 84
- Proper goodbye, 58
- Protective factors, 56
- Psychoanalytic tradition, 3
- Psychodynamic approaches, 329
- Psychoeducation, 334
- Psychopharmacologic intervention, 344, 345
- Psychopharmacologic treatment, preschool children
- ADHD over medication treatments, 290
 - anxiety disorders, 305
 - ASD (*see* Autism Spectrum Disorders (ASD))
 - caregiver–child relationship, 287
 - caregivers, 290
 - clinical humility, 292
 - clinician beliefs, 291
 - culture, race, and community, 292
 - DDAA (*see* Disorder of Dysregulated Anger and Aggression (DDAA))
 - infant mental health, 288
 - interventions, 287
 - medication, 288–290
 - OCD, 308, 309
 - play therapy, 288
 - practice guidelines, 286
 - psychotherapy, 287
 - PTSD, 307, 308
 - rapid development, 293
 - reassessment, diagnosis, 288
 - recommendations and decisions, 291
 - regulatory factors, 295
 - research base guiding clinical decisions, 295
 - sleep (*see* Sleep problems in preschoolers)
 - workforce, 296
- Psychosocial stressors, 264, 265
- Psychotherapeutic approaches, 285
- Psychotherapeutic intervention, 316
- Psychotherapy, 86, 287, 288, 307
- Psychotropic medications, 305
- Puppet-based interviewing techniques, 206
- R**
- Receptive language, 160
- Reciprocal engagement, 189
- Reflective functioning, 50, 77
- Relational scripts, 185, 191
- Relationally-based developmental guidance, 71
- Relationship-based assessments, 144
- Relationship-Specific Disorder, 267
- Research Domain Criteria (RDoC), 255
- S**
- Scoring system, 87
- Screen for Anxiety and Related Emotional Disorders (SCARED), 306
- Selective mutism, 287

- Selective serotonin reuptake inhibitors (SSRI's), 305
 - Self-actualization, 55
 - Self-awareness, 47
 - Sensory Over Responsivity Disorder, 276
 - Separation/reunion situation, 124
 - Sleep problems in preschoolers
 - behavioral interventions, 310
 - clonidine, 310
 - diagnosis, 309
 - diphenhydramine, 310
 - melatonin, 310
 - OTC medications, 309
 - tracking, 309
 - SMART goals approach, 338
 - Social and emotional skills, 10
 - Social-emotional and behavioral (SEB)
 - adult-report rating scales, 217
 - assessment goals, 223
 - behavioral observations, 243
 - caregiver, 218
 - child medical and developmental history, 242–243
 - clinical assessment, 217
 - clinical rating scale
 - comprehensive SEB checklists, 239
 - problem-specific rating scales, 240
 - screeners, 238
 - comprehensive assessments, 227
 - contextual and cultural factors
 - influence, 221
 - interpersonal relationships, 221
 - language and literacy, 221
 - multiple settings, 222
 - neighborhood and community, 222
 - stressors, 222
 - values and norms, 221, 222
 - domains of interest
 - assessing competencies, 228
 - behavioral functioning, 227
 - caregiver concerns, 229
 - cognitive and language delays, 229
 - disruptive behavior problems, 227, 228
 - early childhood, 227
 - impairment, 230
 - trauma and stress, 228
 - formulation and plan, 244, 245
 - history, presenting concerns, 241, 242
 - identifying information and description, 241
 - interpreting rating scales and sharing
 - results, families, 238–241
 - limitations, 219
 - psychometric properties
 - content validity, 233
 - convergent validity, 233
 - cultural adaptation, 237
 - discriminant/between groups validity, 233
 - face validity, 233
 - factor structure, 233
 - influence instrument adoption, 230
 - internal consistency, 231
 - longitudinal developmental context, 234
 - parent–caregiver reports, 230
 - predictive validity, 234
 - sensitivity/specificity, 234
 - standardization, 236
 - statistics and rules, 232
 - test–retest reliability, 231
 - types of, 232
 - rating scale information, 243, 244
 - rating scales, 223
 - screening, 225
 - screening tools vs. comprehensive
 - assessment tools, 223
 - select informants
 - caregivers, 219
 - childcare providers, 220
 - family members and child-care providers, 219
 - non-parental family members, 220
 - parents/guardians, 220
 - social and family history, 242
 - strengths, 218
 - Social phobia, 287
 - Socio-cultural contexts, 41, 42
 - Soft start up, 334–336, 353
 - Special education services, 53
 - Spence Preschool Anxiety Scale (SCAS), 280
 - Standardized evaluation tools, 15
 - Still-Face Paradigm (SFP), 125
 - infant behaviors, 126
 - research tool, 125
 - Story stem technique, 205
 - Strange Situation, 192
 - Strange Situation Procedure (SSP), 83, 85, 126
 - Stranger anxiety, 192
 - Structured clinical assessment approach, 109
 - Structured task situation, 124
 - Substance Abuse and Mental Health Services Administration's (SAMHSA), 339
 - Systemic Lupus Erythematosus (SLE), 175
- T**
- Theory of Mind (TOM), 199
 - Theraplay, 342, 344
 - Trauma, 330
 - historical, 41, 42, 51
 - history-taking, 27, 28
 - intergenerational, 51

- intergenerational transmission, 42
 - political, 41
 - relational, 70
- Trauma-Focused Cognitive Behavioral Therapy (TF-CBT), 294, 340, 344
- Treatment phases, 27
- Treatment planning and intervention, 22
- Treatment plans, 336

U

- Unconscious projections, 80

V

- Video Replay Interview, 125

W

- Working Model of the Child Interview (WMCI), 57, 99, 101, 315

- AAI, 87

- AMBIANCE, 88

- classifications, 88, 89

- disengaged representations, 88

- empirical research, 89

- examination, 87

- infant attachment classification, 90

- knowledge, 86

- mediation analysis, 90

- questions, 87, 88

- treatment planning, 87

- World Association of Infant Mental Health (WAIMH), 2, 270