

Index

A

Alagille syndrome, 340
Anorectal malformations, 112

B

Berliner Gesellschaft für Anthropologie, 5
Bidloo, Govert, 6
Bone morphogenetic protein-2 (BMP-2), 23
Brunner, Johann Conrad, 8, 9

C

Carcinoembryonic antigen (CEA), 176–177
Caudal cell mass (CCM), 25, 26
Center for Disease Control (CDC), 67
Clinical presentation, OSD
 cutaneous stigmata
 aplasia cutis congenita, 102
 asymmetric lesions, 100
 atretic meningocele, 102, 106
 caudal appendages, 102, 105, 107
 coccygeal pit, 106, 109
 cutaneous finding, 100, 103
 DST, 105, 106, 108
 flat capillary hemangiomas, 100
 focal hirsutism, 100, 101
 focal hypertrichosis, 100
 hyperpigmented nevi, 102
 hypopigmented maculae, 102
 infantile hemangiomas, 100, 105
 lipomelomeningoceles, 100
 lipomyelomeningoceles, 100
 medical evaluation and investigation,
 100
 neuroenteric cysts, 100
 silky down, 100
 terminal myelocystoceles, 100
 valsava maneuvers, 100

neurologic symptoms, 109–110
orthopedic deformities, 110–112
urologic dysfunction, 110

Closed spinal dysraphic conditions
 animal models and human genetics
 Currarino syndrome, 339, 340
 junctional neurulation defects, 335
 neural tube closure, 333, 334
 open NTDs, 333, 334
 sacro-caudal agenesis, 338, 339
 sacrococcygeal teratoma, 336, 338
 spinal cord tethering, 334
 spinal lipoma, 335–337
 split cord malformations, 339
 vertebral defects, 340, 341
 backwards extrapolation, 325
 congenital disease
 animal models, 327, 328
 patient-oriented genomic approaches,
 326, 327
 embryonic development
 convergent extension, 330
 gastrulation, 328
 NMPs, 329, 330
 notochord, 332
 primary and secondary neurulation,
 330, 331
 vertebral development, 332
Club foot, 110
Coccygeal ligament, 88
Complex lipomyelomeningocele, 274, 275
Congenital talipes equinovarus (CTEV),
 110, 112
Conspicuous abnormalities, 4
Constructive interference in steady state
 (CISS) imaging, 197, 244
Conus lipoma
 natural history, 317–318
 surgical outcome, 319, 320

- Conus medullaris, 291
 Convergent extension (CE), 22, 23, 330
 Cruveilhier, Jean, 3
 Currarino syndrome, 294, 339, 340
 Currarino triad, 112
 Cutaneous hemangioma, 212
 Cutaneous stigmata, 212, 213
 - aplasia cutis, 77, 78, 102
 - asymmetric lesions, 100
 - atretic meningocele, 70, 73, 102, 106
 - capillary hemangioma, 70, 73–75
 - caudal appendages, 102, 105, 107
 - coccygeal pit, 106, 109
 - complications, 305, 306
 - correlation analysis, 70
 - cutaneous finding, 100, 103
 - dermal sinuses, 76, 77
 - dorsal appendage, 70
 - DST, 105, 106, 108
 - flat capillary hemangiomas, 100
 - focal hirsutism, 70–72, 76, 100, 101
 - focal hypertrichosis, 100
 - hemangiomas, 71, 75
 - human tails, 70, 77
 - hyperpigmented nevi, 102
 - hypertrichosis, 70, 72, 76
 - hypopigmented maculae, 102
 - infantile hemangiomas, 100, 105
 - lipomelomeningoceles, 100
 - lipomyelomeningoceles, 100
 - medical evaluation and investigation, 100
 - neuroenteric cysts, 100
 - silky down, 100
 - skins lesions, 77
 - small dermal sinus, 70, 73
 - spinal dysraphism, 77
 - subcutaneous lipoma, 70, 76
 - terminal myelocystoceles, 100
 - Valsalva maneuvers, 100
 - vascular nevi, 71, 75
- Cytokeratin, 177
- D**
- d'Angers, Ollivier, 11
- Dermal sinus tracts (DSTs), 62, 105, 106, 108, 229, 230
 - clinical presentation
 - infection, 169, 170
 - neurosurgical evaluation, 167
 - superficial abnormalities, 167, 168
 - conservative treatment, 298
 - cutaneous stigmata, 297
 - dermal sinus like stalk, 170
 - dermoid/epidermoid lesions, 250, 251
 - diagnostic study, 170
 - disordered embryogenesis, 166, 167
 - encephaloceles, 298
 - epidemiology, 165
 - Escherichia coli*, 297
 - etiology, 296
 - failure of disjunction, 166
 - Hensen's node, 165, 166
 - infection, 106–108
 - LDM, 250, 252
 - management, 299
 - meningocele manqué, 252
 - MRI, 298
 - natural history, 314
 - pathophysiology, 166, 167
 - primary neurulation, 166
 - squamous epithelium, 166
 - Staphylococcus aureus*, 297
 - superficial dermal sinus ostium, 166
 - surgical outcomes, 314
 - symptoms, 297
 - terminal syringomyelia, 299
 - treatment
 - outcomes, 172
 - postoperative care, 171, 172
 - surgical technique, 168, 171
 - timing of surgery, 170, 171
 - ultrasonography, 297
- Dermal-sinus-like stalk (DSLS), 229
- Diastematomyelia, 11–13, 63, 209, 292
- Diffusion tensor imaging (DTI), 244, 259
- Diffusion-weighted imaging (DWI), 184, 259
- Diplomyelia, 12, 40, 63, 210, 213, 258, 292
- Dorsolateral hinge points (DLHP), 20
- Dysraphia
 - clinical findings, 13
 - clinical implications, 13
 - high morbidity and mortality, 13
 - management and outcomes, 13
 - meningocele manqué, 13, 14
 - surgical intervention, 13
- E**
- E-cadherin, 18
- Embryology
 - blastocystic cavity
 - amniotic cavity, 17, 18
 - primitive streak, 18, 19
 - trophoblast, 17, 18
 - umbilical vesicle, 17
 - conus medullaris, 30, 31, 33

- dysraphic malformations (*see* Neural tube defects (NTD))
- gastrulation, 19
- junctional neurulation
- mapping studies, 29
 - NSB, 29, 30
 - PCP pathway, 29
 - SSEP, 29, 30
- notochordal process, 19–21
- primary neurulation
- anterior neuropore, 22
 - caudal brainstem/cranial spinal cord, 20
 - caudal neuropore, 22
 - convergent extension, 22, 23
 - DLHP, 20
 - elevation and convergence, 23, 24
 - filopodia, 24, 25
 - floor plate reaction, 23, 24
 - homeobox gene *Pax3*, 25
 - lamellipodia, 25
 - neural crest cells, 20
 - neural tube defects, 22
 - occipito-parietal region, 22
 - pseudostratified columnar epithelium, 20, 22
 - Schwann cells, 20
 - Shh* expression, 23
 - Wnt/ β -catenin PCP pathway, 25
- secondary neurulation
- in animal models, 28
 - avian morphogenesis, 28
 - CCM, 25, 26
 - mammalian neurulation, 26, 28
 - N-CAM, 28
 - overlap zone, 26
 - secondary neural tube, 26
 - Wnt/ β -catenin pathway, 28
- Epithelial membrane antigen (EMA), 176
- Excalation, 20, 21
- Exercitationum Anatomico-Chirurgicarum*, 6
- F**
- Fast Imaging Employing Steady-state Acquisition (FIESTA), 244
- Fatty filum terminale, 291, 292
- Fetal MRI, 260
- Fibroblast Growth Factor 8 (FGF8), 18
- Filum lipomas, 308
- natural history, 316
 - surgical outcome, 316–317
- Filum terminale, 212
- conus medullaris, 86
 - duplicated filum terminale, 92, 93
- embryonic development
- congenital malformations, 85
 - dedifferentiation, 85, 86
 - posterior neuropore, 85
 - primary neurulation, 84, 85
 - sacral nerve roots, 86
 - secondary neurulation, 85
 - ventriculus terminalis, 85
 - vertebral canal, 85
 - vertebral column, 85, 86
- fatty infiltration, 90–92
- FTE, 88
- FTI, 82, 84, 86–88
- history
- abnormal stretching, 83
 - antiquity, 81–83
 - De Humani Corporis Fabrica*, 81, 84
 - intermittent incontinence, 82
 - sacral lipoma, 83
 - spinal cord dysfunction, 83
 - stretch-induced symptomatology, 84
 - tethering-induced spinal cord lesions, 82
- myxopapillary ependymoma, 93, 94
- premature and term neonates, 86
- radiology, 95
- SCM, 92
- surgery, 95, 96
- TCS, 90, 91
- variations, 89
- ventriculus terminalis, 87, 94, 95
- Filum terminale externum (FTE), 88
- Filum terminale internum (FTI), 82, 84, 86–88
- Filum terminale syndrome/cord-traction syndrome, 10
- Fluid-attenuated inversion recovery (FLAIR) sequences, 184
- Focal hypertrichosis, 212
- G**
- Garceau, 10
- General anesthesia, 202, 203
- Generalized auto-calibrating partially parallel acquisition (GRAPPA), 259
- Glial fibrillary acidic protein (GFAP), 177
- H**
- Hairy patch, 100
- Hemimyelia, 284, 285
- Hemimyelomeningoceles, 282, 284
- Hemivertebrae, 110
- Hensen's node, 18
- Hydrocephalus, 9

I

Immunohistochemistry (IAC), 180
 International Mouse Phenotyping Consortium (IMPC), 327
 Intradural extramedullary (IDEM) compartment, 175–176
 Intramedullary cyst, 9
 Intra-operative physiological monitoring (IOPM), 201

J

Johnson, Athol, 9
 Junctional neurulation disorders, 335

K

Klippel-Feil syndrome, 228

L

Lifelong Specialty Care, 308, 309
 Limited Dorsal Myeloschisis (LDM), 250, 252
 Lipomas
 conus lipomas
 natural history, 317–318
 surgical outcome, 319, 320
 filum lipomas
 natural history, 316
 surgical outcome, 316–317
 symptomatic and asymptomatic lesions, 311
 Lipomyelomeningoceles (LMCs), 7, 256
 benign teratomas, 194
 Chiari malformation, 294
 classification schemes, 194
 classification systems
 Arai scheme, 197
 caudal LMC, 197
 chaotic lipomas, 198
 dorsal LMC, 197, 198
 lipogenic progenitor cells, 198
 subcutaneous cystic meningeal sac, 197
 transitional LMC, 197, 198
 clinical deficit, 194
 clinical presentation, 195, 196
 conus medullaris, 194
 Currarino syndrome, 294
 endoscopic resection, 205
 fatty-fibrous masses, 194
 fatty-fibrous tissue, 194
 imaging, 196, 197
 incidence, 293
 lipomatous OSDs, 255

longitudinal stress, 195
 micro-trauma, 195
 motor and sensory symptoms, 294
 MRI, 254
 musculoskeletal and neural spinal elements, 194–195
 neural placode reconstruction, 203, 204
 non-lipomatous OSDs, 255
 PELVIS syndrome, 294
 postovulatory day, 293
 primary neurulation, 293
 progressive incontinence, 195
 sensorimotor dysfunction, 195
 sirenomelia, 295
 subtraction osteotomy, 205
 surgical approaches
 asymptomatic patients, 200
 children harboring, 200
 classification, 200
 conventional classic principles, 200
 conventional resection, 202, 203
 disease progression, 199
 intraspinal pathology, 198
 microsurgical dissection, 200
 natural history, 199, 200
 neurological decline, 198, 199
 neurological impairment, 199
 normal neurologic exams, 199
 operative procedure, 200
 Pang's techniques, 201
 partial resection, 200
 pathophysiology, 198
 prophylactic untethering, 199
 radical resections, 201
 sac ratios, 201
 sphincteric and lower extremity function, 199
 statistical significance, 201
 surgical group, 200
 surgical team, 199
 surgical treatment, 200
 surgical untethering, 199
 visualization, 198
 wound-related complication rates, 200
 surgical outcomes and complications, 204, 205
 T1-weighted and T2-weighted images, 254, 255
 TCS, 294
 TSC, 194
 X-rays, 253
 Long tethering tracts (LTTs), 229
 Lumbosacral dysraphism, 9
 Lumbosacral hypertrichosis, 3

M

Magnetic resonance spectroscopy (MRS), 184

Meningocele manqué (MM), 13, 14

atretic meningoceles, 219

classifications, 228–230

clinical presentations, 226, 227

congenital dorsal bands, 219

conus medullaris, 228

dermoid cysts, 228

DST, 252

embryology, 224–226

imaging, 223, 227

Klippel-Feil syndrome, 228

MMC, 285

vs. SCM, 228

syringohydromyelia, 228

tethering tracts, 220–224

treatment, 230, 231

Mermaid syndrome, 295

Mesenchyme-to-epithelium transition

(MET), 334

Mesoderm, 5

Miller-Dieker syndrome, 291

Morgagni, Giovanni, 9

Multivariate regression model, 146

Muscatello, Giuseppe, 9

Myelocystocele, 239

Myelomeningocele (MMC), 9

clinical outcome, 287

clinical presentation, 283, 284

hemimyelomeningoceles, 282

incidence, 282

management, 285–287

Pang's unified theory, 281

vs. SCM, 282, 283

spina bifida, 281

Myosin light chain (MLC), 23

N

National Spina Bifida Patient Registry

(NSBPR), 66

Natural history

asymptomatic patients, 60

characterization and classification, 67

DSTs, 62

long-term effectiveness, 60

neurological disability, 60

outcome parameters, 67

principles, 61

prophylactic surgery, 60

retrospective single institution centers, 66

SCMs, 62, 63

spinal cord lipomas

clinical decision making, 64

conus medullaris, 63, 64

dural sleeve reconstruction, 65

fat and fibrous masses, 63

fatty infiltration, 63

follow-up, 65

iatrogenic neurological injury, 64

inherent operative risks, 65

long-term protection, 65

minimal operative morbidity, 64

neurological decline, 64

older patients, 64

operative and observation treatment

arms, 64

operatively-treated cohort, 64

pre-operative progressive decline, 64

radical lipoma resection, 65

radical resection and reconstruction, 64

signs and symptoms, 64, 65

technical demands, 65

technical performance, 64

surgical decision making, 65, 66

symptomatic patients, 60

Neural crest (NC), 330

Neural tube defects (NTD)

amniotic tissue, 35

associated cutaneous anomalies, 33

caudal agenesis

associated limb anomalies, 49

associated visceral malformations,

49, 50

caudal urogenital and gastrointestinal

systems, 50

Currarino's triad, 50, 51

dysplastic glial nodule, 48, 49

early embryogenesis, 50

glycoproteins, 50

vertebral anomalies, 48, 49

caudal neuropore, 34, 35

combined (anterior and posterior) spina

bifida, 41, 43

complex dysraphic malformations, 41,

44–46

cutaneous ectodermal elements, 33

dermal sinus-like tracts, 33

diastematomyelia, 40

diplomelia, 40

dural coverings, 40

dural sheaths, 40

dystrophic median nerve roots, 40, 42

endodermal-ectodermal adhesion, 42

gastrulation, 44, 45

hemicords, 40

Hensen's node midsagittally, 45

- Neural tube defects (NTD) (*cont.*)
- Hensen's node regression, 44
 - limited dorsal myeloschisis, 33, 35
 - midline fibrous bands, 40, 41
 - multiple somitic abnormalities, 45–46
 - neurenteric canal, 44
 - notochordal duplication, 44
 - osseous/fibrocartilagenous spurs, 40
 - pathological tracts, 35, 36
 - retained medullary cord, 51
 - sacro-coccygeal sinuses, 35
 - SCM, 38, 40
 - separate filum terminale, 34
 - spina bifida, 38
 - spinal lipomas/lipomyelomeningoceles
 - caudal repressive Wnt signals, 38
 - chaotic lipoma, 37
 - Chapman classification, 36
 - conus and filum terminale, 38
 - dermal sinus tract, 37–39
 - extradural tissues, 36
 - fatty spinal cord malformations, 35, 37
 - rostral malformations, 38
 - Type I, 37
 - Type II, 37
 - Type III, 37
 - Type IV, 37
 - terminal myelocystoceles, 46–48
 - thoracolumbar junction, 46
 - vertebral anomalies, 44
- Neurenteric canal, 20
- Neurenteric cysts (NECs), 257, 258, 295, 296
- clinical presentation
 - acute onset of symptoms, 177
 - anomalies, 177
 - benign lesions, 177
 - bimodal age distribution, 177
 - contrast T1 sagittal MR, 177, 179, 181
 - intradural extramedullary neurenteric cyst, 177, 181, 182
 - intra-operative images, 177, 180, 181
 - location, 177
 - lower clivus, 177
 - malignant transformation, 177
 - nonspecific symptoms, 177
 - post-operative T2 sagittal, 177, 181
 - radiological findings, 177, 178, 181
 - T1 sagittal, 177, 181, 183
 - T2 coronal MR images, 177, 181, 183
 - T2 sagittal MR image, 177, 181, 183
 - ventrally-placed foramen magnum, 177
 - diagnostic imaging
 - CT imaging, 181
 - lobulated cystic lesion, 181
 - NAA, 184, 185
 - differential diagnosis, 185
 - early embryogenesis, 176, 177
 - epidemiology, 175, 176
 - histopathological classification, 178, 179, 184
 - history, 175
 - IHC, 180
 - intracranial location, 176
 - surgical management, 185–187
 - surgical outcome and recurrence, 187
- Neurocutaneous melanosis (NCM), 292
- Neuroectoderm, 5
- Neurologic deterioration
- in adult population, 109
 - deep tendon reflexes, 109
 - extension movements, 109
 - hydrocephalus, 110
 - Lhermitte sign, 110
 - lower motor neuron symptoms, 109
 - nonhealing ulcers, 109
 - pathologic fixation, 109
 - in pediatric population, 109
 - repetitive flexion, 109
 - scoliotic deformity, 110
 - sensory loss, 109
 - upper motor neuron symptoms, 109
- Neuro-mesodermal progenitors (NMPs), 329, 330
- Neuro-orthopedic syndrome, 63, 110, 315
- Node-streak border (NSB), 29, 30
- Notochord, 332
- Notochordal canal, 20
- O**
- Occult tethered cord syndrome (OTCS), 274
- conservatively-managed patients, 159
 - cutaneous/anatomical hallmarks, 156
 - data-driven changes, 159
 - diagnostic tests, 160
 - evidence-based medicine, 161
 - filum sectioning, 159
 - general population, 156, 157
 - non-thick filum terminale, 154
 - normally-positioned conus, 154–156
 - pediatric surgical series, 156–158
 - refractory urinary incontinence, 159
 - signs and symptoms, 159
 - standard of care, 160, 161
 - surgical indications, 156
 - tethering lesions, 159
- Omphalocele, exstrophy, imperforate anus and spinal defects (OEIS syndrome), 112, 272

- Orthopedic complications, 306, 307
- Orthopedic deformities
- foot deformities, 112
 - habitual posturing/positioning, 110
 - leg length discrepancy, 112
 - muscular imbalance, 110
 - scoliosis, 112
 - spinal column and lower extremities, 110, 111
- P**
- Palpable depression, 6
- Phenotypic abnormalities, 5
- Polar cell polarity (PCP) pathway, 29
- Postovulatory day (POD), 293
- Prickle-1*, 29
- Primitive streak (PS), 18, 19
- Progression-free survival (PFS), 204
- Pseudotails, 77
- R**
- Reissner's fiber (RF), 95
- Ruysch, Frederik, 1
- S**
- Sacro-caudal agenesis, 338, 339
- Sacrococcygeal teratoma, 336, 338
- Scoliosis
- definition, 138
 - diagnostic work-up, 138, 139
 - follow-up, 139
 - with hump, 137, 139–141
 - low back and left leg pain, 141–143
 - symptoms, 138
 - treatment, 139
- Short tethering tracts (STTs), 229
- Somatic ectoderm (cutaneous), 5
- Somatosensory evoked responses (SSEP), 29, 30
- Sonic hedgehog (*Shh*) pathway, 23
- Spina bifida, 224
- cystica, 3
 - occulta, 5, 9
- Split cord malformations (SCMs), 11–13, 38, 40, 62, 63, 92, 116, 121, 123, 127, 130, 131, 197, 258, 259
- anatomy, 210
 - characteristics, 293
 - classification, 210–212
 - clinical outcomes, 216
 - cutaneous stigmata and conditions, 212, 213
 - diastematomyelia, 209, 292
 - embryology, 213, 214
 - history, 214
 - vs. meningocele manqué, 228
 - MRI, 259
 - natural history, 315
 - symptoms, 292
 - surgical outcome, 315, 316
 - treatment, 215, 216
 - Type 1 SCM, 258
 - Type 2 SCM, 259
- Surgical 'untethering', 9, 199
- Syringoarachnoid shunt (SAS), 233
- Syringohydromyelia, 228, 298
- Syringomyelia, 9
- T**
- Terminal syringomyelia
- clinical presentation, 235
 - hydromyelia, 298
 - imaging, 235, 236
 - incidence, 233, 234
 - management, 237–239
 - myelocystocele, 239, 240
 - pathogenesis, 234, 298
 - scoliosis and motor weakness, 299
- Terminal syrinx (TS), 233
- Tethered cord syndrome (TCS), 9, 90, 91, 243
- in adults
 - asymptomatic TC, 144
 - benign and malignant tissue, 123
 - bowel and bladder disturbances, 143
 - clinical presentation, 119–121
 - complications, 146, 147
 - CT imaging, 124–127
 - degenerative processes, 118
 - dermal sinus tract, 116, 134
 - diastematomyelia, 116, 121–123, 130, 131
 - early postoperative CT imaging, 126, 129
 - early postoperative MRI, 126, 128
 - early surgical intervention, 143
 - fatty accumulations, 132
 - filum terminale, 117, 133
 - follow-up, 129, 147
 - foot anomalies, 117
 - gender and age, 119, 144
 - herniation, 123
 - histopathological examinations, 126
 - intradural dermoid masses, 116
 - kidney malformations, 117
 - lipomeningocele, 131

- Tethered cord syndrome (TCS) (*cont.*)
- lipomyelomeningocele, 131–134, 136
 - low back pain, 120, 122
 - lumbar MRI, 123
 - malformations, 117, 121–123
 - meningocele manqué, 134
 - midline cutaneous abnormalities, 120, 122
 - mild static neurological deficits, 117
 - motor weakness, 144
 - muscular weakness, 144
 - neurological deficits, 122
 - neurological examination, 130, 145
 - neurological scoring system, 126, 129
 - neurosurgical deficits, 122
 - normal daily activity, 117
 - pathological conditions, 122
 - pathophysiology, 118, 119
 - patient history, 118
 - plain X-rays, 123
 - rehabilitation programs, 144
 - SCMs, 116, 121–123, 127, 130, 131
 - scoliosis (*see* Scoliosis)
 - single-channel cystometrography, 124
 - spinal cord (intradural) lipomas, 132
 - spinal neurenteric cyst, 133
 - standard management protocol, 117, 120, 134–136
 - surgical approaches, 145, 146
 - surgical outcomes, 145
 - surgical procedure, 132, 136–138
 - symptoms, 144
 - T1-weighted sagittal MRI, 126, 127
 - terminal myelocystocele, 133
 - thoracolumbar scoliosis, 126–129, 131, 132, 134, 135
 - three-dimensional CT, 124, 126
 - untethering procedure, 145
 - urodynamic study, 124, 125, 146
 - anatomical correlations, 152, 153
 - anatomy and embryology, 290
 - Chiari I malformations, 291
 - clinical improvement, 152
 - congenital spinal defects, 151
 - distal cord anomalies, 154
 - dysraphic lesions, 154
 - fatty filum terminale, 245, 248, 290–292
 - folic acid supplementation, 115
 - Fuchs's hypothesis, 290
 - imaging technologies, 115
 - LMMC, 290
 - low conus medullaris, 153, 245, 246
 - lumbosacral cutaneous findings, 249
 - non-diagnostic imaging, 248
 - OTCS (*see* Occult tethered cord syndrome (OTCS))
 - physiological/experimental, 153
 - stretch-induced cord dysfunction, 154
 - surgical series, 153, 154
 - thickened filum terminale, 245
 - VACTERL, 291
 - ventriculus terminalis, 247
- Tethered spinal cord (TSC), 10, 11, 194
- natural history, 312, 313
 - surgical treatment, 313, 314
- Thoracolumbar scoliosis, 126–129, 131, 132, 134, 135
- Tight filum, 274
- Tulp, Nicolaes, 1
- U**
- Urologic dysfunction
- bowel incontinence, 110
 - chronic constipation, 110
 - detrusor hyperactivity, 110
 - detrusor-sphincter dyssnergia, 110
 - progressive bladder dysfunction, 110
 - sexual dysfunction, 110
 - subclinical findings, 110
- Urology
- bladder development and function, 266
 - clinical outcome, 275
 - complications, 307, 308
 - follow-up, 276
 - imaging, 267
 - incidence and prevalence, 265
 - neurosurgical intervention
 - asymptomatic occult spinal dysraphism, 273
 - complex lipomyelomeningocele, 274, 275
 - spinal malformations, 275
 - “tight filum”/occult tethered cord syndrome, 274
 - newborn with urogenital/anorectal anomalies, 272
 - newborns and infants with cutaneous stigmata, 272
 - pathophysiology, 266, 267
 - in toddlers and young children, dysfunctional voiding, 273
 - urodynamics, 268, 269, 271

V

Ventriculus terminalis (VT), 87, 94, 95
Vertebral anomalies, anorectal defects, cardiac
defects, tracheoesophageal fistula,
renal abnormalities and limb
deformity (VACTERL), 112, 272
Vertebral development, 332
Virchow, Rudolf, 5, 6
Voiding dysfunction, 266, 274
von Recklinghausen, Friedrich Daniel, 7

X

X-rays, 13, 123, 138–140, 156, 158, 215, 243,
253, 260

Y

Yamada, Shokei, 14