

INDEX FOR RECT VOL. 235

- Accumulation in bivalves, paralytic shellfish toxins, **235: 9**
- Accumulation rates, paralytic shellfish toxins, **235: 10**
- Ag:Si (dissolved), vs. oxygen profiles in the Pacific Ocean (diags), **235: 40**
- Agricultural soils in Newport Bay, DDT concentrations (table), **235: 54**
- Agricultural soils, chlordane, **235: 137**
- Agricultural soils, DDT, **235: 54**
- Agricultural soils, toxaphene (table), **235:146**
- Algal toxin, saxitoxin, **235: 2**
- American avocet, DDT residues, **235: 70**
- Amphipod toxicity, metals & organics in sediment (table), **235: 142**
- Analysis methods, paralytic shellfish toxins, **235: 10**
- Anchovies, DDE residue decreases (table), **235: 122**
- Atlantic menhaden, DDT & metabolite residues (table), **235: 123**
- Avian threat, DDT residues over time, **235: 93**
- Benthic triad, impairment analysis, **235: 141, 142**
- Bioaccumulation, DDT in Oregon, **235: 109**
- Bioassay methods, paralytic shellfish toxins, **235: 12**
- Biogeochemical cycling, oceanic silver, **235: 27 ff.**
- Biogeochemistry, silver cycling, **235: 29**
- Biomagnification factor, ospreys, **235: 114**
- Biotransformation, paralytic shellfish toxins, **235: 15**
- Biotransformation, paralytic shellfish toxins, **235: 1ff.**
- Bird eggshell thickness (table), **235: 132**
- Bird residues, DDT, **235: 61**
- Bivalve accumulation & depuration, paralytic shellfish toxins, **235: 9**
- Bivalve detoxification, paralytic shellfish toxins (table), **235: 11**
- Black skimmers, DDT residues, **235: 69**
- Black-necked stilts, DDT residues, **235: 69**
- Brown pelican eggs, DDE levels (diag.), **235: 122**
- Brown pelican eggs, DDE residue decreases (table), **235: 122**
- Brown pelican eggs, DDT & metabolites (diag.), **235: 126**
- Brown pelican eggs, organochlorine residue decline, **235: 125**
- Brown pelican eggshell thinning, DDE dose-response (diag.), **235: 120**
- Brown pelican recovery, California (table), **235: 121**
- Brown pelican reproduction, recovery (table), **235: 124**
- Brown pelican reproductive effects, DDE threshold (diag.), **235: 128**
- California fish guidance on DDT, human health protection, **235: 135**
- California fish guidance on PCBs, human health protection, **235:152**
- Canadian fish guidance, wildlife protection, **235: 129**
- Cetaceans & DDT residues, P450 isozymes, **235: 81**
- Chemical assay methods, paralytic shellfish toxins, **235: 13**
- Chlordane hazard assessment, Newport Bay & watershed, **235: 49 ff**

- Chlordane levels, water column, fish & mussels, **235**: 139
- Chlordane residues, mussels (diag.), **235**: 141
- Chlordane, described, **235**: 50
- Chlordane, environmental levels, **235**: 137
- Chlordane, *Hyallolela azteca* toxicity (table), **235**: 145
- Chlordane, in agricultural soils (table), **235**: 138
- Chlordane, in agricultural soils, **235**: 137
- Chlordane, in Newport Bay & watershed sediments (diag.), **235**: 138
- Chlordane, red shiner levels (diags.), **235**: 140
- Chlordane, toxicity thresholds, **235**: 144
- Climate described, Newport Bay & watershed, **235**: 51
- Coastal waters, dissolved silver concentrations (table), **235**: 32
- Contaminant mixture issues, TEL data points, **235**: 86
- Copper concentrations, Mid-Atlantic ridge plumes (table), **235**: 32
- Copper, vs. silver in oceanic waters (diag.), **235**: 36
- Cormorant eggs, DDE residues (table), **235**: 66
- Cormorant eggshells, thickness (table), **235**: 66
- Cormorants, DDT residues, **235**: 64
- Cormorants, eggshell thinning & DDE (diag.), **235**: 65
- DDD residues, osprey prey fish (table), **235**: 105
- DDE & DDT, osprey eggshell thinning, **235**: 96
- DDE bioaccumulation, Oregon, **235**: 109
- DDE dose-response, brown pelican eggshell thinning (diag.), **235**: 120
- DDE egg residues, osprey population status (table), **235**: 97
- DDE egg residues, vs. osprey productivity (diag.), **235**: 102
- DDE exposure effect, kestrel eggshell thickness (diag.), **235**: 132
- DDE levels, brown pelican eggs (diag.), **235**: 122
- DDE pelican egg residues, inverse eggshell thickness (diag.), **235**: 124
- DDE reproductive effects, threshold for brown pelican (diag.), **235**: 128
- DDE residue decline, osprey recovery, **235**: 98
- DDE residue decreases, anchovies & brown pelican eggs (table), **235**: 122
- DDE residue plot, brown pelican eggshell thinning (diag.), **235**: 120
- DDE residues, in Texas cormorant eggs (table), **235**: 66
- DDE residues, largescale suckers (diag.), **235**: 111
- DDE residues, osprey egg (table), **235**: 104
- DDE residues, osprey eggshell thinning (table), **235**: 110
- DDE residues, osprey prey fish (table), **235**: 105
- DDE residues, pelican recovery, **235**: 124
- DDE residues, species variation & eggshell thinning (diag.), **235**: 119
- DDE residues, vs. osprey eggshell thickness (diag.), **235**: 102
- DDE residues, vs. pelican eggshell thickness (diag.), **235**: 127
- DDE, eggshell thinning of pelicans, **235**: 116
- DDE, pelican eggshell thinning (diag.), **235**: 118
- DDT & eggshell thickness, birds (table), **235**: 132
- DDT & human health, California fish guidance, **235**: 135
- DDT & metabolites, brown pelican eggs (diag.), **235**: 126
- DDT & metabolites, in emitted Atlantic menhaden (table), **235**: 123
- DDT & peregrine falcons, wintering ground exposures (table), **235**: 106
- DDT & porpoises, stress hormone induction, **235**: 79
- DDT agricultural soil samples, locations (illus.), **235**: 56
- DDT concentration trends, fish species, **235**: 59
- DDT concentrations, agricultural soils, **235**: 54
- DDT concentrations, Newport Bay agricultural soils (table), **235**: 54
- DDT dissipation, process summarized, **235**: 54
- DDT effect data sets, fresh & marine water sediments (table), **235**: 85
- DDT effect discrepancies, fresh- & marine-water sediments, **235**: 85
- DDT hazard assessment, Newport Bay & watershed, **235**: 49 ff.
- DDT levels, in the environment, **235**: 53
- DDT levels, Newport Bay & watershed water column (table), **235**: 58
- DDT release proximity, dolphin residues, **235**: 79

- DDT residue date, in sediments, **235: 55**
- DDT residue differences, stranded vs. harvested whales, **235: 80**
- DDT residue trends, fish, **235: 103**
- DDT residues in cetaceans, P450 isozymes, **235: 81**
- DDT residues in whales, gender differences, **235: 81**
- DDT residues over time, avian threat, **235: 93**
- DDT residues, American avocet, **235: 70**
- DDT residues, and soil erosion, **235: 55**
- DDT residues, black skimmers, **235: 69**
- DDT residues, black-necked stilts, **235: 69**
- DDT residues, cormorants, **235: 64**
- DDT residues, dolphins, **235: 78**
- DDT residues, in birds, **235: 61**
- DDT residues, in fish (table), **235: 94**
- DDT residues, in fish of Great Gull island (table), **235: 91**
- DDT residues, in marine mammals, **235: 82**
- DDT residues, in Montana osprey eggs (table), **235: 100**
- DDT residues, in Newport Bay & watershed mussels (diag.), **235: 60**
- DDT residues, in osprey eggs (table), **235: 93**
- DDT residues, in sea lions, **235: 75**
- DDT residues, in seals, **235: 73**
- DDT residues, in whales, **235: 80**
- DDT residues, killdeer, **235: 70**
- DDT residues, Newport Bay region dolphins, **235: 80**
- DDT residues, Newport Bay sediment data (diag.), **235: 57**
- DDT residues, osprey egg (table), **235: 104**
- DDT residues, ospreys, **235: 93**
- DDT residues, terns, **235: 62**
- DDT residues, vs. toxicity in marine sediments, **235: 87**
- DDT residues, water column, fish & mussels, **235: 58**
- DDT residues, white sucker tissues (diag.), **235: 103**
- DDT safe levels, fresh water organisms, **235: 88**
- DDT safety, NAS recommendation, **235: 92**
- DDT TEL calculation, issues, **235: 89**
- DDT TEL levels, in sediment, **235: 83**
- DDT TEL values, in Newport Bay region, **235: 84**
- DDT TMDL targets, in sediment, fish & water (table), **235: 83**
- DDT TMDL values, in Newport Bay & watershed, **235: 83**
- DDT toxicity, vs. residues, **235: 87**
- DDT, and eggshell thinning, **235: 61**
- DDT, described, **235: 50**
- DDT, in marine mammals, **235: 72**
- DDT, pelicans, **235: 115**
- DDT, red shiner residues in San Diego Creek (diag.), **235: 58**
- DDT, wildlife protection guidance, **235: 114**
- DDT:DDE egg residue ratio, exposure timing, **235: 107**
- Depuration in bivalves, paralytic shellfish toxins, **235: 9**
- Detoxification in bivalves, paralytic shellfish toxins (table), **235: 11**
- Detoxification rates, paralytic shellfish toxins, **235: 10**
- Detoxification, paralytic shellfish toxins, **235: 10**
- Dissolved silver concentrations, estuarine & coastal waters (table), **235: 32**
- Dolphin residues, of DDT, **235: 78**
- Dolphin residues, proximity to DDT release, **235: 79**
- Dolphins & DDT residues, from Newport Bay, **235: 80**
- Effects range median (ERM), chlordane (table), **235: 143**
- Eggshell thickness & DDT, birds (table), **235: 132**
- Eggshell thickness, changes in New Jersey (table), **235: 97**
- Eggshell thickness, osprey eggs (table), **235: 104**
- Eggshell thickness, vs. DDE residues, **235: 124**
- Eggshell thinning & DDE, cormorants (diag.), **235: 65**
- Eggshell thinning & species variation, DDE residues (diag.), **235: 119**
- Eggshell thinning in brown pelican, DDE dose-response (diag.), **235: 120**
- Eggshell thinning in brown pelicans, DDE residue plot (diag.), **235: 120**
- Eggshell thinning in ospreys, DDE residues (table), **235: 110**
- Eggshell thinning in ospreys, DDE, **235: 96**
- Eggshell thinning of pelicans, DDE, **235: 116**
- Eggshell thinning, and DDT, **235: 61**
- Environmental effects, harmful algal blooms, **235: 7**
- Environmental levels, chlordane, **235: 137**
- Environmental levels, toxaphene, **235: 146**
- Environmental release, silver, **235: 28**

- ERM (effects range median), chlordane (table), **235: 143**
- Estuarine waters, dissolved silver concentrations (table), **235: 32**
- Eutrophication, influences on harmful algal blooms, **235: 6**
- Exposure timing, DDT:DDE egg residue levels, **235: 107**
- Fish concentration trends, DDT, **235: 59**
- Fish DDT residue trends, 1970's, **235: 103**
- Fish fillets from Newport Bay, PCB residues (table), **235:151**
- Fish levels, chlordane, **235: 139**
- Fish residues, DDT (table), **235: 94**
- Fish residues, DDT in Newport Bay area, **235: 58**
- Fish residues, of DDT (table), **235: 91**
- Fish residues, toxaphene, **235:147**
- Fish TMDL targets, for DDT (table), **235: 83**
- Fluxes, of oceanic silver, **235: 27 ff.**
- Fresh water safe levels, DDT residues, **235: 88**
- Fresh water sediments, effect data sets for total DDT (table), **235: 85**
- Great Lakes, osprey recovery (diags.), **235: 108**
- Harmful algal bloom species, toxins produced (table), **235: 5**
- Harmful algal blooms, defined, **235: 2**
- Harmful algal blooms, environmental effects, **235: 7**
- Harmful algal blooms, eutrophication effects, **235: 6**
- Harmful algal blooms, factors influencing toxicity, **235: 6**
- Harmful algal blooms, light intensity effects, **235: 8**
- Harmful algal blooms, nutrient effects, **235: 7**
- Harmful algal blooms, temperature effects, **235: 8**
- Human hazard assessment, DDT in Newport Bay & watershed, **235: 49 ff.**
- Human health protection & DDT, California fish guidance, **235: 135**
- Human health protection, California guidance on PCBs, **235:152**
- Hyallela azteca* toxicity, chlordane (table), **235: 145**
- Hydrology described, of Newport Bay and watershed, **235: 51**
- Immunosorbant assays, paralytic shellfish toxins, **235: 14**
- Intoxication symptoms, paralytic shellfish toxins, **235: 2**
- Kestrel eggshell thickness, DDE exposure effect (diag.), **235: 132**
- Killdeer, DDT residues, **235: 70**
- Light intensity effects, harmful algal blooms, **235: 8**
- Marine mammals, and DDT residues, **235: 82**
- Marine mammals, DDT residues, **235: 72**
- Marine sediment, silver cycling, **235: 39**
- Marine sediments, silver concentrations (table), **235: 31**
- Marine toxins, paralytic shellfish, **235: 1ff.**
- Marine water sediments, effect data sets for total DDT (table), **235: 85**
- Metals in sediments, amphipod toxicity (table), **235: 142**
- Mid-Atlantic ridge plumes, Ag, Cu & silicate concentrations (table), **235: 32**
- Mode of action, paralytic shellfish toxins, **235: 2**
- Mussel levels in Newport Bay region, toxaphene (diag.), **235:148**
- Mussel levels, chlordane (diag.), **235: 141**
- Mussel levels, chlordane, **235: 139**
- Mussel residues of DDT, Newport Bay & watershed (diag.), **235: 60**
- Mussel residues, DDT in Newport Bay area, **235: 58**
- Mussels, toxaphene, **235:147**
- NAS (National Academy of Science) guidance, to protect wildlife, **235: 90**
- NAS fish guidance on toxaphene, wildlife protection, **235:149**
- NAS recommendation, DDT safety, **235: 92**
- National Academy of Science (NAS), wildlife protection, **235: 90**
- New York sediment guidance, toxaphene, **235:149**
- Newport Bay & watershed mussels, DDT residues (diag.), **235: 60**
- Newport Bay & watershed water column, DDT levels (table), **235: 58**
- Newport Bay & watershed, chlordane in mussels (diag.), **235: 141**
- Newport Bay & watershed, DDT, chlordane, toxaphene & PCB hazard assessment, **235: 49 ff.**
- Newport Bay & watershed, map (illus.), **235: 51**
- Newport Bay & watershed, TMDL values for DDT, **235: 83**
- Newport Bay agricultural soils, DDT concentrations (table), **235: 54**
- Newport Bay and watershed, described, **235: 51**
- Newport Bay area, DDT in fish & mussels, **235: 58**

- Newport Bay area, DDT residues in birds, **235: 61**
- Newport Bay dolphins, DDT residues, **235: 80**
- Newport Bay region sediment, chlordane levels (diag.), **235: 138**
- Newport Bay region, DDT TEL values, **235: 84**
- Newport Bay sediment date, DDT residues (diag.), **235: 57**
- Newport Bay, land use, **235: 52**
- Newport Bay, water quality, **235: 52**
- Nutrient effects, harmful algal blooms, **235: 7**
- Oceanic contaminant, silver, **235: 28**
- Oceanic contamination profile, silver (illus.), **235: 31**
- Oceanic contamination, by silver (table), **235: 30**
- Oceanic contour plot, silver concentrations (illus.), **235: 37**
- Oceanic data, silver contamination, **235: 29**
- Oceanic fixation, silver, **235: 41**
- Oceanic silver, data comparisons, **235: 42**
- Oceanic silver, measurements, **235: 33**
- Oceanic silver, vertical concentration profiles (diag.), **235: 34**
- Organ elimination, bivalve paralytic shellfish toxins, **235: 10**
- Organics in sediments, amphipod toxicity (table), **235: 142**
- Organochlorine residue decline, brown pelican eggs, **235: 125**
- Osprey egg residues, DDE & DDD (table), **235: 104**
- Osprey eggs, DDT residues (table), **235: 100**
- Osprey eggs, DDT residues (table), **235: 93**
- Osprey eggs, nesting productivity (table), **235: 100**
- Osprey eggshell thickness, vs. DDE residues (diag.), **235: 102**
- Osprey eggshell thinning, DDE residues (table), **235: 110**
- Osprey population status, DDE egg residues (table), **235: 97**
- Osprey populations, reproductive success (table), **235: 95**
- Osprey prey fish, DDE & DDD residues (table), **235: 105**
- Osprey productivity, New Jersey (table), **235: 105**
- Osprey productivity, vs. DDE egg residues (diag.), **235: 102**
- Osprey recovery, from DDT effects, **235: 98**
- Osprey recovery, Great Lakes (diags.), **235: 108**
- Osprey recovery, in Connecticut-Long Island (diag.), **235: 99**
- Osprey reproduction, Newport Bay, **235: 112**
- Osprey reproduction, San Joaquin Wildlife Sanctuary (table), **235: 113**
- Ospreys & DDE, eggshell thinning, **235: 96**
- Ospreys, biomagnifications factor, **235: 114**
- Ospreys, DDT residues, **235: 93**
- Ospreys, declining populations, **235: 94**
- Ospreys, eggshell thickness (table), **235: 104**
- Ospreys, population size & productivity (diag.), **235: 101**
- Ospreys, population status (table), **235: 95**
- Ospreys, wintering locations (illus.), **235: 111**
- P450 isozymes, cetaceans & DDT residues, **235: 81**
- Pacific Ocean oxygen profiles, vs. Ag:Si (dissolved) (diags.), **235: 40**
- Paralytic shell fish toxins, organisms responsible, **235: 3**
- Paralytic shellfish toxin composition, salinity effect, **235: 8**
- Paralytic shellfish toxins, analysis methods, **235: 10**
- Paralytic shellfish toxins, bioassay methods, **235: 12**
- Paralytic shellfish toxins, biotransformation & detoxification, **235: 1ff.**
- Paralytic shellfish toxins, biotransformation, **235: 15**
- Paralytic shellfish toxins, chemical assay methods, **235: 13**
- Paralytic shellfish toxins, clinical symptoms, **235: 2**
- Paralytic shellfish toxins, detoxification in bivalves (table), **235: 11**
- Paralytic shellfish toxins, immunosorbant assays, **235: 14**
- Paralytic shellfish toxins, mode of action, **235: 2**
- Paralytic shellfish toxins, organ elimination & accumulation rates, paralytic shellfish toxins, **235: 10**
- Paralytic shellfish toxins, paralytic shellfish poisoning, **235: 2**
- Paralytic shellfish toxins, structure & toxicity (diag., table), **235: 4**
- Paralytic shellfish toxins, treatments, **235: 3**
- PCB (polychlorinated biphenyls), sport fish levels, **235: 150**
- PCB hazard assessment, Newport Bay & watershed, **235: 49 ff**
- PCB residues, fish fillets from Newport Bay (table), **235: 151**

- PCBs, California guidance to protect human health, **235:152**
- PCBs, described, **235: 50**
- Pelican eggshell thickness, vs. DDE residues (diag.), **235: 127**
- Pelican eggshell thickness, vs. DDE residues, **235: 124**
- Pelican eggshell thinning, DDE (diag.), **235: 118**
- Pelican eggshell thinning, DDE, **235: 116**
- Pelican recovery, DDE residues, **235: 124**
- Pelicans, DDT, **235: 115**
- Peregrine falcons & DDT, wintering ground exposures (table), **235: 106**
- Polychlorinated biphenyls (PCBs), sport fish levels, **235:150**
- Population status, ospreys (table), **235: 95**
- Porpoises & DDT, stress hormone induction, **235: 79**
- Red shiner levels, chlordane (diags.), **235: 140**
- Red shiner levels, toxaphene (diags.), **235:147, 148**
- Red shiner residues in San Diego Creek, DDT (diag.), **235: 59**
- Red tide, described, **235: 1**
- Reproductive success, U.S. osprey populations (table), **235: 95**
- Salinity effect, paralytic shellfish toxin composition, **235: 8**
- Salinity effect, silver concentration in estuarine waters, **235: 33**
- San Joaquin Wildlife Sanctuary, osprey reproduction (table), **235: 113**
- Saxitoxin, algal toxin, **235: 2**
- Saxitoxins, categories, **235: 3**
- Sea lions, DDT residues, **235: 75**
- Seals, DDT residues, **235: 73**
- Sediment data, DDT in Newport Bay (diag.), **235: 57**
- Sediment date, DDT, **235: 55**
- Sediment guidance, toxaphene, **235:149**
- Sediment levels, chlordane, **235: 138**
- Sediment loads, vs. fate of organochlorines, **235: 53**
- Sediment TMDL targets, for DDT (table), **235: 83**
- Sediments, toxaphene levels, **235:146**
- Selenium concentrations, vs. silver in water & aerosols (diags.), **235: 38**
- Silicate concentrations, Mid-Atlantic ridge plumes (table), **235: 32**
- Silicate, vs. silver in oceanic waters (diag.), **235: 35**
- Silver concentrations in N. Pacific, temporal variations (diag.), **235: 37**
- Silver concentrations, marine sediments (table), **235: 31**
- Silver concentrations, Mid-Atlantic ridge plumes (table), **235: 32**
- Silver concentrations, oceanic contour plot (illus.), **235: 37**
- Silver concentrations, seawater reference material (table), **235: 43**
- Silver concentrations, ultraviolet light effects, **235: 42**
- Silver concentrations, vs. selenium in water & aerosols (diags.), **235: 38**
- Silver contaminant profiles, oceans (illus.), **235: 31**
- Silver contamination, oceanic data, **235: 29**
- Silver contamination, oceans (table), **235: 30**
- Silver cycling, marine sediment, **235: 39**
- Silver fixation, ocean, **235: 41**
- Silver in the oceans, data comparisons, **235: 42**
- Silver oceanic distribution, low oxygen effect, **235: 39**
- Silver profiles, vertical oceanic concentrations (diag.), **235: 34**
- Silver, as oceanic contaminant, **235: 28**
- Silver, biogeochemical cycling, **235: 29**
- Silver, early production and use, **235: 28**
- Silver, oceanic biogeochemical cycling, **235: 27 ff.**
- Silver, oceanic sources & fluxes, **235: 27 ff.**
- Silver, sources in environment, **235: 28**
- Silver, toxicity issues, **235: 28**
- Silver, vs. dissolved copper in oceanic waters (diag.), **235: 36**
- Silver, vs. silicate in oceanic waters (diag.), **235: 35**
- Silver, water concentrations, **235: 33**
- Soil erosion, and DDT residues, **235: 55**
- Sources & cycling, oceanic silver, **235: 27 ff.**
- Species variation & eggshell thinning, DDE residues (diag.), **235: 119**
- Stress hormone induction, DDT in porpoises, **235: 79**
- Suckers (largescale), & DDE residues (diag.), **235: 111**
- TEL (threshold effect levels) for DDT, in sediment, **235: 83**
- TEL calculation issues, DDT residues, **235: 89**
- TEL data points, contaminant mixtures, **235: 86**
- TEL use issues, TMDL sediment target, **235: 90**

- TEL values for DDT, in Newport Bay region, **235: 84**
- Temperature effects, harmful algal blooms, **235: 8**
- Temporal variations, silver concentrations in N. Pacific (diag.), **235: 37**
- Terns, DDT residues, **235: 62**
- Threshold effect levels (TEs) for DDT, in sediment, **235: 83**
- TMDL (total maximum daily load) targets for DDT, in Newport Bay & watershed, **235: 83**
- TMDL targets for DDT, in sediment, fish & water (table), **235: 83**
- Total maximum daily load (TMDL) targets for DDT, in Newport Bay & watershed, **235: 83**
- Toxaphene decay, influencing factors, **235:149**
- Toxaphene levels, Newport Bay & watershed mussels (table), **235:148**
- Toxaphene levels, sediments, **235:146**
- Toxaphene residues, water column, fish & mussels, **235:147**
- Toxaphene, described, **235: 50**
- Toxaphene, environmental levels, **235:146**
- Toxaphene, hazard assessment, Newport Bay & watershed, **235: 49 ff**
- Toxaphene, in agricultural soils (table), **235:146**
- Toxaphene, NAS fish guidance, **235:149**
- Toxaphene, New York sediment guidance, **235:149**
- Toxaphene, red shiner levels (diags), **235:147, 148**
- Toxicity issues, silver, **235: 28**
- Toxicity of harmful algal blooms, influencing factors, **235: 6**
- Toxicity thresholds, chlordane, **235: 144**
- Toxicity, paralytic shellfish toxins (table), **235:4**
- Treatment, paralytic shellfish toxins, **235: 3**
- U.S. EPA fish guidance, wildlife protection, **235: 134**
- Ultraviolet light effects, silver concentrations, **235: 42**
- Water column guidance on DDT, to protect wildlife, **235: 114**
- Water column levels, chlordane, **235: 139**
- Water column of Newport Bay & watershed, DDT levels (table), **235: 58**
- Water column, toxaphene, **235:147**
- Water concentrations, silver, **235: 33**
- Water quality, Newport Bay, **235: 52**
- Water TMDL targets, for DDT (table), **235: 83**
- Whale DDT residues, stranded vs. harvested specimens, **235: 80**
- Whales, DDT residues, **235: 80**
- White sucker tissues, DDT residues (diag.), **235: 103**
- Wildlife hazard assessment, DDT in Newport Bay & watershed, **235: 49 ff.**
- Wildlife protection, Canadian fish guidance, **235: 129**
- Wildlife protection, NAS guidance, **235: 90**
- Wildlife protection, U.S. EPA fish guidance, **235: 134**
- Wildlife protection, U.S. EPA water column guidance on DDT, **235: 114**
- Wintering ground exposures to DDT, peregrine falcons (table), **235: 106**