

Conclusion

The production of high quality noodles and pasta products have been associated largely with particular physicochemical properties of the protein components in the dough. Other flour constituents and added ingredients also influence the desirable characteristics and organoleptic qualities. The optimized conditions of constituents, ingredients, and processing techniques would give the optimum quality of yellow alkaline noodles.

References

- Ahmad I, Anjum FM, Butt MS, Bajwa BE (2007) Improvement in spring wheat quality in Pakistan. *Pakistan J Agric Res* 20(1/2):1–6
- Akashi H, Takahashi M, Endo S (1999) Evaluation of starch properties of wheats used for Chinese yellow-alkaline noodles in Japan. *Cereal Chem* 76(1):50–55
- Arif S, Ahmed M, Khazada KK (2006) Quality evaluation of some sindh (Pakistan) wheat varieties. II. Correlation among various quality traits. *Pak J Sci Ind Res* 49(4):285–289
- Asenstorfer RE, Wang Y, Mares DJ (2006) Chemical structure of flavonoid compounds in wheat (*Triticum aestivum* L.) flour that contribute to the yellow colour of Asian alkaline noodles. *J Cereal Sci* 43(1):108–119
- Asenstorfer RE, Appelbee MJ, Mares DJ (2009) Physical–Chemical analysis of non-polyphenol oxidase (Non-PPO) darkening in yellow alkaline noodles. *J Agric Food Chem* 57(12):5556–5562
- Asenstorfer RE, Appelbee MJ, Mares DJ (2010) Impact of protein on darkening in yellow alkaline noodles. *J Agric Food Chem* 58:4500–4507
- Asenstorfer RE, Appelbee MJ, Kuszniir CA, Mares DJ (2014) Towards an understanding of mechanisms involved non-PPO darkening in yellow alkaline noodles (YAN). *J Agric Food Chem* 62:4725–4730
- Baik BK, Lee MR (2003) Effect of starch amylose content of wheat on textural properties of white salted noodles. *Cereal Chem* 80:304–9
- Bellido GG, Hatcher DW (2009) Stress relaxation behaviour of yellow alkaline noodles: effect of deformation history. *J food Eng* 93(4):460–467
- Bellido GG, Hatcher DW (2011) Effects of a cross-linking enzyme on the protein composition, mechanical properties, and microstructure of Chinese-style noodles. *Food chem* 125(3):813–822
- Bhattacharya M, Corke H (1996) Selection of desirable starch pasting properties in wheat for use in white salted or yellow alkaline noodles. *Cereal Chem* 73(6):721–728
- Chakraborty M, Hareland GA, Manthey FA, Berglund LR (2003) Evaluating quality of yellow alkaline noodles made from mechanically abraded sprouted wheat. *J Sci Food Agric* 83(5):487–495
- Chang HC, Wu LC (2008) Texture and quality properties of Chinese fresh egg noodles formulated with green seaweed (*Monostroma nitidum*) powder. *J Food Sci* 73(8):S398–S404
- Chin CK, Huda N, Yang TA (2012) Incorporation of surimi powder in wet yellow noodles and its effects on the physicochemical and sensory properties. *Int Food Res J* 2:701–707
- Chung CE, Lee KW, Cho MS (2010) Noodle consumption patterns of American consumers: NHANES 2001–2002. *Nutr Res Pract* 4(3):243–251
- Cole ME (1991) Prediction and measurement of pasta quality. *Int J Food Sci Technol* 26(2):133–151

- Crosbie GB, Ross AS, Moro T, Chiu PC (1999) Starch and protein quality requirements of Japanese alkaline noodles (Ramen). *Cereal Chem* 76(3):328–334
- Deng Z, Tian J, Zhao L, Zhang Y, Sun C (2008) High temperature-induced changes in high molecular weight glutenin subunits of Chinese winter wheat and its influences on the texture of Chinese noodles. *J Agron Crop Sci* 194(4):262–269
- Devlieghere F, Vermeiren L., Debevere J (2004) New preservation technologies: possibility and limitation. *Int Dairy J* 14:273–285
- Edwards NM, Scanlon MG, Kruger JE, Dexter JE (1996) Oriental noodle dough rheology: relationship to water absorption, formulation, and work input during dough sheeting. *Cereal Chem* 73:708–711
- Ekthamasut K (2006) Effect of tomato seed meal on wheat pasting properties and alkaline noodle qualities. *Aust J Technol* 9:147–152
- Foo WT, Yew HS, Liong MT, Azhar ME (2011) Influence of formulations on textural, mechanical and structural breakdown properties of cooked yellow alkaline noodles. *Int Food Res J* 18(4):1295–1301
- Fu BX (2008) Asian noodles: history, classification, raw materials, and processing. *Food Res Int* 41(9):888–902
- Fuerst EP, Anderson JV, Morris CF (2006) Delineating the role of polyphenol oxidase in the darkening of alkaline wheat noodles. *J Agric Food Chem* 54(6):2378–2384
- Galvez FCF, Resurreccion AV (1992) Reliability of the focus group technique in determining the quality characteristics of mungbean [*Vigna radiata* (L.) wilczec] noodles. *J Sens Stud* 7(4):315–326
- Gan CY, Ong WH, Wong LM, Easa AM (2009) Effects of ribose, microbial transglutaminase and soy protein isolate on physical properties and in-vitro starch digestibility of yellow noodles. *LWT-Food Sci Technol* 42(1):174–179
- Garbutt J (1997) *Essentials of food microbiology*. Hodder Headline Group, London, pp 124–129
- Ghaffar S, Abdulmir AS, Bakar FA, Karim R, Saari N (2009) Microbial growth, sensory characteristic and pH as potential spoilage indicators of Chinese yellow wet noodles from commercial processing plants. *Am J Appl Sci* 6(6):1059
- Gulia N, Khatkar BS (2014) Relationship of dough thermomechanical properties with oil uptake, cooking and textural properties of instant fried noodles. *Food Sci Technol Int* 20(3):171–182
- Gulia N, Dhaka V, Khatkar BS (2014) Instant noodles: processing, quality, and nutritional aspects. *Crit Rev Food Sci Nutr* 54(10):1386–1399
- Hatcher DW, Anderson MJ (2007) Influence of alkaline formulation on oriental noodle color and texture 1. *Cereal Chem* 84(3):253–259
- Hatcher DW, Kruger JE, Anderson MJ (1999) Influence of water absorption on the processing and quality of oriental noodles. *Cereal Chem* 76:566–572
- Hatcher DW, Anderson MJ, Desjardins RG, Edwards NM, Dexter JE (2002) Effects of flour particle size and starch damage on processing and quality of white salted noodles. *Cereal Chem* 79(1):64–71
- Hatcher DW, Bellido GG, Dexter JE, Anderson MJ, Fu BX (2008a) Investigation of uniaxial stress relaxation parameters to characterize the texture of yellow alkaline noodles made from durum and common wheats. *J Texture Stud* 39:695–708
- Hatcher DW, Edwards NM, Dexter JE (2008b) Effects of particle size and starch damage of flour and alkaline reagent on yellow alkaline noodle characteristics. *Cereal Chem* 85(3):425–432
- Hatcher DW, Dexter JE, Anderson MJ, Bellido GG, Fu BX (2009a) Effect of blending durum wheat flour with hard white wheat flour on the quality of yellow alkaline noodles. *Food Chem* 113(4):980–988
- Hatcher DW, Bellido GG, Anderson MJ (2009b) Flour particle size, starch damage, and alkali reagent: impact on uniaxial stress relaxation parameters of yellow alkaline noodles. *Cereal Chem* 86(3):361–368
- Heo H, Kang CS, Woo SH, Lee KS, Choo BK, Park CS (2012) Characteristics of yellow alkaline noodles prepared from Korean wheat cultivar. *Food Sci Biotechnol* 21(1):69–81
- Hou G (2001) Oriental noodles. *Adv Food Nutr Res* 43:141–193

- Hou GG (2010) Asian noodles: science, technology, and processing. Wiley
- Hou G, Kruk M (1998) Asian noodle technology. *Tech Bull* 20(12):1–10
- Huang YC, Lai HM (2010) Noodle quality affected by different cereal starches. *J Food Eng* 97(2):135–143
- Huang S, Morrison WR (1988) Aspects of proteins in Chinese and British common (hexaploid) wheats related to quality of white and yellow Chinese noodles. *J Cereal Sci* 8(2):177–187
- Hung PV, Hatcher DW (2011) Ultra-performance liquid chromatography (UPLC) quantification of carotenoids in durum wheat: influence of genotype and environment in relation to the colour of yellow alkaline noodles (YAN). *Food Chem* 125(4):1510–1516
- Ito M, Ohta K, Nishio Z, Tabiki T, Hashimoto N, Funatsuki W, Miura H, Yamauchi H (2007) Quality evaluation of yellow alkaline noodles made from the Kitanokaori wheat cultivar. *Food Sci Technol Res* 13(3):253–260
- Jensen N, Hocking AD, Miskelly D, Berghofer LK (2004) Microbiological safety of high moisture noodles. 1. Marketplace survey of noodles sold in Australia. *Food Aust* 56(3):71–74
- Jiang H, Martin J, Okot-Kotber M, Seib PA (2011) Color of whole-wheat foods prepared from a bright-white hard winter wheat and the phenolic acids in its coarse bran. *J Food Sci* 76(6):C846–C852
- Jun WJ, Seib PA, Chung OK (1998) Characteristics of noodle flours from Japan. *Cereal Chem* 75:820–825
- Keyimu XG (2013) The effects of using seaweed on the quality of Asian noodles. *J Food Process Technol* 4(216):2
- Kim BY, Lee JY, Ha SD (2011) Growth characteristics and development of a predictive model for *Bacillus cereus* in fresh wet noodles with added ethanol and thiamine. *J Food Prot* 74(4):658–664
- Konik CM, Mikkelsen LM, Moss R, Gore PJ (1994) Relationship between physical starch properties and yellow alkaline noodle quality. *Starch* 46:292–299
- Kruger JE, Matsuo RR, Preston K (1992) A comparison of methods for the prediction of Cantonese noodle colour. *Can J Plant Sci* 72:1021–1029
- Kruger JE, Hatcher DW, DePauw R (1994) A whole seed assay for polyphenol oxidase in Canadian prairie spring wheats and its usefulness as a measure of noodle darkening. *Cereal Chem* 71(4):324–326
- Li M, Zhu K, Guo X, Peng W, Zhou H (2011) Effect of water activity (a_w) and irradiation on the shelf-life of fresh noodles. *Innov Food Sci Emerg Technol* 12(4):526–530
- Li M, Peng J, Zhu KX, Guo XN, Zhang M, Peng W, Zhou HM (2013) Delineating the microbial and physical-chemical changes during storage of ozone treated wheat flour. *Innovative Food Sci Emerg Technol* 20:223–229
- Li M, Zhu KX, Peng J, Guo XN, Amza T, Peng W, Zhou HM (2014) Delineating the protein changes in Asian noodles induced by vacuum mixing. *Food Chem* 143:9–16
- McCormick KM, Panozzo JF, Hong SH (1991) A swelling power test for selecting potential noodle quality wheats. *Crop Pasture Sci* 42(3):317–323
- Miskelly DM, Moss HJ (1985) Flour quality requirements for chinese noodle manufacture. *J Cereal Sci* 3(4):379–387
- Morris CF, Jeffers HC, Engle DA (2000) Effect of processing, formula and measurement variables on alkaline noodle color-toward an optimized laboratory system. *Cereal Chem* 77(1):77–85
- Moss HJ, Miskelly DM, Moss R (1986) The effect of alkaline conditions on the properties of wheat flour dough and Cantonese-style noodles. *J Cereal Sci* 4(3):261–268
- Okusu H, Otsubo S, Dexter J, Hou GG (2010) Wheat milling and flour quality analysis for noodles in Japan. In: Hou GG (ed) Asian noodles: science, technology, and processing. Wiley, New Jersey, pp 57–73
- Ong YL, Ross AS, Engle DA (2010) Glutenin macropolymer in salted and alkaline noodle doughs. *Cereal Chemistry* 87(1):79–85
- Park CS, Baik BK (2004) Relationship between protein characteristics and instant noodle making quality of wheat flour. *Cereal Chem* 81:159–164

- Park CS, Hong BH, Baik BK (2003) Protein quality of wheat desirable for making fresh white salted noodles and its influences on processing and texture of noodles. *Cereal Chem* 80(3):297–303
- Park CS, Kang CS, Jeung JU, Woo SH (2011) Influence of allelic variations in glutenin on the quality of pan bread and white salted noodles made from Korean wheat cultivars. *Euphytica* 180(2):235–250
- Rahman MS (2007) *Handbook of food preservation*, 2nd edn. CRC Press, Boca Raton
- Ramli S, Alkarkhi AF, Shin Yong Y, Min-Tze L, Easa AM (2009) Effect of banana pulp and peel flour on physicochemical properties and in vitro starch digestibility of yellow alkaline noodles. *Int J Food Sci Nutr* 60(s4):326–340
- Rayas-Duarte P, Francisco C, Payton ME, Bellmer DD, Carver BF, Huang WN (2009) Alkaline noodles and flour/gel properties of hard red and white winter wheat. *J Food Qual* 32(5):627–643
- Ross AS, Crosbie GB (2010) Effects of flour characteristics on noodle texture. In: Hou GG (ed) *Asian noodles: science, technology, and processing*. Wiley, New Jersey, pp 313–330
- Ross AS, Ohm JB (2006) Sheeting characteristics of salted and alkaline Asian noodle doughs: comparison with lubricated squeezing flow attributes. *Cereal Foods World* 51:191–196
- Ross AS, Quail KJ, Crosbie GB (1997) Physicochemical properties of Australian flours influencing the texture of yellow alkaline noodles. *Cereal Chem* 78:814–820
- Rosyid TA, Karim R, Adzahan NM, Ghazali FM (2011) Antibacterial activity of several Malaysian leaves extracts on the spoilage bacteria of yellow alkaline noodles. *Afr J Microbiol Res* 5:898–904
- Shelke K, Dick JW, Holm YR, Loo KS (1990) Chinese wet noodle formulation: a response surface methodology study. *Cereal Chem* 67(4):338–342
- Shewry PR (2009) Wheat. *J Exp Bot* 60(6):1537–53
- Takashima H, Miyakawa Y, Kanno Y (2007) Microwave sterilization with metal thin film coated catalyst in liquid phase. *Mater Sci Eng C* 27(4):898–903
- Thammathongchat S, Fukuoka M, Watanabe H (2005) An innovative noodle: gelatinized at the core, leaving the surface ungelatinized. *J Food Eng* 70(1):27–33
- Toyokawa H, Rubenthaler GL, Powers JR, Schanus EG (1989a) Japanese noodle qualities. I. Flour component. *Cereal Chem* 66:382–386
- Toyokawa H, Rubenthaler GL, Powers J. R, Schanus EG (1989b) Japanese noodle qualities. II. Starch component. *Cereal Chem* 66:387–391
- Tsimogiannis H, Mmolawa P, Davos D, Tribe IG (2001) An outbreak of *Salmonella typhimurium* phage type 29 linked to a noodle restaurant in South Australia. *Commun Dis Intell Q Report* 25(2):72–76
- Ugarcic-Hardi Z, Jukic M, Komlenic DK, Sabo M, Hardi J (2007) Quality parameters of noodles made with various supplements. *Czech J Food Sci* 25:151–157
- Veraverbeke WS, Delcour JA (2002) Wheat protein composition and properties of wheat glutenin in relation to breadmaking functionality. *Crit Rev Food Sci Nutr* 42(3):179–208
- Vijayakumar TP, Boopathy P (2012) Optimization of ingredients for noodle preparation using response surface methodology. *J Food Sci Technol* 51(8):1501–1508
- Wang YH, Lu XT, Wang JW (2008) Relationship between the wheat-powder descent amount and the damaged starch content and the bud rate. *J Chang Chun Univ Technol (Natural Science Edition)* 29:455–458
- World Instant Noodle Association Expanding Market (2013) World instant noodle association. Osaka, Japan. <http://instantnoodles.org/noodles/expanding-market.html>
- Wu J, Beta T, Corke H (2006) Effects of salt and alkaline reagents on dynamic rheological properties of raw oriental wheat noodles. *Cereal Chem* 83(2):211–217
- Xu Y, Hall C III, Wolf-Hall C, Manthey F (2008) Fungistatic activity of flaxseed in potato dextrose agar and a fresh noodle system. *Int J Food Microbiol* 121(3):262–267
- Xue C, Sakai N, Fukuoka M. (2008) Use of microwave heating to control the degree of starch gelatinization in noodles. *J Food Eng* 87(3):357–362

- Xue DAN, Gao H, Zhao LEI, Yin J (2010) Relationships between instrumental measurements and sensory evaluation in instant noodles studied by partial least squares regression. *J Texture Stud* 41(2):224–241
- Yamauchi H, Ito M, Nishio Z, Tabiki T, Kim SJ, Hashimoto N, Hideho Miura H, ISM Z (2007) Effects of high-molecular-weight glutenin subunits on the texture of yellow alkaline noodles using near-isogenic lines. *Food Sci Technol Res* 13(3):227–234
- Ye Y, Zhang Y, Yan J, Zhang Y, He Z, Huang S, Quail KJ (2009) Effects of flour extraction rate, added water, and salt on color and texture of Chinese white noodles. *Cereal Chem* 86(4):477–485
- Yeoh SY, Alkarkhi AF, Ramli SB, Easa AM (2011) Effect of cooking on physical and sensory properties of fresh yellow alkaline noodles prepared by partial substitution of wheat flour with soy protein isolate and treated with cross-linking agents. *Int J Food Sci Nutr* 62(04):410–417
- Zeb A, Zahir A, Ahmad T, Abdumanon A (2006) Physiochemical characteristics of wheat varieties growing in the same and different ecological regions of Pakistan. *Pakistan J Agric Biol Sci* 9(9):1823–1828
- Zhang SB, Lu QY, Yang H, Meng DD (2011) Effects of protein content, glutenin-to-gliadin ratio, amylose content, and starch damage on textural properties of Chinese fresh white noodles. *Cereal Chem* 88(3):296–301
- Zhao LF, Seib PA (2005) Alkaline-carbonate noodles from hard winter wheat flours varying in protein, swelling power, and polyphenol oxidase activity. *Cereal Chem* 82(5):504–516