

Index

A

- Abdomen, 32, 33, 59, 80, 168, 226, 229
- Abdominal aortic aneurysm (AAA), 227, 228, 280
- Abdominal pain, 38, 52, 60, 62, 117. *See also*
 - Acute abdominal pain (AAP)
- Abuse, 39, 40, 120, 169
- Acute abdominal pain (AAP), 62, 217
 - AA, 229
 - AAA, 227
 - age-related physiological changes, 218
 - AMI, 225
 - AP, 230
 - bile tract diseases, 230
 - differential diagnosis, 220–221
 - diverticulosis, 228–229
 - epidemiology, 218
 - extra-abdominal diseases, 231
 - history and physical examination, 218–220
 - imaging studies, 222–223
 - intestinal obstruction, 228
 - laboratory studies, 221–222
 - management, 223–224
 - NSAP, 231
 - PUD, 231
- Acute appendicitis (AA), 229
- Acute brain failure, delirium, 51
- Acute care for the elderly (ACE) model
 - units, 94, 95
 - inhibitors, 116
- Acute coronary syndrome (ACS), 52
 - diagnosis, 252
 - differential diagnosis, 253
 - NSTE-ACS, 251
 - prognosis and outcomes, 255
 - STEMI, 251
 - symptoms, 252
 - treatment, 253–254
 - cardiac catheterization, 254
 - pharmacological treatment, 254
 - prasugrel, 255
 - ticagrelor, 255
- Acute geriatric units, 9
- Acute kidney injury (AKI), 302
- Acute mesenteric ischemia (AMI), 225–227
- Acute myocardial infarction, 231
- Acute pain, 7, 278–280, 315, 316, 319, 321, 322
- Acute pancreatitis (AP), 230–231
- Acute respiratory failure (ARF), 34, 261, 262
- Adjuvant treatments, 283
- Advance care planning (ACP), 374, 383
- Advanced idiopathic skeletal hyperostosis, 167
- Advance directives (ADs), 374, 375
- Adverse drug events (ADEs), 54, 76, 101, 120, 299, 306, 307
- Ageing/aging, 24, 180, 292, 293, 300, 301
- Age-related degeneration, 148
- Airway, 164
 - assessment, 26–27
 - management, 28–31
 - protective reflexes, 59
- Airway, breathing, circulation (ABC)
 - approach, 8, 24, 40, 56
- Amiodarone, 301
- Analgesics, topical use, 321
- Antibiotics, 115, 117, 128, 183, 187, 229, 236, 237, 239, 242, 266, 270
- Anticoagulation, 270
- Antidepressants, 213–214, 283, –284
- Antihypertensives, 112, 150, 152, 305
- Apnea, 30, 32, –33
- Appendectomy, 229
- Arterial blood gas, 61, 221
- Arthritis, 31, 92, 389
- Asthma, 264, 265
- Asymptomatic bacteriuria (ASB), 115, 237
- Atrial fibrillation, 38, 154

- Autonomy, patients
 - delirium, 372
 - diagnostic/therapeutic intervention, 372
 - health care, decision-making, 371
 - informed consent, 371–373
 - and mental capacity, 371
 - paternalistic physician–patient relationship, 371
 - patient refusal, 372
 - principle of, 371
- B**
- Back pain, 59, 117. *See also* Low back pain (LBP)
- Bag-valve-mask device (BVM), 32
- Basel nonspecific complaints (BANC) study, 128
- Baseline oxygen saturation, 30
- Basic activities of daily living (BADL), 55
- Beneficence, principle of
 - ethical obligation, 379
 - health-care providers, 379
 - paternalistic, 379
 - patient confidentiality, 380
- Benzodiazepines, 143, 205, 301, 321
- Beta-2 agonist, 269
- Bile tract diseases, 230
- Bioethics, 370, 376, 388
- Biomedical ethics, 370
- Bisphosphonates, 283
- Blood pressure, 20, 32, 37, 38, 57, 111, 112, 143, 150, 152, 153, 220, 249, 294, 303
- Blood urea nitrogen, 60
- Body temperature, 113
- Boerhaave's syndrome, 29
- BONES mnemonic, 29
- Boston Syncope Criteria, 155
- Bradycardias, 149
- Breathing, 32–36, 164
 - assessment, 32
 - management
 - apnea and irregular respirations, 32–33
 - respiratory distress, 33–34
 - respiratory failure, 34–36
- Breathlessness, 261
- Brief Confusion Assessment Method (bCAM), 203
- Bronchodilators, 269–270
- B-type natriuretic peptide (BNP), 61
- C**
- Carbamazepine, 302
- Cardiac catheterization, 254
- Cardiogenic shock, 37
- Cardiopulmonary resuscitation (CPR), 37, 41–43, 384, 386, 387
- Cardiorespiratory system, age-related changes, 261, 262
- Cardiovascular disease, 149
- Cardiovascular system, 184
- Care home services, 332
- Care homes, 332
- Care transitions
 - within emergency department
 - disposition, 338–339
 - initial assessment, 336
 - optimization, 332–333
 - patient discharge, 340–341
 - pre-hospital care transitions
 - care home teams, 331
 - emergency department receiving team, 333–334
 - emergency medical services and care home, 331
 - inter-professional communication, 335–336
 - non-specific presentation, 334
 - paramedic and technician staff, 331
 - paramedics and care home, 333
 - technology, 335
 - triage, 334–335
- Carotid sinus hypersensitivity (CSH), 151
- Carotid sinus massage (CSM), 151
- Carotid sinus syndrome (CSS), 151
- Cementoplasty, 284
- Central venous oxygen saturation (ScvO₂), 186, 191
- Central venous pressure (CVP), 186
- Chest pain, 52, 247
 - ACS, 251
 - diagnosis, 252
 - differential diagnosis, 253
 - NSTE-ACS, 251
 - prognosis and outcomes, 255
 - STEMI, 251
 - symptoms, 252
 - treatment, 253–255
 - differential diagnosis, 248, 250, 251
 - PE, 248–249
 - tension pneumothorax, 249–250
 - type A aortic dissection, 249
 - management, 256, 257
- Cholecystitis, 230
- Chronic illness, 26
- Chronic pain, 276, 317, 322, 323
- Circulation
 - assessment, 36–37
 - management, 37–38

- Clarithromycin, 301
Clostridium difficile, 244
 Cognition, 101
 Cognitive impairment (CI), 20, 82, 101, 199, 200
 - delirium diagnostic criteria, 201
 - detection, 202–203
 - management, 204
 - prevention, 205
 - risk factors, 203–204
 - subtypes, 202
 dementia, 205
 - diagnosis and management, 206
 - subtypes, 205–206
 ED, 200
 epidemiology, 200–201
 neurocognitive disorders, 201
 service reconfiguration with, 206–207
 Coma, 58
 Comprehensive geriatric assessment (CGA)
 - emergency department
 - conduct, 100
 - evidence base, 95
 - evidence base, 94–95
 - homeostatic reserves assessment, 55
 - functional disability, 55
 - indicators, 55–56
 - morbid conditions, 55
 - life-threatening conditions, 50–52
 - practical examples, 99–100
 - preadmission functional and health status
 - assessment
 - biological pitfalls, 53
 - clinical pitfalls, 53
 - electrocardiogram, 53–54
 - medication history, 54
 - secondary assessment, 56
 - abdominal symptoms, 62
 - care with cognitive impairment, 62–63
 - context-dependent testings, 60–61
 - electrocardiogram, 60, 61
 - geriatric emergency health-care centers, 63–64
 - head to toe examination, 58–59
 - liaison service, 63
 - systematically ordered, 60
 - trauma, 61–62
 - vital signs and general appearance, 57–58- Computed tomography (CT), 117, 222
 - dyspnoea, 267
- Computed tomography pulmonary angiography (CTPA), 248
- Computerised physician order entry (CPOE)
 - systems, 310
- COPD exacerbations, 263, 265
- Coronary artery disease, 119
- C-reactive protein, 60
- Creatine kinase, 60
- Crystalloid fluid resuscitation, 8
- Cumulative Illness Rating Scale (CIRS), 59
- Cytochrome p450 enzyme, 301, 302

D

 - D-dimer, 61, 266
 - Delirium, 51, 82, 118, 199
 - diagnostic criteria, 201
 - detection, 202
 - management, 204
 - prevention, 205
 - risk factors, 203
 - subtypes, 202
 - Dementia, 82, 118, 199, 205, 318, 388, 389
 - diagnosis and management, 206
 - palliative and end of life care
 - communications planning, 358
 - decision making, 357
 - families and carers involvement, 359
 - Gold Standards Framework, 355–357
 - maintaining dignity, 359
 - management, 358
 - patient transfer, 357–358
 - subtypes, 205
 - Depression, 58, 101
 - antidepressant treatment, 213–214
 - diagnosis, acute and emergency care
 - setting, 210–211
 - epidemiology, 209–210
 - risk assessment, 211–212
 - suicide, factors, 212
 - symptoms, 211
 - therapeutic intervention, 212–213
 - Disposition, syncope, 156, 157
 - Distributive justice, 381
 - Distributive/vasodilatory shock, 37
 - Diuretics, 143, 269
 - Diverticulosis, 228, 229
 - Do not attempt resuscitation (DNAR), 386, 387, 391
 - Domino effect, 50–51
 - Drug-disease interactions, 304
 - Drug-drug interactions, 304
 - Dyspepsia, 52
 - Dyspnoea, 261–262
 - aetiology, 262, 263
 - diagnosis, 263–264
 - arterial blood gas, 267
 - biomarkers, 266–267
 - chest radiography, 265
 - CT, 267

- Dyspnoea (*cont.*)
 electrocardiogram, 265
 laboratory tests, 265
 symptoms and signs, 264–265
 ultrasound, 267
 risk stratification, 271
 treatment, 268
 algorithm, 268, 269
 antibiotics, 270
 anticoagulation, 270
 bronchodilators, 269
 comorbidities, management, 271
 diuretics, 269
 oxygen and NIV, 268
- Dysrhythmias, 38, 149
- E**
- Early goal-directed therapy (EGDT), 186
 Echocardiography, 37
 Edinburgh Postnatal Depression Scale, 210
 Effective communication, 78, 373
 EGSYS, 156
 Electrocardiogram, 54, 265
 Electroconvulsive therapy (ECT), 214
 Emergency department (ED), 9
 atypical presentations
 abdominal pain, 117
 abuse and neglect, 120
 ADEs, 120
 back pain, 117
 common fluid and electrolyte problems, 116
 coronary artery disease, 119
 delirium and dementia, 118–119
 discharge planning, 121
 falls, 117–118
 fever and common infections, 114
 hidden diseases, social admissions and search for, 121
 PE, 116
 pneumonia, 114–115
 thyroid disease, 116
 UTI and ASB, 115
 care plan, 103
 care transitions within
 disposition, 338
 initial assessment, 336
 team members, handover, 336–338
- CGA
 conduct, 100
 evidence base, 95
 cognition, 101
 conduct, 100
 problem detection
 assessment, domains, 100
 high-risk population, identification, 100
 depression, 101
 early discharge planning, 102
 evidence-based solutions
 care of older people, 92
 follow-up, 93
 frail older people, 93
 history of, 93
 interdisciplinary diagnostic process, 92
 multidimensional assessment, 92
 treatment, coordinated and integrated plan, 93
- falls, 102
 follow-up, 103
 frail older people in, 91
 functionality, 101
 polypharmacy, 102
 receiving team, 333
 triage
 issues, 18
 priorities, determination, 17
 process, 17
 triage environment, 18–20
 vital signs
 blood pressure, 112
 body temperature, 113
 HR, 112
 RR, 113
- Emergency medical services, 331–332
 Emergency physicians (EPs), 24, 369, 379, 386
 Emergency Severity Index (ESI), 19
 Empirical antibiotic therapy, 187, 188
 End of life care, 362–365
 dying people, priorities, 362
 individual care plans, 365
 involvement, 364
 providing support, 364
 recognising, 363
 sensitive communication, 363
 eating and drinking, 366
 environment, 365
 hospice care, 366
 organ donation, 366
 pain relief and symptom control, 365
 religion and spiritual care, 366
- End-tidal carbon dioxide concentration (ETCO₂), 8
Enterococcus faecalis, 239
Escherichia coli, 239
 Ethics, medical practice, 384, 389
 clinical decision, 370
 clinical trial, 389

- cultural and political implications, 370
 - definition, 370
 - end of life decisions, 382, 383
 - human culture diversity, 390
 - informed consent, 389, 390
 - psychological and social vulnerability, 370
 - research, emergencies, 390
 - resuscitation process, 384
 - biological age, 384
 - informed consent, 389
 - neurological outcome, 384
 - survival rates, 384
 - survival to discharge, 384
 - vulnerable populations, 390
 - Evidence-based medicine, 291–292
 - Exercise therapy, 285
 - Exposure, 165–166
 - Extra-abdominal diseases, 231
 - Extremities, 168
- F**
- 4 As Test (4AT), 202
 - Faintness, 51
 - Fall, 85, 102, 117, 137, 303
 - environment, 143
 - epidemiology, 138
 - management, 141–143
 - medication, 143
 - multidisciplinary and holistic approach, 137
 - multifactorial intervention, 143
 - multi-factorial management programme, 144
 - poor processes and lack of access, 144
 - risk factors, 138–141
 - Family witnessed resuscitation, 384–386
 - Fecal impaction, 52
 - Fever
 - and common infections, 114
 - definition, 132
 - Foreign body aspiration, 31
 - Four-item geriatric depression scale, 214
 - Frailty, 77, 78, 165, 293
 - Functional decline, 9, 51
 - Functional disability, 55
 - Futility in medicine, 376–377
- G**
- Gamma-glutamyl transferase (GGT), 60
 - Gangrenous cholecystitis, 230
 - Geriatric emergency health-care centers, 63
 - Geriatric evaluation and management (GEM) units, 94
 - Glasgow Coma Scale (GCS), 25, 165
 - Glomerular filtration rate (GFR), 302
 - Glycemic control, 188
 - Grab sheet, 332
- H**
- Head computed tomography scan, 154
 - Head examination, 58
 - Head injury, 61, 166
 - Head-tilt-chin-lift maneuver, 29
 - Heart rate (HR), 112, 133
 - Helicobacter pylori* (*H. pylori*), 231
 - High-sensitivity troponin, 61
 - Hip fracture, 50, 61, 62, 117, 143, 204
 - Homeostatic reserves assessment
 - functional disability, 55
 - morbid conditions, 55
 - Horton's disease, 279
 - Hospice care, 366
 - Hospital Anxiety and Depression (HAD) scale, 210–211
 - Hyperglycemia, 188
 - Hyperkalemia, 116
 - Hypertension, 31, 111, 227
 - Hypoactive delirium, 200
 - Hypoglycemia, 188
 - Hyponatraemia, 116
 - Hypotension, 37
 - Hypothermia, 41, 183
 - Hypovolemic shock, 37, 220
 - Hypoxemia, 34, 35
- I**
- Identification of seniors at risk (ISAR), 86
 - IL-6, 181
 - Impaired mental status, 36
 - Inappropriate prescribing (IP), 306–307
 - Infections
 - clinical manifestation, 132
 - examination, 132
 - fever, 132
 - pulmonary infection, 132
 - urinary tract infections, 132
 - working hypothesis, 132
 - Inflammatory low back pain, 279
 - Instrumental activities of daily living, 55
 - Interdisciplinary geriatric consultation services (IGCS), 94
 - Intestinal obstruction, 228
 - Intoxication, 40–41
 - Irregular respirations, 32

J

- Jaw-thrust maneuver, 29
- Justice, principle of, 381
 - fairness, opportunity, 381
 - finite resources, equitable allocation, 381, 382

K

- Klebsiella* sp., 237, 239

L

- Laxatives, 283
- LEMON mnemonic, 29
- Lewy bodies (LBD), 206
- Liaison service, 63
- Lidocaine, 323
- Life-threatening conditions, 23
 - acute brain failure
 - delirium, 51
 - faintness/syncope and general malaise, 51
 - atypical presentations, 52
 - digestive symptoms, 52
 - domino effect, 50
 - functional decline, 51
 - referred pain and symptoms, 52
 - secondary assessment, CGA, 56
 - abdominal symptoms, 62
 - care with cognitive impairment, 62
 - context-dependent testings, 61
 - electrocardiogram, 60, 61
 - head to toe examination, 58
 - systematically ordered, 60
 - trauma, 61
 - vital signs and general appearance, 57
- Life-threatening emergencies, 25–26
- Low back pain (LBP), 275–276
 - autonomy and functional performance, 282
 - clinical presentation, 277
 - evaluating pain, 280–281
 - history, 277–278
 - inflammatory, 278–279
 - lumbar spinal canal stenosis, 279–280
 - older patients, behavior change, 280
 - osteoarthritis, 280
 - osteoporotic fracture, 279
 - physical examination, 278
 - epidemiology, 276
 - investigations, 281–282
 - management, 285
 - pathophysiology, 276–277
 - treatment, 282

- adjuvant treatments, 283–284
- exercise therapy, 285
- nonopioid analgesics, 282–283
- opioids, 283
- surgical treatment, 284

- Lower limbs, 59
- Lumbar degenerative spondylolisthesis, 283
- Lumbar puncture, 61
- Lumbar spinal canal stenosis, 279

M

- Magnetic resonance imaging (MRI), 168, 229, 281
- Manchester Triage System (MTS), 19, 79
- Mean arterial blood pressure, 37
- Medicine
 - ageing process, 292
 - clinical guidelines, 291
 - Evidence-based medicine, 291
 - older patients, 293–295
 - physician knowledge and experience, 296–297
 - reconciliation, 308
 - scientific evidence, 295
- Melancholic somatic symptoms, 210
- Mental health disorders (MHDs), 209
- Microstomia, 31
- Mild cognitive impairment (MCI), 205
- Morphine, 283
- Multidrug-resistant organisms (MDROs), 239

N

- Natriuretic peptide type B, 266
- Neck examination, 59
- Neck stiffness, 54
- Neglect, 120, 169
- Neurologic disorder, 36
- Neuropathic pain, 317, 323
- Nitrates, 143, 150, 242
- Nitrous oxide-oxygen gas mixture (N₂O/O₂), 321
- Nociceptive pain, 317
- Noninvasive positive-pressure ventilation (NIPPV), 33, 34
- Non-invasive ventilation (NIV), 268
- Non-maleficence principle, 375, 376
 - futile medical treatment, 376, 377
 - implications, 376
 - life-prolonging treatment, 378, 379
- Non-occlusive mesenteric ischemia (NOMI), 225, 227
- Nonopioid analgesics, 282, 283

- Non-prescription, drug use, 300
 Nonspecific abdominal pain (NSAP), 231
 Nonspecific complaints (NSCs), older patients
 adverse outcomes, 127, 128
 BANC, 128
 causes, 128–129
 diagnostic approach
 diagnostic workup, 127, 129, 130
 heart failure, 133
 infections, 132
 localised vs. generalised weakness, 130
 medical history, 130–131
 physical examination, 130–131
 specific symptoms, 129
 water and electrolyte disorder, 132–133
 disease presentation, 128
 emergency department, 127
 in-hospital mortality, 127, 128
 lack of community support, 128
 Non-steroidal anti-inflammatory drugs
 (NSAIDs), 282, 320
 NSTE-ACS, 251, 254, 255
- O**
- Obstructive shock, 37
 Older abuse, 387, 388
 Older patients
 care, models of, 8–9
 acute geriatric units, 9
 community and primary care, 9–10
 ED, 9
 EMS/EMT/ambulance care, 11
 hospital at home, 10–11
 out of hours' care, 10
 traditional inpatient management
 models, 9
 impact on services, 4
 pathologies
 pain, 7
 recognition, 6
 time critical, 6–7
 pre-hospital geriatric patient
 anatomical and physiological differ-
 ences, 5
 disability and function, 5
 epidemiology, 6
 pre-hospital personnel training, 4–5
 Older people, 75–86
 communication, with frailty, 82
 functional assessments, 87–88
 geriatric giants
 dementia and delirium, 82–83
 falls and syncope, 85
 injuries, 85
 medication, 86
 pain, 85–86
 sepsis, 83–84
 management
 CGA, 77
 differential challenge, 76–77
 functional decline and altered
 homeostasis, 76
 multiple comorbidities, 76
 non-specific presentations, 75
 older abuse, assessing for, 87
 risk assessment, 86–87
 urgent care, identifying with frailty, 77
 carers and families, communication
 with patients, 78–79
 history taking, 79
 physical examination, 80
 professionals, communication between,
 80–81
 Opioids, 283, 320, 321
 Oral transmucosal fentanyl citrate (OFTC),
 321
 Organ donation, 366
 Orthostatic hypotension (OH), 150, 151
 Osservatorio Epidemiologico sulla Sincope nel
 Lazio (OESIL) score, 155
 Osteoarthritis, 277, 280
 Osteoarticular pain, 275
 Osteoporotic fracture, 279, 283
 Oxygen supplementation, 268
 Oxygen therapy, 33, 268
 Oxygenation, 32, 267
- P**
- Pain, 7, 85, 315, 316, 319–322
 assessment, 317–318
 behaviours, 319
 care quality indicators, 316
 chronic pain, exacerbation, 322
 classification, 317
 definition and types, 316–317
 dementia and assessment, 318–319
 drugs, 323
 management, 322
 analgesics, topical use, 321
 interventional techniques, 322
 N₂O/O₂, 321
 principles, 319–320
 treatment, 320–321
 neuropathic, 323
 optimal management, 315
 under- treatment of, 316

- Painful diverticular disease, 229
- Palliative care, 362–365
 - dying people, priorities, 362–363
 - individual care plans, 364–365
 - involvement, 364
 - providing support, 364
 - recognising, 363
 - sensitive communication, 363–364
 - eating and drinking, 366
 - environment, 365
 - hospice care, 366
 - organ donation, 366
 - pain relief and symptom control, 365
 - religion and spiritual care, 366
- Paracetamol, 320
- Patient communication, 373–375
- Patient response/mental status assessment, 25
- Patient's comorbidity, 59
- Patient-centred care, 95
- Pelvis, 168
- Peptic ulcer disease (PUD), 231
- Percutaneous epidural adhesiolysis, 284
- Permissive hypotension, 8
- Phenytoin, 301, 302
- Plain abdominal X-rays, 222
- Pneumatosis intestinalis, 227
- Pneumonia, 114, 180
- Polypharmacy, 76, 101, 102, 304
- Postural homeostasis, 150
- Potentially inappropriate medication (PIM), 306
- Prasugrel, 255
- Prescription
 - adverse drug events, 307–308
 - CGA, 309
 - CPOE, 310
 - drug-drug and drug-disease interactions, 304–306
 - epidemiology of, 300
 - inappropriate prescribing, 306
 - medicines reconciliation (MR), 308–309
 - pharmacokinetics and pharmacodynamics, 300–303
 - recommendations, 309
- Primary care, communicating with, 341
- Procalcitonin (PCT), 61, 184, 266
- Procedural pain, 317
- Prodromes, 150, 154
- Pulmonary and heart examination, 59
- Pulmonary embolism (PE), 116, 248
- Pulmonary Embolism Severity Index (PESI), 271
- Pulmonary infection, 132
- Pyuria, 240
- Q**
- Quality emergency care, 391
- quickSOFA (qSOFA), 178, 179
- R**
- Rapid sequence intubation (RSI), 30
- Rectal examination, 59
- Referred pain, 52
- Rehabilitation, 345
 - collaborative inter-disciplinary working, 350
 - conceptual framework, 350
 - geriatric emergency medicine, 347–349
 - ICF, 346
 - inter-disciplinary working, 349–350
 - interpersonal and communication skills, 351
 - vulnerability, to poorer outcomes, 347
 - WHODAS 2.0, 346–347
- Religion care, 366
- Reperfusion, 226
- Respiratory distress, 33
- Respiratory failure, 34, 262
- Respiratory rate (RR), 113
- Resuscitation, 41–43, 164
- Richmond Agitation Sedation Scale (RASS), 25, 58
- Rockwood Frailty Index, 55
- S**
- San Francisco Syncope Rule (SFSR), 155
- Scientific evidence, 295–296
- Senescent immune remodeling, 180
- Sensitive communication, 363
- Sensory exam, 59
- Sepsis, 83, 152, 177, 186–189
 - causes, 180
 - clinical presentation and diagnosis, 182–185
 - epidemiology of, 179–180
 - management, 184–185
 - glycemic control, 188
 - initial resuscitation, 186–187
 - principles, 188–189
 - source control and antimicrobial treatment, 187–188
 - treatments, 188
 - pathophysiology, 180–181
 - prognosis and outcomes, 189–190
 - risk factors, 181–182
- Septic shock, 177, 178, 184, 186–188
- Sequential (sepsis-related) organ failure assessment (SOFA) score, 178

- Serum creatinine, 53, 60
 Serum electrolytes, 60
 Severe low back pain, 59
 Silver Code (SC), 87
 Single Question in Delirium (SQiD), 202
 SIRS, 179, 191
 Social admissions, 82, 121
 Spinal stenosis, 284
 Spine fusion, 284
 Spiritual care, 366
 Spondylodiscitis, 279
 Standard medical therapy (SMT), 34
 Standardised handover tool, 332
Staphylococcus, 239
 ST-elevation myocardial infarction (STEMI), 251
 Steroids, 283
 Stick blood glucose, 60
 Superior mesenteric artery (SMA), 225
 Surviving Sepsis Campaign (SSC), 184
 Syncope, 51, 85, 147
 background and epidemiology, 148–149
 disposition, 156–157
 etiologies, 149–152
 evaluation, 152–153
 multidimensional geriatric assessment, 148
 scoring tools, 154–156
 testing, 153–154
 Syncope units (SUs), 156
- T**
 Tachycardias, 149
 Tension pneumothorax, 249, 250
 Thorax, 167
 Thyroid disease, 116
 Thyroid-stimulating hormone, 61
 Ticagrelor, 254, 255
 Traditional inpatient management models, 9
 Tramadol, 283
 Transaminases, 60
 Trauma, 38, 39, 61, 163
 abdomen, 168
 abuse and neglect, 169
 airway and breathing, 164–165
 clinical assessment, 166
 disability, 165
 exposure, 165–166
 extremities, 168–169
 head injury, 166–167
 imaging, 166
 neck and spine, 167
 pelvis, 168
 resuscitation, 164
 thorax, 167
 Trauma Audit Research Network (TARN), 143
 Triage, ED
 issues, 18
 priorities, determination, 17
 process, 17
 triage environment, 18–20
 Troponin, 54, 251, 266
 Type A aortic dissection, 249
- U**
 Ultrasonography (USG), 222, 228, 229
 Ultrasound, dyspnoea, 267
 Urinalysis, 61, 132, 222
 Urinary tract infection (UTI), 115, 132
 diagnosis
 community-dwelling older adults, 240–242
 long-term care facilities, diagnostic algorithm, 243
 long-term indwelling urinary catheter, 243–244
 patients with communication barriers, 241–242
 microbiology, 239–236
 prevention, 244–245
 risk factors, 239–240
 symptoms, 236–237
 asymptomatic bacteriuria, 237–238
 symptomatic, 238–239
 treatment, 243–244
- V**
 Vasovagal syncope, 148, 150
- W**
 Warfarin, 120, 143, 301, 303, 307
 Water and electrolyte disorder, –133, 132
 World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0), 346
- X**
 X-rays, 132, 166, 204, 222, 227, 250, 264, 281