Index

A	fluoride ion release, 99–100
Abrasion, 37–38	intra-orifice barrier and restorative
Acidic polymer, 13–14	material, 105-106
Adhesion	nonsurgical applications, 104
delivery method, 38–40	perforation repair, 106–108
early research, 18–20	root canal sealers, 104-105
Aluminosilicate poly(acrylic acid) (ASPA), 9	root-end filling material, 106
Amorphous calcium phosphate (ACP), 141	surgical applications, 104
Atraumatic restorative treatment (ART)	Controlled-release polymer technologies, 129
and clinical performance, 48–50	
GIC and, 89–90	
restorative care, 116	D
sealant, 50, 51	Dental silicate cement, 3–4
Attenuated total reflectance (ATR), 15	composition, 4
Auto-cure glass-ionomer cement	physical properties, 6
bonding amalgam to enamel and dentine, 77–78	setting reaction, 4–6
carrier for medicaments, 75–77	solubility and ion release, 6–7
cervical hypersensitivity management, 63-64	
co-curing, 73–74	
collapsing dentition, 72	E
dental adhesives, 73	Electron probe microanalysis (EPMA), 5, 6, 58
fissure protection, 62–63	Endodontic care, 120
luting cements, 64	Erosion, 36–37
OII-type cavity, 65–66	
OI-type cavity, 64–65	T.
PI-type cavity, 67–69	F
resin-modified, 72–73	Fluoride release
silver-sintered auto-cure, 71–72	clinical benefits, 58
tooth structure, 75	early research, 17
unsupported cusps, 69–71	endodontics, properties, 99–100
	pharmacological approaches, 140–141
D	Fracture toughness, 35, 132–133
Biofilm formation 44, 141, 142	
Biofilm formation, 44, 141–142 Bioremineralisation, 141	G
Bioteninieransation, 141	_
	Gelatin hydrogels, 130 G200 glass, 9–10, 12
C	Glass-Ionomer cements (GICs), 126
Chlorhexidine, 142, 143	adhesion, vs. resin composite, 92–93
Clicker system, 29, 32	aesthetic improvements, 139, 140
Contemporary endodontics, 97	amalgam, 126–127
adhesive properties, 97–99	antibiotic additions, 143–144
antibacterial properties, 103	antimicrobial properties, 142–143
biocompatibility, 100–103	and ART, 89–90
	,

Glass-Ionomer cements (GICs) (cont.)	root canal sealers, 104–105
auto-cure glass-ionomer cement	root-end filling material, 106
bonding amalgam to enamel and dentine, 77–78	surgical applications, 104
carrier for medicaments, 75–77	controlled-release vehicle, 129–130
cervical hypersensitivity management, 63-64	delivery method
co-curing, 73–74	abrasion, 37–38
collapsing dentition, 72	adhesion, 38–40
dental adhesives, 73	capsules types, 29, 30
fissure protection, 62–63	clicker system, 29, 32
luting cements, 64	erosion, 36–37
OII-type cavity, 65–66	fracture toughness, 35
OI-type cavity, 64–65	glass, 31–33
PI-type cavity, 67–69	ion release (see Ion release)
resin-modified, 72–73	liquid, 33
silver-sintered auto-cure, 71–72	paste/paste dispenser, 29, 31
tooth structure, 75	powder/liquid RM-GIC lining material, 28-29
unsupported cusps, 69-71	shear punch strength, 35–36
auto-cure/resin-modified, 62	strength, 33–35
biofilm alteration, 141–142	types, 29, 31
bio-protection, 142–143	delivery systems, 136
bioremineralisation/ biopromoting	dental silicate cement, 3–4
improvements, 141	composition, 4
clinical applications	physical properties, 6
condition/etch, 62	setting reaction, 4–6
surface preparation, 62	solubility and ion release, 6–7
clinical benefits	dentine and enamel, adhesion, 133–136
adhesion, 58	early research
aesthetics, 60	acidic polymer component, 13–14
contouring, 60	adhesion, 18–20
fluoride release, 58, 59	clinical applications, 20–21
gingival biocompatibility, 60	composition and structure of glasses, 11–13
ion exchange layer and fluorapatite	fluoride release, 17
formation, 58, 59	maturation and water role, 16–17
marginal caries protection, 60	physical properties, 17–18
remineralization, 58, 60	tartaric acid, 14–16
clinical performance	fibre reinforcement, 128–129
ART, 48–50	filler systems, improvements, 127
coating, 48	fluoride release, 140–141
NCCL, 45–47	fracture toughness, improvements, 132–133
reinforced, 48	invention, 9–10
restorations, 47–48	limitations
sealants, 50–51	buffering oral acids, 61
clinical studies of, 87–89	residual HEMA, 61–62
composition and classification	wear resistance, 61
lining/base cements, 27–28	MID, 81–82
luting and bonding (RM-GIC adhesive)	minimally invasive dentistry, 82–83
materials, 26–27	minimum intervention care, 81
restorative materials, 27	multi-arm poly(acrylic acid-co-itaconic) acids,
contemporary endodontics, 97	131–132
adhesive properties, 97–99	nanoparticle technology, 127–128
antibacterial properties, 103	pharmacological approaches, participation, 144
biocompatibility, 100–103	pioneering studies, 10
fluoride ion release, 99–100	pit and fissure sealing, 144
intra-orifice barrier and restorative	polymeric filler carriers, incorporation, 129
material, 105–106	porosity reduction, 132
nonsurgical applications, 104	and pulp response, 92
perforation repair, 106–108	and remineralization, 86–87

restorations, longevity, 90–91	0
self-cleaning glass technology, 142	OII-type cavity, 65–66
self-healing technology, 130–132	OI-type cavity, 64–65
spherical particles, 128	Open-sandwich restorations, 70–71
tooth structure	Orthodontic care, 120–121
caries-affected dentine, 85–86	
enamel, 83	
fluoride and mineralized tissues, 83-85	P
sound dentine, 83–85	Paediatric dentistry
wear improvements, 136–139	approximal surfaces, effects, 116
zinc polycarboxylate cements, 7–9	contemporary carious tissue removal
Gold nanoparticles, 130	cavity design, 117, 119
Gord nanoparticles, 130	clinical scenarios, 119–120
	placement, restoration, 118
H	sandwich technique, 117
High-viscosity glass-ionomer cement (HVGIC), 89, 90	treatment modality, 116
Hydrogel technology, 129–130	endodontic care, 120
2-Hydroxyethylmethacrylate (HEMA), 61–62	orthodontic care, 120–121
2-Hydroxyethyllilethacrylate (HEMA), 01–02	•
	parental concerns, 121
т	prevention, caries, 114–116
I	restorative care, 116
Ion release	Paste/paste dispenser, 29, 31
advantage of, 41	PI-type cavity, 67–69
biofilm formation, 44	Polyacrylic acid (PAA), 13, 83
fluoride release and, 41–43	Polymer carriers, 129
solubility and, 6–7	Porosity reduction, 132
т	R
\mathbf{L}	
L Liquid, 33	
Liquid, 33	Remineralization, 86–87
Liquid, 33	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62
Liquid, 33 Luting cements, 64 M	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62
Liquid, 33 Luting cements, 64 M	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID))	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85 Minimum intervention dentistry (MID), 81–82	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T Tartaric acid, 14–16
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85 Minimum intervention dentistry (MID), 81–82 minimally invasive dentistry, 82–83	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T Tartaric acid, 14–16 Titanium dioxide (TiO ₂), 142
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85 Minimum intervention dentistry (MID), 81–82 minimally invasive dentistry, 82–83 minimum intervention care, 81	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T Tartaric acid, 14–16 Titanium dioxide (TiO ₂), 142
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85 Minimum intervention dentistry (MID), 81–82 minimally invasive dentistry, 82–83 minimum intervention care, 81	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T Tartaric acid, 14–16 Titanium dioxide (TiO ₂), 142
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85 Minimum intervention dentistry (MID), 81–82 minimally invasive dentistry, 82–83 minimum intervention care, 81	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T Tartaric acid, 14–16 Titanium dioxide (TiO ₂), 142 Z Zinc oxychloride cement, 2
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85 Minimum intervention dentistry (MID), 81–82 minimally invasive dentistry, 82–83 minimum intervention care, 81 N Nano-ionomer, 28 Nanoparticle technology, 127–128	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T Tartaric acid, 14–16 Titanium dioxide (TiO ₂), 142
Liquid, 33 Luting cements, 64 M Minimally invasive (MI) dentistry, 82–83 Minimum intervention (MI) caries management adhesion, vs. resin composite, 92–93 and ART, 89–90 caries-affected dentine, 85–86 clinical studies of, 87–89 enamel, 83 fluoride and mineralized tissues, 83–85 MID (see Minimum intervention dentistry (MID)) and pulp response, 92 and remineralization, 86–87 restorations, longevity, 90–91 sound dentine, 83–85 Minimum intervention dentistry (MID), 81–82 minimally invasive dentistry, 82–83 minimum intervention care, 81	Remineralization, 86–87 Residual HEMA (2-Hydroxyethylmethacrylate), 61–62 Resin-modified glass-ionomer (RMGIC), 26–27, 61–62 S Sealants, 50–51 Self-cleaning glass technology, 142 Self-healing technology, 130–132 Shear punch strength, 35–36 Single-use dual-barrel paste-paste delivery system, 136, 137 Single-use powder-liquid delivery system, 136, 137 Spherical particles, 128 T Tartaric acid, 14–16 Titanium dioxide (TiO ₂), 142 Z Zinc oxychloride cement, 2