

# Index

- ADI. *See* alternating directions implicit
- adjoint
  - differential-algebraic equation, 273
  - equation, 270
- alternating directions implicit, 134
- approximation
  - $\mathcal{H}_2$ -optimal, 132
  - interpolation-based, 129
  - Padé, 129
- balanced truncation
  - bounded real, 122
  - LQG, 128
  - Lyapunov, 118
  - positive real, 121
  - stochastic, 126
- BDF, 87
  - method, 3, 243
- behavior, 166
  - distributional, 167
  - ITP, 168
  - weak, 166
- behaviorally
  - controllable (*see* controllable)
  - detectable (*see* detectable)
  - observable (*see* observable)
  - stable (*see* stable)
- bounded real, 122
- canonical form, 184
  - feedback (*see* feedback canonical form)
  - kronecker (*see* Kronecker canonical form)
  - Weierstrass (*see* Weierstrass canonical form)
- characteristic values
  - bounded real, 122
  - LQG, 127
  - positive real, 120
  - stochastic, 125
- Clohessy-Wilshire equations, 253
- closed kinematic loop, 10
- completely
  - controllable (*see* controllable)
  - detectable (*see* detectable)
  - observable (*see* observable)
- condition
  - jump, 278, 282
  - real-time, 264
  - Signorini contact, 285
  - switching, 278, 283
- constraint
  - at acceleration level, 11, 227
  - hidden, 49, 71
  - holonomic, 44
  - inconsistent, 59
  - Jacobian, 10
  - manifold, 21
  - matrix, 45, 71
    - rank-deficient, 56, 59
  - at position level, 9, 227
  - redundant, 59
  - rheonomic, 51
  - scleronomic, 49
  - at velocity level, 227
- contractive, 121
- controllable
  - behaviorally, 194
  - in the behavioral sense, 111
  - C-, 111

- completely, 111, 194
- I-, 111
- impulse, 111, 194
- Inf-, 111
- at infinity, 111, 194
- R-, 111, 179
- S-, 111
- strongly, 111, 194
- coordinate
  - minimal, 10
  - partitioning, 53
  - position, 43
  - velocity, 43
- co-simulation, 256
  
- DAE. *See* differential-algebraic equation
- DASSL, 80, 90
- derivative array, 25, 49, 239
- detectable
  - behaviorally, 207
  - in the behavioral sense, 111
  - completely, 207
  - R-, 111
  - RS behaviorally, 207
  - RS completely, 207
  - RS strongly, 207
  - S-, 111
  - strongly, 111, 207
- differential-algebraic equation
  - adjoint, 273
  - asymptotically stable, 110
  - Hessenberg, 226
  - implicit, 4, 5, 15
  - index, 110
  - linear, 108
    - constant coefficient, 6
    - implicit, 4
  - overdetermined, 233
  - parametric, 239, 282
  - quasilinear, 222
  - Riccati, 18
  - second-order, 146
  - semi-explicit, 20, 23, 139, 222, 224
  - stochastic, 34
  - Stokes-like, 145
- differential equation on manifold, 20
- distributional
  - evaluation, 167
  - restriction, 167
- distributions, piecewise smooth, 167
- Drazin inverse, 24
- drift off effect, 230, 266
  
- duality, 182
  - of detectability and stabilizability, 208
  - of observability and controllability, 191
  
- eigenvalue
  - finite, 110
  - generalized, 110
  - at infinity, 110
- Euler–Lagrange equations, 10
  
- feedback canonical form, 164
- force vector, 45, 75
  - condensed, 79
- friction cone, 285
- functional mock-up interface, 256
  
- Gauss-Seidel co-simulation scheme, 258
- generalized- $\alpha$  method, 82
- Gramian
  - bounded real, 122
  - controllability, 115
  - improper, 116
  - LQG, 127
  - observability, 115
  - positive real, 120
  - proper, 116
  - stochastic, 125
  
- Hamilton’s principle, 44, 64, 72
- Hankel singular value, 115
  - improper, 117
  - proper, 117
- Hautus test, 195, 208
- HEDOP5, 94
- Hessenberg
  - differential-algebraic equation (*see* differential-algebraic equation)
- HHT- $\alpha$  method, 82
- hidden constraints. *See* constraints
  
- ILTIS vehicle, 252
- impact model, 286
- impulse
  - controllable (*see* controllable)
  - observable (*see* observable)
- impulsive part, 168
- increment function, 246
- IND. *See* internal numerical differentiation

- index
  - body mass, 75
  - differential, 5, 28
  - 3 differential-algebraic equation in
    - Hessenberg form, 54, 60, 81
  - differentiation, 224
  - 1 formulation, 94
  - 2 formulation, 91
  - 3 formulation, 90
  - of a matrix pencil, 210
  - nilpotency, 7
  - perturbation, 15, 228
  - reduction, 90, 210
  - tractability, 20
- initial trajectory problem, 168
- initial value
  - consistent, 8, 53, 168, 236, 237
  - inconsistent, 168
  - problem, 222
- input, 163
- internal numerical differentiation, 268
- interpolation
  - Hermite, 130, 131
  - tangential, 132
- ITP-behavior, 168
  
- jacobi co-simulation scheme, 258
- jump. *See* condition
  
- Kalman
  - criterion, 199
  - decomposition, 203
- kinetic energy, 11
- Kronecker canonical form, 164
- Krylov subspace
  - block, 129
  - extended, 135
  - rational, 136
  
- Lagrange multiplier, 11, 44
- Lagrangian, 44
- Lie group, 62
- LQG. *See* linear-quadratic Gaussian
- linear-quadratic Gaussian, 127
- local parametrization, 22
- loop, closed kinematical, 80
- low-rank
  - ADI, 134
  - Newton, 138
- Lur'e equation
  - projected
    - bounded real, 122
    - positive real, 120
- Lyapunov equation
  - generalized, 128
  - projected
    - continuous-time, 116, 133
    - discrete-time, 117, 132, 150
- Markov parameter, 113
- mass matrix, 10, 43, 79
  - body, 75
  - condensed, 79
  - rank-deficient, 44, 55
- matrix
  - constraint, 45, 71
    - rank-deficient, 56, 59
  - mass (*see* mass matrix)
  - pencil (*see* matrix pencil)
  - sensitivity, 267
- matrix pencil, 245
  - regular, 108, 163
  - singular, 163
  - stable, 110
- maximum dissipation principle, 287
- MDOP5, 95
- method
  - ADI (*see* alternating directions implicit)
  - alternating directions implicit, 134
  - BDF, 3, 243
  - Generalized- $\alpha$ , 82
  - HHT- $\alpha$ , 82
  - multi-step, 242
  - Newton, 137
    - low-rank, 138
    - semi-smooth, 289
  - one-step, 242
  - projection, 18
  - Rosenbrock, 23
  - Rosenbrock-Wanner, 247
  - Runge-Kutta, 16, 245
    - half-explicit, 92, 95, 250
    - implicit, 86
    - linearly-implicit, 247
    - partitioned, 251
  - Smith, 245
- mode
  - forward, 267
  - reverse, 267
- model order reduction, 108
- Moebius transformation, 123
- moment, 113, 129
  - matching, 129
  - multi-point, 131

- multibody formalisms, 72, 73
  - $\mathcal{O}(N)$ , 77
  - explicit, 80
  - recursive, 75
  - residual, 80
  
- Newton-Euler equations, 73
- Newton method. *See* method
- non-smooth equation, 289
- normal form, 184
  
- observability. *See* observable
- observable, 161
  - behaviorally, 171, 198, 201
  - in the behavioral sense, 111
  - C-, 111
  - completely, 111, 173, 198, 201
  - I-, 111
  - impulse, 111, 171, 197, 201
  - Inf-, 111
  - at infinity, 111, 173, 197, 201
  - R-, 111
  - RS behaviorally, 175
  - RS completely, 176, 202
  - RS impulse, 175, 202
  - RS at infinity, 175, 202
  - RS strongly, 175, 202
  - S-, 111
  - strongly, 111, 171, 198, 201
  - weakly behaviorally, 202
  - of zero, 172
- ODE. *See* ordinary differential equation
- OI. *See* output injection
- one-step method, 242
- ordinary differential equation, 162
  - underlying, 224
- output, 163
  - injection, 180
    - equivalence, 180
    - normal form, 182
  
- parametric
  - constrained least-squares problem, 239
  - sensitivity analysis, 238, 266
- partial
  - differential-algebraic equation, 29
  - realization, 130
- passive, 119
- PDAE. *See* partial differential-algebraic equation
  
- piecewise smooth distributions. *See* distributions
- pole
  - finite, 112
  - at infinity, 112
- Popov-Belevitch-Hautus test. *See* Hautus test
- positive real, 119
- power spectrum, 123
- principle
  - Hamilton's, 44, 64, 72
  - maximum dissipation, 287
- projection
  - Galerkin, 115
  - methods, 18
  - Petrov-Galerkin, 115
  - techniques, 97
- projector-based analysis, 19
  
- quaternions, 253
  
- RADAU5, 89–91
- R-controllable. *See* controllable
- real-time condition. *See* condition
- realization, 111
  - minimal, 111
- reflexive inverse, 124, 135
- Riccati equation
  - bounded real, 123
  - differential-algebraic, 18
  - generalized, 127
  - positive real, 121
  - projected, 137
- Rosenbrock method. *See* method
- Rosenbrock-Wanner method. *See* method
- RS
  - behaviorally detectable (*see* detectable)
  - behaviorally observable (*see* observable)
  - completely detectable (*see* detectable)
  - completely observable (*see* observable)
  - impulse observable (*see* observable)
  - observable at infinity (*see* observable)
  - strongly detectable (*see* detectable)
  - strongly observable (*see* observable)
- Runge-Kutta method. *See* method
  
- saddle point problem, transient, 30
- satellite docking maneuver, 252
- sensitivity, 282
  - DAE, 241, 271
  - matrix, 267
  - theorem, 240

- Signorini contact condition. *See* condition
- Smith method, 133
- solution
  - distributional, 167
  - weak, 166
- SO(3). *See* special orthogonal group
- special orthogonal group, 62
- spectral projector, 110
- stabilizable
  - in the behavioral sense, 111
  - R-, 111
  - S-, 111
  - strongly, 111
- stabilization, 211
  - Baumgarte, 97, 233, 266
  - Gear–Gupta–Leimkuhler, 234
- stabilized index-2
  - formulation, 99
  - system, 13
- stable, behaviorally, 210
- state, 163
  - feedback, 185
  - space form, 22
- strongly
  - controllable (*see* controllable)
  - detectable (*see* detectable)
  - observable (*see* observable)
- switching
  - condition, 278, 283
  - function, 278
- system
  - behavioral, 163
  - bilinear, 153
  - complementarity, 288
  - descriptor, 162
    - bilinear, 153
    - linear time-invariant, 108
    - periodic discrete-time, 150
  - hybrid, 278
  - mechanical multibody, 11, 227
    - with contact and friction, 285
  - pencil, 112
  - regular, 184
  - singularly perturbed, 23
  - stabilized index-2, 13
- tangent space, 21
- tilde operator, 63
- topology, 73
- transfer function, 109
  - bounded real, 122
  - improper, 112
  - minimum phase, 113
  - positive real, 119
  - proper, 112
  - strictly proper, 112
- tree structure, 73
- Weierstrass canonical form, 7, 109, 164
- Wong sequence, 200, 210
  - restricted, 201
- zero
  - finite, 112
  - at infinity, 112