
Index

A

- Acculturation, 6
 - AFD, 10
 - American vs. native Russian culture, 193
 - (see also Latino youth)
 - bullying, 39
 - cultural identification, 38, 39
 - description, 10, 38
 - East Asian youth, 117
 - emerging adulthood/transitional age, 39
 - familial acculturative stress, 39
 - immediate and extended family networks, 184
 - individuals, 38
 - Jewish culture, 184
 - ME population, 153
 - parental mental illness, 40
 - parent-child relationship, problematic, 39
 - psychological distress, 191
 - as risk factor, Latino youth, 71
 - Russian cultural norms, 185
 - separation and individuation, 192
 - stress, 38
 - suicidality and depressive symptoms, 39
 - Acculturative family distancing (AFD), 6,
 - 10, 28
 - cultural orientations, 121
 - East Asian youth, 120, 121
 - Acculturative stress, 39, 44
 - definition, 38
 - East Asian youth, 117
 - environment surrounding, 38
 - SSD and ADHD, 40
 - Adolescent suicide
 - cause of death, 1, 2
 - distress/impulsivity, 3
 - DSM-IV diagnosis, 3
 - risk factors, 4
 - suicide deaths, 2
 - on Western culture, 2
 - Adolescent Suicide Prevention Project, 231, 232
 - Adverse childhood experiences (ACEs), 209
 - African-American (AA)
 - AAPI youth, 9
 - AFD impacts, 10
 - Asian cultures, 9
 - Black population, 32
 - brutality, 33
 - child maltreatment, rate of, 40
 - child welfare and juvenile justice systems, 42
 - Chinese culture, 9
 - chronic suicidality and self-injury, case studies, 31, 32
 - church, utilization of, 41
 - cohort of urban, 40
 - collectivist orientation, 9
 - cultural mistrust, government agencies, 35
 - cultural values and attitudes, on youth suicide
 - AIAN, 8, 9
 - Black church, 8
 - help-seeking behavior, 8
 - Latino, 10, 11
 - NHW/Caucasian/European, 11, 12
 - vs. NHW youth, 8
 - racism and discrimination, 8
 - risk variables, suicidal, 7, 13
 - socioeconomic status, 8
 - description, 32
 - enculturation, 10
 - epidemiological studies, 36, 37
 - family's experience, 2
 - hanging and suffocation, rate of, 33
 - historical trauma, 34
 - interventions, 43, 44
 - LGBT youth, 40
 - mental health, 34
 - MH utilization, 10
 - protective factors, 33, 41
 - psychiatric evaluation, 42, 43
 - religion, 33
 - risk factors, adolescents, 37
 - stereotyping, 10
 - suicidal ideation, in girls, 36
 - suicide rates, 33
 - suicidology, 34
 - surveillance, 35
- Alaskan Native (AN), children and adults, 59
- American Academy of Child and Adolescent Psychiatry (AACAP), 213

- American Indian (AI)
 acute hospitalization, case studies, 51, 52
 arrangements, 61
 aware, 61
 MH treatment, 61
 perceived discrimination, 58
 psychiatric day treatment program, 60
 on reservations, 64
- American Indian Alaskan Native (AIAN)
 beliefs, 59
 cultural values and beliefs, 58
 demographics, 57
 depression/ASD, 60
 diagnosis, depression and grief reaction, 60
 DSM5 criteria, 60
 enculturation, 9
 ethnocide, 54
 historical trauma, 53
 hospitalization, 60
 intervention approaches
 aware, 61
 caring, 63
 community priorities, 62
 community-wide interventions, 65
 cultural activities, 62
 cultural continuity, 62, 63
 enculturation, 64
 evidence-based connect suicide prevention program, 64
 kueanna program, 63
 logistical and economic barriers, 61
 MH treatment, 61
 reservation, 64
 risk and protective factors, 63
 substance use/abuse, screening for, 63
 weaving, 62
 Joey's method of suicide, 58
 mental health stigma, 9
 MH services, 59
 physical resources, loss of, 55
 population groups, 52
 poverty, 57
 rate of suicidality, 9
 substance and alcohol abuse, 9
 suicidal ideation, 59
 suicide, cause of death, 53
 suicide rate, 56
 traditional religions, 9
 trauma history, 53, 54
 unemployment, 57
- American Indian Alaskan Native/Native Hawaiian Pacific Islander (AIAN/NHPI)
 cultural integrity, 55
 ethnocide, 55
 historical trauma, 56
 phases, historical unresolved grief, 55–56
 substance abuse, 57
 suicidal individuals, 57
 suicide rate, 56
 trauma history, 53, 54
- American Indian Life Skills Development Curriculum (AILSDC) program, 234
- Asian acculturation, 87, 91, 94, 95
- Asian-American/Pacific Islanders (AAPI) youth, 9
- Asian immigrants
 “aliens ineligible for citizenship”, 86
 differences, 85
 Hart-Celler Act, 86
 immigration policies, 85
 Luce-Celler Act, 86
 minority population, USA, 85 (*see also* South Asian (SA) youth)
 “model minorities”, 86
- Attachment-based family therapy (ABFT)
 description, 225
 theories of attachment, 225, 228
 traditional cultural values, minority groups, 228
- Attention deficit hyperactivity disorder (ADHD), 34, 40
- Autism spectrum disorder (ASD), 52, 60
- B**
- Beck Depression Inventory (BDI), 77
- Behavior Risk Factor Surveillance System (BRFSS), 203
- Bipolar disorder, 3
- Brief strategic family therapy (BSFT), 28, 79, 228
- Brief Symptom Inventory, 212
- Buddhism, 5
- Bullying, 39
- C**
- Cambodia, SEA, 104
- CARE, *see* Connect, Accept, Respond, Empower (CARE) model
- Caribbean
 church, utilization of, 42
 described, “black”, 32
 epidemiological studies, 36
 maternal education, 42
 psychoeducation, 35
 rate of suicide, 33
 stigma and mistrust, 35
- 1986 Chernobyl nuclear reactor, 190
- Child and adolescent psychiatrists (CAP), 229
- Child mental health services
 changes and differences, 21
 cultural competence and differences, 22
- Children of Refugees' Education, Acculturation Teaching and Empowerment (CREATE), 144
- Children's Depression Inventory (CDI), 77, 109
- Christianity, 5
- Church
 children's and youth, 32
 utilization, 41
 youth group activities, 32
- Cognitive Behavioral Intervention for Traumatic Stress (CBITS), 28

- Cognitive behavioral therapy (CBT), 44
 in adults, 160
 East Asian American youth, 128
- Cognitive Behavior Therapy for Suicide Prevention (CBT-SP), 194
- Collectivist orientation, 9
- Columbia-Suicide Severity Rating Scale (C-SSRS), 24, 25, 141
- Communalism, 6
- Community outreach
 Adolescent Suicide Prevention Project, 231, 232
 CBPR model, 231
 community-specific models, 230
 community strengths, 233–235
 description, 230
 gatekeepers, 233
 individual providers and community leaders, 230
 religious leaders, 232
 self-destructive symptoms, 231
 support, 230
- Community readiness assessment, 234, 235
- Confucianism, 116
- Connect, Accept, Respond, Empower (CARE) model, 224
- Conscientization, 63
- Crisis Safety Plan, 115
- Cultural Accommodation Model (CAM), 128
- Cultural Assessment of Risk for Suicide (CARS), 128
- Cultural competence, 22
- Cultural integrity, 55, 56
- Culturally competent mental health services
 cultural competence and differences, 22
 family therapy/involvement, 28
 practice parameter, 22, 23
- Cultural syndrome (*ataque de nervios*), 76
- Culture
 AA family, 8
 acculturation, 10
 AIAN group, 9
 Asian cultures, 9
 “coming of age” impacts, 13
 Cuban-American population, 11
 cultural cohesion, 8
 definition, 6
 families’ cultural beliefs, 16
 health beliefs, 13
 hispanic/Latino Americans, 10
 non-hispanic Whites/Caucasian/European, 11
 of suicidal behavior, 5, 6
 Western culture, 2
- Cyberbullying, 39
- D**
- Dallas Independent School District (DISD), 144
- Death, 5, 7, 8
- Department of Children and Families (DCF), 85
- Depression, 34, 38–43
 description, 175
 and emotional numbness, 175
- ME countries
 abuse history, 154
 age of girls and severity, 155
 diagnosing, 158
 gender differences, 154
 nicotine initiation, continuation, 155
 prevalence, 150, 151
 protective factors, 156
 and PTSD, in T.A. case, 161
 R-CBT, 160
 T.A. case, 152
- Diagnostic and Statistical Manual of Mental Disorders-IV (DSM -IV), 3, 36
- Dialectical behavioral therapy (DBT), 44, 78
- Discrimination, 6, 8
 acculturative stress, 39
 anti-LGBT, 40
 racism, 38
- Disenfranchised grief, 65
- Disparities
 ethnic, 12, 13
 MH, 15, 16
 treatment and services, 13, 14
- Diverse youth
 AFD, 28
 cognitive-behavioral interventions, 27
 cultural backgrounds, 22
 cultural formulation for suicidality, 24
 culturally informed treatment, 26
 disparities, 21
 evaluation of, suicidality, 23
 interview process, 24
 professionalism, 27
 suicidal ideation/motivation, 24
- DSM Cultural Formulation, 24
- E**
- East Asian youth
 angry feelings, 126
 Asian population, 115
 behavioral addiction, 127
 CARS, 128
 case study
 cognitive behavioral therapy, 115
 crisis safety plan, 115
 pro-social behavior, 114
 psychiatric evaluation, 114
 psychiatric history, 113
 suicide attempt, 114
 treatment planning, 115
- Chinese beliefs, 116
- comorbidities, 125, 126
- confucianism, 116
- cultural factors, 128
- diagnosis, 127
- DSM, 127
- GLS program, 129
- Immigration and Nationality Act of 1965, 115
- internalization, 126

- East Asian youth (*cont.*)
- intervention approaches, 128, 129
 - perceived parental support, 125
 - PIU, 127
 - proficiency, in English language, 125
 - risk factors
 - age and gender, 124, 125
 - cross-cultural, 117
 - family relationships, 119
 - intergenerational conflict, 120–122
 - internalized model minority stereotype, 123, 124
 - parent-child conflict, 118
 - parenting styles, 120
 - peer and school relationship, 118, 119
 - perceived discrimination, 122, 123
 - “psychiatric” and “non-psychiatric” subtype, 117
 - sexual orientation, 124
 - in the United States, proportion of life, 119
 - seriously considered attempting suicide, 116, 117
 - social support, 125
 - somatic symptom presentation, 126
 - somatization, 126
 - suicidal ideation, 116, 117
- Eastern European youth
- Balkans, 167
 - Belgrade government, 168
 - biases/prejudices/privileges, 177
 - “bullycide”, 175
 - constituent countries, 167
 - cultural contexts, 175
 - discriminatory stress and migratory stress, 175
 - economic stagnation, 175
 - emotional numbness, 175
 - ethnicities, 167
 - family and family-integrated models, 177
 - firearm safety, 176
 - groups and forms, 176
 - human right violations and atrocities, 168
 - mental health and illness, 171, 172
 - Multiphase Model of Psychotherapy, 177
 - NATO, 175
 - northern Baltic nations, 167
 - politicians, 168
 - primary stressors, 175
 - risk and protective factors
 - family factors, 170, 171
 - individual factors, 170
 - loss and injustice, 169, 170
 - migration, 168, 169
 - socioeconomic and health, 171
 - (*see also* Suicidality)
 - self-injurious behavior risk, 176
 - self-reporting MH issues, 176
 - services, 176
 - skilled multicultural practitioners, 177
 - valuable resources, 176
 - war-torn decades, 168
 - “Yugonostalgia”, 168
 - “Yugosphere”, 168
- Eating disorder, 3
- Elluam Tungiinun* (Toward Wellness) prevention program, 64
- Enculturation, 6, 10, 58, 62, 64
- Ethnic affiliation, 73
- Ethnicity, 2, 5, 6, 13
- European-origin population
 - birth rate and progressive aging, 21
 - child mental health services, 21
 - health care system, 21
- F**
- Familism, 73, 76, 79
- Familismo, 6
- Family therapy, 28
- Family treatment, *see* Prevention of suicides
- First-generation South Asians
 - agricultural laborers, 86
 - lack of assimilation, 89
 - mental health professional, 93
- Formative Method for the Adapting Psychotherapy (FMAP), 128
- Former Soviet Union (FSU), 183, 187, 190
- G**
- Garrett Lee Smith Memorial Suicide Prevention Program (GLS program), 129
- Gay & Lesbian Alliance Against Defamation (GLAAD), 204
- Gay, Lesbian, and Straight Education Network (GLSEN), 204, 208
- Generalized anxiety disorder (GAD), 103, 150, 151
- H**
- High-risk individuals prevention
 - ABFT, 225, 228
 - CBT and DBT, 225
 - CBT-SP phases, 225–227
 - interdependent relationships encouragement, 225
 - SSSE, 228
- Hinduism, 6
- Historical trauma, AIAN/NHPI trauma, 34, 35, 37, 53, 54
- Hodja (Imam), 134
- I**
- Immigrants, from ME origin, 6
 - acculturation, 153
 - alcohol-related disorders, 156
 - cigarette smoking, 155
 - communication, 158
 - diagnosing depression, 158
 - first- and second-generation, 151, 153
 - intervention programs, 159
 - Muslim, 156
 - and refugees, 147
 - risk and protective factors, 157
 - screening, 152

- Immigration, Turkish American youth, 139
 Immolation, 151, 152
 Institute of Medicine (IOM), 203
 Integration, 6
 Intensive community-based treatment services, 28
 Interventions
 - cognitive behavior therapy, 44
 - dialectical behavior therapy, 44
 - evidence-based therapies, 44, 45
 - mentalization-based therapy, 44
 - MST, 43
 - NREPP, 44
 - SAMHSA, 44
 Islam, 7
- J**
 Judaism, 7
- L**
 Latino Americans (Hispanic/Latino Americans)
 - cultural heritage, 10
 - family closeness and relations, 10
 - immigration and acculturation processes, 11
 - nervios (nerves), 11
 - parent's language barrier, 11
 - personalismo, 10
 Latino youth
 - acculturation
 - acculturative stress, 71, 72
 - AFD, 72
 - cultural differences, 72
 - cultural diffusion, 71
 - discrimination, 73
 - etiological factors, 71
 - familism, 72
 - individuals, 71
 - Mexican-born, 71, 72
 - as risk factor, 71
 - suicidal attempts, 71
 - suicidal ideation, on immigrant adolescents, 71
 - case management
 - aspirin tablets, overdose of, 75
 - behavioral problems, history of, 76
 - depression, recurrent with psychosis and ADHD, 74, 75
 - individuation, attainment of, 76
 - as "perfect storm", 75
 - cause of death, 70
 - clinical approach
 - family involvement and intervention, 77
 - individual psychotherapy, 77
 - linguistic support, 78
 - verbalization and action, 77
 - culturally related risk factors, 73
 - epidemiological data, 69
 - ethnic group, 69
 - evidence-based therapies
 - BSFT, 79
 - cost-effective intervention, 79
 - SSSE, 79
 - TADS, 80
 - family dysfunction and socioeconomic status, 70
 - females, adolescent, 70, 71
 - growth rate, 69
 - immigrants, 70
 - minority groups, 73
 - vs. non-Latino Whites, 73
 - origins, 69
 - protective factors against suicide
 - ethnic affiliation, 73
 - familism, 73
 - personalismo, 74
 - religiosity and moral objections, 73
 - psychosocial treatments, 78, 79
 - risk factors, 70, 71
 - suicidality, 70
 - traditional Latino families, 71
 Lesbian, gay, bisexual and transgender (LGBT) youth, 37–40
- M**
 Major depressive disorder (MDD), 3, 11, 103, 151
 Malaysia, SEA, 103
 Marginalization, 6
 Marianismo, 6
 Massachusetts Child Psychiatry Access Project (MCPAP), 229
 Media, adolescent suicide
 - excessive attention, 236
 - lack of coverage, 236
 - overidentification, 236
 - repeated and explicit coverage, 236
 Mental health (MH), 14–16
 - disparities
 - contextual variables, 15
 - depression treatment, 14
 - families' cultural beliefs, 16
 - in ethnic minorities, 14
 - monolingual Hispanic and AA patients, 16
 - outpatient treatment, 15
 - service use, 16
 - social aggression, 15
 - help-seeking behavior, 13
 - immigration journeys and acculturative stress, 11
 - NCS-A, 5
 - utilization, AA, 10
 Mental health stigma, 8, 9, 13
 Mentalization-based therapy, 44
 Middle East (ME)
 - case presentation
 - cultural evaluation, 150
 - depressive symptoms, 148
 - insomnia and nightmares, 150
 - investigative tests, routine, 149
 - prevalence, mental illness, 152
 - psychiatric treatment, 149

- Middle East (ME) (*cont.*)
- PTSD symptoms, 148
 - stigma and sense of shame, 149
 - cause of death, 151
 - cultural difference, 148
 - culturally bound healing practices, 159
 - definition, 147
 - diagnosis
 - depression, anxiety, and PTSD, 158
 - of mental illness, 159
 - over-dependence and enmeshment, 158
 - family integration and rehabilitation, 159
 - gender difference, 148
 - immigration and mental health, 157, 158
 - immolation, 151, 152
 - incidence and prevalence, 150
 - intervention approaches
 - cultural complexity, 160
 - cultural sensitivity and understanding, 160
 - culturally orientated programs, 161
 - hospital treatment, 161
 - parental and adolescent acculturation, 160
 - physical contact, 160
 - rudimentary programs, 159
 - supportive social factors, 160
 - MDD, 151
 - mental illness, 147
 - OCD, 151
 - risk and protective factors
 - acculturation, 153
 - age, 155
 - alcohol-related disorders, 156
 - cigarette smoking, 155
 - country, 154
 - description, 152, 153
 - first-generation adolescent immigrants, 153
 - gender, 154, 155
 - incarceration, 155
 - second-generation adolescent immigrants, 153
 - social support, religion and cognitive functions, 156, 157
 - socioeconomic status, 156
 - substance abuse, 156
 - war, traumatic experience and suicide, 153, 154
 - vs. Swedish- born population, 151
 - treatment settings, 159
 - Migration stress
 - family's peri-migration, 194
 - Perez-Foster hypothesis, 190
 - Minority youth
 - ethnic, 225
 - gatekeeper training program, 233
 - generic treatments, 222
 - MH services and cultural stigma, 230
 - providers, 229
 - screening tools, 230
 - suicide hotline services, 224
 - Multisystemic therapy (MST), 28, 43, 44
- N**
- National Child Traumatic Stress Network (NCTSN), 144
 - National Comorbidity Study-Adolescent Supplement (NCS-A), 36
 - National Comorbidity Survey Replication-Adolescent Supplement (NCS-A), 3, 5
 - National Health and Nutrition Examination Surveys (NHANES), 203
 - National Registry of Evidence-Based Programs and Practices (NREPP), 44
 - National Survey of American Life (NSAL), 36
 - National Survey on Family Growth (NSFG), 203
 - Native Hawaiian (NH), population groups, 53
 - Native Hawaiian Pacific Islander (NHPI)
 - assimilation for, 54
 - demographics, 57
 - physical resources, loss of, 55
 - population groups, 53
 - strangulation, 56
 - suicide, cause of death, 53
 - suicide rate, 56
 - New York City (NYC), 136
 - Non-Hispanic White (NHW) group
 - vs. AAs youth, 8
 - boys vs. girls, 14
 - correlations, 12
 - depressive symptoms, for girls, 13
 - individualism and freedom, 12
 - "island of misery" hypothesis, 12
 - MH treatment, 14, 15
 - rate of suicide, 2
 - social aggression, 15
 - "the tip of an iceberg" of suffering, 12
 - Western societies and culture, 11
 - NSAL-Adolescent Supplement (NSAL-A), 36
- O**
- Obsessive-compulsive disorder (OCD), 151
 - Oppositional defiant disorder (ODD), 3
- P**
- Parent-child interaction, East Asian youth, 118
 - Parents and Friends of Lesbians and Gays (PFLAG), 204
 - Passive coping, 41
 - Personalismo, 6, 74
 - Pharmacotherapy, 26
 - Philippines, SEA, 105, 106
 - Positive and Negative Suicide Ideation Inventory (PANSI), 104
 - Post-traumatic stress disorder (PTSD), ME countries, 3, 11, 150, 151
 - Poverty, 53, 56, 57
 - Prevention of suicides
 - AILSDC program, 234, 236
 - American Academy of Child and Adolescent Psychiatry's model, 221, 222
 - challenges, 221
 - challenges, families of minority, 222

collectivist value orientation, 221
 (see also Community outreach)
 Golden Gate Bridge Patrol, 223
 gun locks and locations education programs, 223
 (see also High-risk individuals prevention)
 media, role, 236, 237
 NHW youth, 224
 previous attempts and ideations, predictions, 224
 suicide hotline numbers, 224
 text crisis line, 223
 training workshops, 224
 Trevor Project, 223, 224 (see Youth/family
 educational approach)
 Problematic Internet use (PIU), 127
 Project SHIFA in Boston, 230
Promotora de Salud model, 28
 Psychoeducation, 35
 Psychotherapy
 cognitive-behavioral interventions, 27
 confidentiality, 27
 cultural adaptations, 27
 goal, 27
 in individual sessions, 27
 professionalism, 27
 suicidality, 27
 value-neutral approach, 27

Q

Quran, 7

R

Race, 2, 6, 9, 10, 12, 13
 Racial discrimination, 89, 93, 95
 Racism, 8
 African-American descendants, 34
 and discrimination, 38
 parents and prosocial peer support, 41
 parents' experiences, 38
 Refugee, 6
 Jewish religious and, 183
 resilience concept, 189
 sectarian violence, 184
 Religion
 African religions, 7
 Buddhism, 5
 Christianity, 5
 Hinduism, 6
 Islam, 7
 Judaism, 7
 in North America, 7
 Religiosity, Turkish American youth, 73, 138, 139
 Risk factors, 126, 136
 East Asian youth (see East Asian youth)
 suicide, 38
 acculturation (see Acculturation)
 African-American adolescents, 37
 in Black youth, 37
 early adverse events, 37

“postmodern society”, in urban communities, 38
 racism and discrimination, 38
 societal impact, 37
 Turkish American (see Turkish American youth)
 Runaway behavior, 3
 Russianization process, 183
 Russification process, 183

S

Schizophrenia spectrum disorders (SSD), 40
 Second-generation South Asians
 social organizations, 87
 transnational ties, 88
 Separation, 6
 Sexual minority youth
 ACEs, 209
 adolescence life stage, 203
 adverse events, 209
 anxiety, 204–206
 asexual, pansexual and polysexual, 202
 BRFS, 203
 CDC official, 213
 child psychiatry, 212
 degree of victimization or discrimination, 208
 depression, anxiety and substance abuse, 212
 emotional problems and self-directed violence, 212
 evidence-based interventions, 213–216
 family rejection, 209
 gender and sexual identity, 201
 gender fluid/nonbinary, 206–208
 gender nonconforming, 201
 homelessness, high rates, 203
 impression management, 204
 internalized homophobia, 211
 internalized homophobia/transphobia, 204
 International law, 210
 IOM, 203
 loss anticipation, 211
 minority stress paradigm, 204
 national studies, 203
 nonbinary gender identity, 209
 positive outcomes, 210
 “propaganda laws”, 210
 racial groups, 210
 romantic and sexual being, 202
 same-sex unions, 213
 in school communities, 209
 self-directed violence, 204
 shame/victimization, 211
 social-cultural stereotypes, 210
 stigma consciousness, 204
 terminology and definitions, 201, 202
 and trans*-youth suicide, 211
 US Secretary of Health and Human Services, 212
 YRBS, 212
 Shame, 122, 123
 Somatization, 126
 Somatization in immigrants, Russian-Jewish
 immigrants, 192

- South Asian mental health, 94, 95
 - interventions, 91
 - recommendations, 93
 - sexuality, 93
 - specialist services, 92
- South Asian (SA) youth
 - cultural factors, providers, 93
 - culturally specific symptoms, 92
 - deliberate self-harm, 90, 91
 - interventions research, 93–95
 - presentation among suicidal, 90, 91
 - protective factors, 91
 - racism, 89
 - salience identification, 86, 87
 - sexuality, 87–88
 - violence, family issue, 89, 90
 - weak ethnic identities, 88
- Southeast Asia (SEA)
 - Cambodia, 104
 - cognitive-behavioral interaction, 109
 - countertransference and transference issues, 109
 - drug use disorder, 100
 - epidemiology, 101, 102
 - Filipino immigrants, 100
 - gatekeepers, 108
 - hopelessness, 107
 - impulsiveness, role of, 108
 - integration and cultural assimilation, 103
 - malaysia, 103
 - medical conditions, 100
 - model minority myth, 103, 106
 - novel identification strategies, 108
 - Philippines, 105, 106
 - predisposing and risk factors, 100
 - psychological autopsy, 107
 - psychotherapy, 109
 - psychotropic medications, 109
 - public awareness and media portrayal, 108
 - risk factors, 102, 107
 - role of discrimination, 100
 - school-based interventions, 108
 - social connection and support, 103
 - SPI and STOPS, 107
 - suppressive leadership, 100
 - Thailand, 104
 - Vietnam, 104, 105
- Sovietization process, 183
- Soviet-Jewish youth
 - acculturation progress, 185
 - American peers and higher grades, 185
 - antidepressant medication, 192
 - anti-Muslim prejudice and racism, 184
 - bicultural identity and skills, 187
 - bullying and peer aggression, 187
 - 1986 Chernobyl nuclear reactor, 190
 - child-raising practices, 184
 - comprehensive and pervasive disruptions, 186
 - cultural and educational learning opportunities, 184
 - cultural precedence, 183
 - depression and somatization, 191
 - diagnosis, 192, 193
 - discriminatory experiences and negative perceptions, 187
 - educational and professional goals, 187
 - emotional distress, 191
 - epidemiological data and multivariate statistical analysis, 189–190
 - ethnic minorities, 188
 - family difficulties and underemployment, 191
 - Former Soviet Union (FSU), 183
 - friendship networks, 189
 - growth and authentic living, 185
 - “Havurah”, Hebrew for “Fellowship”, 189
 - immigrant vs. native-born adolescents, 191
 - immigrants, 186
 - immigration and naturalization service, 183, 187, 191
 - intervention approaches, 193–195
 - Israeli cultural norms, 187
 - native-born peers, 192
 - nonacademic extracurricular activities, 185
 - paradoxical dichotomy, balancing, 185
 - parentification, 185
 - parenting styles, 184
 - Perez-Foster hypothesis, 190
 - prevalence and risk factors, 190
 - primary care setting, 191
 - psychological distress, 191
 - relational intimacy and mutuality, 189
 - religious and political refugees, 183
 - research, 186
 - resilience concept, 189
 - risk suicidal thoughts, 184
 - russification process, 183
 - scholarly discussions, 188
 - school environment, 189
 - social identity and culture, 186
 - somatic complaints, 189
 - somatic manifestations, 191
 - somatic symptoms, 184
 - suicidal ideation, Jewish adolescents, 188
 - territories, 183
 - traumatic history, 183
 - troublemakers, 187
 - workforce and education, 183
- Stigma, vi, viii, 5, 8, 16, 22, 123
 - African Americans and stigma, 35, 43, 44
 - East Asians and stigma, 122, 123
 - Eastern Europeans and stigma, 172
 - family and community interventions and stigma, 221, 223, 224, 229, 230, 234, 237
 - gatekeeper model, media, 233
 - Latinos and stigma, 77
 - LGBT and stigma, 204
 - case, 204, 209, 212–214, 216
 - Mental Health stigma, 8, 9, 11, 13, 15, 93, 95, 123, 136, 224, 229
 - Middle Eastern and stigma, 149–161
 - case, 159, 161
 - South Asians and stigma, 83–95

- Southeast Asians and stigma, 100
 - transgender, 102, 104
 - suicide stigma, 24–26
 - Turkish and stigma, 136
 - Strategic structural systems engagement (SSSE), 79, 228
 - Strengths and Difficulties Questionnaire (SDQ), 77
 - Substance abuse, Latino youth, 71, 72, 77, 78, 80
 - Substance Abuse and Mental Health Services Administration (SAMHSA), 44
 - Suicidal behavior, 6
 - description, 3
 - disparities, in MH treatment, 14
 - ethnic disparities, 12, 13
 - female adolescents, 4
 - help-seeking behavior, 13
 - MH disparities, 15, 16
 - predictors of, 3
 - prevalence of, 3
 - psychosocial stressors, 4
 - risk factors, 13
 - somatic symptoms, 13
 - symptoms severity and concern level, 3
 - treatment-seeking behavior, 5
 - in Turkish American youth, 136, 137, 139, 143
 - Suicidal ideation, Turkish American youth, 70, 71, 77
 - careful attention, 140
 - direct inquiry, 141
 - ethnic groups versus majority adolescents, 138
 - and friendship, 138
 - Pessimism, 137
 - risk factors, 136
 - Suicidality
 - AIAN youth, 9
 - among Turkish American youth immigrants, 136, 140
 - assessment, culturally informed, 24
 - acculturation and assimilation, Berry's model of, 23
 - diverse youth (*see also* Diverse youth)
 - DSM Cultural Formulation, 24
 - explanatory models approach and interview questions, 24
 - family members, 25
 - family's language, use of, 26
 - interview process, 24
 - non-judgmental approach, 24
 - post- and pre-suicide attempt, 23
 - precipitating factors, 23
 - suicidal and community crisis services, 25
 - sustaining culturally related factors, 23
 - translated written materials, 26
 - Durkheim's theory, 172, 173
 - effect of war, 172
 - ethnic minority, 12
 - firearms, 173
 - male-to-female ratio, 173
 - in NHW, 12
 - religious influences increase, 173
 - repetition of attempt, 174
 - resiliency factor, Latino males and females, 10
 - social stressors, 172
 - treatment, culturally informed
 - cultural consultants, use of, 26
 - cultural factors, 26
 - pharmacotherapy, 26
 - psychoeducation, continued, 26
 - in Turkish adolescents, 138–140
 - types of modes, 172, 173
 - Vietnam syndrome, 173
 - Suicide, 6
 - among Turkish American immigrants, 138, 140, 142
 - among Turkish immigrants, 136
 - definitions, 3, 6
 - "honorable", 5
 - Suicide attempt, 3, 4, 6
 - Latino youth
 - cost-effective intervention, 79
 - culturally related risk factors, 73
 - depression, rates of, 70
 - ethnic affiliation, 73
 - etiological factors, 71
 - Mexican adolescents, 72
 - prevalence, 70
 - psychosocial treatments, 78
 - vs. US-born Latino adolescents, 70
 - Turkish American youth, 137
 - Suicide ideation, 6
 - Suicide Prevention International's (SPI) strategies, 107
- T**
- Teacher's Report Form (TRF), 77
 - Thailand, SEA, 104
 - Treatment for Adolescents with Depression Study (TADS), 80
 - TrevorChat, 224
 - Trevor Lifeline (24/7 hotline), 223
 - Trevor Project, 223
 - TrevorText, 224
 - Turkish American youth
 - case presentation
 - adjustment problems, 134
 - "down, depressed, and lonely", 133
 - medical problems, 133
 - self-harming behaviors and suicidal ideations, 133
 - cultural factors
 - educated, moderately religious family structure, 135
 - educated, secular family structure, 135
 - ethnic groups, Turkish-speaking, 134
 - extended family, 136
 - graduate and postgraduate students, 135
 - Islamic practices, 135
 - mental health stigma, 136
 - Northern Jersey-NYC area, 136
 - Ottoman Empire, 134
 - traditional family structure, 135
 - culturally specific presentations and symptom expression, 139, 140

Turkish American youth (*cont.*)

diagnosis

- careful attention, 140
- C-SSRS, and hospital-wide screening implications, 141
- immigration and acculturation processes, 140
- intolerable anxiety, 141
- loneliness, 141
- suicide attempts, 141

intervention approaches

- cultural differences, 142
- emergency contacts, list of, 143
- in crisis, 142
- individual and family/system categories, 142
- nonjudgmental approach, 144
- school support, 143
- social, cultural, and economic factors, 141
- suicide hotline and text line, 142, 143
- underlying psychopathology, 143
- working with family, 143

risk factors

- alcohol and substance abuse, 139
- family connectedness, 138
- friends and social support, 138
- immigrant youth, 136
- immigration, 136, 139
- previous suicide attempt, 137
- psychiatric illness and self-esteem, 137
- religiosity, 138
- suicidal ideation, 136

suicidality, 137

suicide and mental health disorders, family members, 137

Tuskegee Syphilis Trials, 35, 43

V

Vietnam, SEA, 104, 105

Vietnam syndrome, 173

Violence, 77

W

Weak ethnic identities

- individualism and self-assertion, 88
- risk factor, youth suicidal behavior, 88
- “role conflict” experience, 88
- second generations, 88
- transnational ties, 88

X

Xenophobia, 21, 27

Y

Youth/family educational approach, 229, 230

Youth Risk Behavior Survey (YRBS), 35, 36, 203, 212

Youth Self-Report (YSR), 77

Youth suicide, 12