

Index

A

Abdominal pain

- ACD (*see* Acute colonic diverticulitis (ACD))
- costs, 295
- definition, 294
- ED census, 294
- hospital discharges, 294
- imaging, 295
- National Hospital Discharge Survey, 294, 295
- pancreatitis (*see* Acute pancreatitis)

Abdominal radiographs, 570–571

Abusive head trauma (AHT), 546, 547, 550

Acetabular fractures, 463–465, 468, 472–474

ACJ separation, *see* Acromioclavicular joint (ACJ) separation

ACR appropriateness criteria, 29, 44, 51

Acromioclavicular joint (ACJ) separation, 451–453, 456, 458, 460

Acute abdomen, pregnancy

- abdominopelvic injuries, 408
- appendicitis (*see* Appendicitis)
- bowel obstruction, 403, 404
- ectopic, 406, 407
- fetal radiation exposure, 400
- imaging, 400, 401
- imaging protocols, 408–411
- MeSH terms, 401
- OB/GYN, 400
- ovarian torsion, 407, 408
- physiologic changes, 400
- placental abruption, 407
- renal colic, 404, 405
- Roux-en-Y gastric bypass, 408
- RUQ pain, 403, 408
- SOGC gradation, levels of evidence, 404, 408
- sonography, 400
- trauma, 405, 406

Acute acalculous cholecystitis (AAC)

- cholescintigraphy, 314, 315
- computed tomography (CT), 315
- imaging strategy, 315, 316
- ultrasonography (US), 313, 314

Acute aortic syndrome (AAS) in adults

- CABG procedure, 214

chest radiography, 210, 217

crushing chest pain, 214

CT angiography, 207, 210, 211

CTA chest, 217

definition, 207–209

dual energy, 213

echocardiography, 211, 212

EKG synchronization, 212–213

epidemiology, 209

high pitch, 213

hypotensive, 217

imaging, 209

IRAD investigators, 209

methods, 209

MRI/MRA, 207, 217

nontraumatic, 207

outpatient, presurgical planning, 214

pathophysiology, 207–209

PET/CT, 212

triple-rule-out CT angiogram, 207, 213, 214

Acute back pain

breast cancer, 177

cervical spine fracture, 177

CT, 167

definitions, 167

disk herniation, 172, 173

epidemiology, 167

foot, 176

fractures, 168–172

hands, 175

imaging, 168

imaging modalities, 175

infection, 173, 174

metastatic disease, 175

methods, 168

Michigan lumbar spine protocol, 177

MRI, 167

pathophysiology, 167

post surgery, disk herniation, 176

research, 177

sensitivity and specificity, 176

thoracolumbar spine fracture, 177

uncomplicated, 167

US patients, 168

- Acute biliary disorders
 - AAC, 313–316
 - ACC, 311–313
 - bile duct obstruction, 316–317
 - choledocholithiasis, 318–320
 - cost to society, 310–311
 - definition, 310
 - epidemiology, 310
 - imaging, 311
 - methodology, 311
 - pathophysiology, 310
- Acute calculous cholecystitis (ACC)
 - cholescintigraphy, 312, 313
 - computed tomography (CT), 312
 - imaging strategy, 313, 314
 - magnetic resonance imaging (MRI), 312
 - ultrasonography (US), 311–313
- Acute cholecystitis
 - prevalence, 310
- Acute colonic diverticulitis (ACD), 306
 - clinical guidelines, 305
 - consensus guidelines, 304
 - CT protocols, 304, 306
 - diagnostic accuracy, MRI, 305
 - diagnostic performance
 - CT, 306
 - US, 306
 - intraluminal pressure, 303
 - MEDLINE database, 303
 - mesenteric fat, 304
 - oral, rectal, and intravenous contrast, 305, 306
 - physician needs, 307
 - right lower quadrant abdominal pain, 306, 307
 - sonography, 304, 305
 - symptoms, 304
- Acute colonic pseudo-obstruction, 331
- Acute coronary syndrome (ACS), *see* Chest pain, coronary CTA
- Acute diverticulitis
 - definition, 294
- Acute headache disorders
 - adults with brain metastases, 120, 121
 - adults with migraine/chronic headaches, 119, 120
 - adults with new-onset headache, 116
 - advanced brain imaging, 114
 - ataxia, 128
 - children with headache, 121–123
 - colloid cyst, 128
 - cost-effectiveness, neuroimaging, 127
 - cough and exertion, 128
 - CT and MR imaging, 123
 - CT imaging, 113, 128
 - CTA, 113
 - definition, 114
 - direct and indirect annual cost of migraine, 115
 - DSA, 113
 - epidemiology
 - adults, 115
 - children, 115
 - and guidelines, 128
 - high-risk adults with new onset headache, 116–119
 - imaging, 116
 - MDCT angiography, 113
 - methods, 116
 - MR imaging, 128
 - MRI, 113
 - pathophysiology, 114
 - primary, 125–127
 - subarachnoid hemorrhage, 123–125
- Acute hematogenous osteomyelitis (AHOM)
 - Bartonella henselae*, 511, 512
 - bone rarefaction and periosteal reaction, 500
 - bone scintigraphy, 502
 - costs, 500
 - definition, 497
 - diabetic pedal osteomyelitis, 503
 - diagnostic performance characteristics, 507
 - fever and septicemia, 498
 - ¹⁸F-FDG, 502
 - imaging algorithm, 507, 508
 - left femoral osteomyelitis, 509, 510
 - left proximal humerus osteomyelitis, 511
 - MRI, 501
 - MRSA, 498
 - pathologies, 498
 - periosteum, 498
 - populations, 499
 - proximal left tibia osteomyelitis, 509, 510
 - radiography, 501
 - SCD, 503, 504
 - transphyseal vessels, 498
 - tuberculous osteomyelitis, 498
 - ultrasound, 501–502
 - WBC scintigraphy, 502
- Acute intestinal ischemia
 - atrial fibrillation, 378, 379
 - celiac artery, 374
 - classification, 374
 - CT (*see* Computed tomography (CT))
 - definition, 373
 - diagnosis, 373
 - endovascular revascularization (*see* Endovascular revascularization)
 - epidemiology, 375
 - goals, 375
 - hypodense thrombus, 375
 - inferior mesenteric artery, 374
 - lactic acidosis, 374
 - local and systemic complications, 374
 - methodology, 375–376
 - multidetector array scanner, 378
 - multiplanar reformations, 380
 - phases, 379
 - pneumoperitoneum, 375
 - presplanchnic, splanchnic and postsplanchnic, 374, 375
 - serum lactate, 374
 - SMA, 375

- stages, 374
 - superior mesenteric artery, 374
 - symptoms, 374
 - vasoconstriction, 375
 - vasodilation, 375
 - vomiting, diarrhea and fever, 378, 380
- Acute intramural hematoma (IMH), 207, 208, 210, 211, 215, 216
- Acute ischemia, 96, 97
- Acute lower GI bleeding
 - catheter angiography, 359–360
 - CTA, 357–358
 - nuclear medicine, 358–359
 - radiological management
 - ACR Appropriateness Criteria, 361
 - catheter angiography, 360
 - colonoscopy, 360
 - CTA, 360
 - efficacy and safety of superselective microcoil embolization, 360
 - hemodynamically unstable patient, 360
 - malignancy, 361
 - randomized controlled trial, 360
 - rebleeding rates, 361
 - superselective microcoil embolization, 361
 - transcatheter embolization, 361
- Acute musculoskeletal infections
 - AHOM (*see* Acute hematogenous osteomyelitis (AHOM))
 - cellulitis, 499, 500, 507, 509
 - extremity MR imaging, 514
 - imaging, 500
 - incidence, 500
 - MEDLINE search, 500
 - radiographs, 512–514
 - radionuclide bone scintigraphy, 514
 - SA (*see* Septic arthritis (SA))
 - soft tissue abscess (*see* Necrotizing fasciitis (NF))
 - ultrasound hip joint, 515
 - whole-body MR imaging, 514
- Acute neurological deficits, 107
- Acute pancreatitis
 - APACHE II scoring system, 298
 - Atlanta classification system, 296
 - choledocholithiasis, 296
 - clinical assessment, 297
 - CT protocols, 301
 - CT scoring protocols, 296
 - CT scoring systems, 297
 - definition, 294
 - diagnosis and evaluation, MRI, 299, 300
 - ERCP *vs.* MRCP, 300
 - gallstones and alcohol abuse, 296
 - grading and scoring systems, 297
 - IAP/APA, 298
 - laboratory testing, 296
 - MEDLINE database, 295
 - modified CT severity index, 297
 - MRI protocol, 301–303
 - radiological scoring systems, 298
 - sonography, 299
- Acute pelvic pain
 - acute abdominal pain and urinary retention, 429, 431
 - clinical presentation, 419
 - congenital vaginal obstruction, 428, 429
 - definition, 416
 - differential diagnoses, 429
 - gynecological process, 416
 - imaging protocols, 429–431
 - pediatric emergency setting, 421
 - pediatric population, 421
 - physical examination, and laboratory testing, 419
 - radiography, 419
 - sensitivity and specificity, ultrasound, 423, 429
 - serum β -hCG-negative patients, 421
 - serum β -hCG-positive patients, 420, 421
 - symptoms, 420
 - transvaginal and transabdominal US, 420
- Acute pyelonephritis (APN), 616, 620, 622, 623
 - See also* Urinary tract stones
- Acute respiratory illness (ARI)
 - acute bronchitis, 262
 - causes of pulmonary infection, 275
 - CD4 count, HIV/AIDS, 264, 273–275
 - CT
 - CAP, 271
 - CXR (*see* Chest radiograph (CXR))
 - definition, 262
 - imaging, 264 (*see also* Pneumonia)
 - PubMed, MEDLINE, and Google Scholar search databases, 264
 - viral causes, 268
- Acute scrotal pain
 - causes, swelling, 436, 439
 - clinical and radiographic examinations, 437
 - costs, 436
 - definition, 435
 - epididymitis, 437
 - ischemia, 436
 - spermatic cord torsion, 436
 - testicular torsion (*see* Testicular torsion)
- Acute sinusitis
 - acute myeloid leukemia, 198, 200
 - allergic fungal infection, 192, 199
 - allergy/asthma, 184
 - American Academy of Pediatrics clinical guidelines, 195
 - amoxicillin, 195
 - anatomical variations, 194
 - antibiotics, 185, 186, 191
 - CEA, 194
 - complications, 191
 - contrast-enhanced CT/MR, 191, 192
 - costs and adverse effects, 194
 - CT and MRI, 187
 - definition, 198, 199
 - description, 184
 - effectiveness of antibiotic therapy, 195

- Acute sinusitis (*cont.*)
- endoscopic surgery, 193
 - fungal infection, CT scans, 192, 199
 - guidelines, children, 187, 188
 - headache and mental status change, 193, 200
 - imaging and clinical examinations, 198, 200
 - nasal swab and culture, 187
 - orbital and intracranial complications, 186
 - orbital cellulitis, 191
 - ostia obstruction, 184
 - physical examination, 187
 - radiographs cost, 186
 - recommendations, 194
 - respiratory symptoms, 186, 187
 - sinus radiography and CT, 188, 189
 - Streptococcus pneumoniae* and *Haemophilus influenzae*, 184
 - surgery, 191
 - symptoms, 190, 191
 - viral upper respiratory infection, 198, 199
- Acute spine trauma
- Atlanto-occipital joint, 162
 - cervical spine CT, 151
 - cervical spine in after blunt trauma, 153–155
 - cervical spine in blunt trauma, 155–160
 - clinical prediction rules, 164
 - CT protocol, 163, 164
 - definition, 151, 152
 - imaging, 152
 - interarticularis fracture, 162
 - methods, 152
 - NEXUS and Canadian cervical spine rules, 151
 - pathophysiology, 151, 152
 - radiography protocol, 163
 - spinal cord injuries in the USA, 152
 - thoracic and lumbar spine after blunt trauma, 160, 161
 - thoracic and lumbar spine in blunt trauma, 161, 162
 - thoracolumbar spine imaging, 151
- Acute spine trauma
- cervical spine in blunt trauma, 156
- Acute upper GI bleeding
- CTA, 361, 362
 - Nuclear Medicine, 362–363
 - radiological management
 - CTA, 363
 - endoscopic management, 363
 - retrospective study, 363
 - superselective microcoil embolization, 363
 - TIPSS, 363, 364
- Adnexal/ovarian torsion
- adnexal mass, 427
 - causes, 426
 - CT and MRI, 427
 - definition, 417
 - Doppler flow, 427
 - follicles, 427
 - menarche, 418
 - neonatal and antenatal period, 418
 - ultrasound, 426
 - whirlpool sign, 427
- Adults
- hip fracture, 481, 482
 - new-onset headache, 116, 117
- Advanced pulmonary CT angiographic techniques, 239, 240
- Advanced Trauma Life Support (ATLS), 249
- AHT, *see* Abusive head trauma (AHT)
- Air enema, 567, 570, 572, 573, 575–578, 580
- Air vs. liquid enema, 573–575
- Alberta Stroke Program Early CT (ASPECTS), 91, 97–99, 103, 110
- American Academy of Otolaryngology-Head and Neck Surgery (AAO-HNS), 194
- American Academy of Pediatrics (AAP), 615, 616, 621, 624, 625, 627, 630
- American Association of Family Physicians (AAFP), 267
- American College of Gastroenterology, 361
- American College of Radiology, 361, 377
- American Society for Gastrointestinal Endoscopy, 361
- Aneurysms, 113, 114, 116, 124, 125
- Angiography, 357
- CTA (*see* CT angiography (CTA))
- Ankle and foot injuries
- acute tendinous rupture, 489
 - ankle sprains, 488
 - calcaneal apophysitis, 487
 - distal fibular fractures, 487
 - distal tibia, 486
 - epiphyseal injury, 486
 - fractures of phalanges, 487
 - intra-articular fractures, 488
 - juvenile Tillaux fracture, 487
 - Lisfranc joint disruption, 488
 - metatarsals, 488
 - OAR, 486, 489, 490
 - osteochondral lesions, 488
 - Sever's lesion, 487
 - talar fractures, 488
 - triplane fracture, 487, 490, 494
- Ankylosis, 152
- Annals of Internal Medicine Clinical Guidelines, 361
- Anti-cysticercus antibody, 139
- Aortic dissection (AD), 207–212, 214
- Aortic rupture, 207, 209, 216
- Appendicitis
- ACR guidelines, 290, 291
 - adverse effect, 288
 - clinical and laboratory examination, 287
 - cost- effectiveness, 402–403
 - CT protocols, 290
 - decision-making, 288
 - definition, 281–282
 - diagnosis, 402
 - diagnostic performance, MRI, 284
 - emergency department, 284

- imaging goals, 282
 - IV contrast, 290
 - laparoscopic surgery, 402
 - lifetime incidence, 282
 - meta-analyses, 282
 - MRI cost and scan time, 283
 - MRI utilization and accuracy, 284
 - negative appendectomy rate, 288, 289
 - pediatric patients
 - clinical decision rule, 287
 - cost-effectiveness analysis, 286
 - limitations, 285
 - MRI utilization, 286
 - negative appendectomy rates, 286
 - US and CT, 285, 286
 - pregnancy, 282
 - PubMed search, 282
 - recursive partitioning analysis, 287
 - RLQ abdominal pain and nausea, 289, 290
 - sensitivity and specificity, CT diagnosis, 283, 284
 - societal cost, 282
 - sonography, 402
 - ultrasound and MRI, 402, 409
 - wall ischemia, 282
 - Applying the evidence, 13–16
 - ARI, *see* Acute respiratory illness (ARI)
 - Arnold-Chiari type I malformation, 120
 - Arteriovenous malformations, 114, 116, 119, 120, 122
 - Artery susceptibility sign, 528
 - As Low as Reasonably Achievable (ALARA), 39
 - Assessing medical literature, 5–10
 - Ataxia, 129
- B**
- Balloon-occluded retrograde transvenous obliteration (BRTO), 364
 - Barium, 587–589
 - Bayes' theorem, 17
 - BCVI
 - screening criteria, 81
 - Bias
 - detection and correction, 22
 - observer, 22
 - reference standard, 23
 - risk of, 22
 - in screening, 24, 25
 - selection, 22
 - spectrum, 22
 - systematic, 19
 - type of, 22
 - Bile duct obstruction
 - computed tomography (CT), 317
 - EUS, 317
 - MRCP, 317
 - ultrasonography (US), 317
 - Blunt cerebrovascular injury (BCVI)
 - duplex sonography, 83
 - high risk, 80
 - incidence rate, 80
 - probability of, 80
 - screening and diagnosis, 82
 - screening criteria, 86
 - Blunt neck trauma
 - patients, 79
 - vascular injury
 - CT angiography, 82
 - DSA, 82
 - digital subtraction angiography, 82
 - duplex sonography, 82–83
 - MRI and MR angiography, 83
 - Blunt trauma
 - abdomen
 - allergic reaction, 254
 - arterial phase images, 254
 - CT examinations, 252
 - diagnostic evaluation, 252
 - FAST (*see* Focused assessment with sonography for trauma (FAST))
 - hemodynamic status and blood loss, 251
 - intraabdominal hemorrhage, 252
 - microscopic hematuria, 255
 - motor vehicle accident, 255, 258
 - prevalence, 251
 - signs and symptoms, 251
 - whole-body CT, 254
 - cause of death, 248
 - definition, 247
 - hospitalizations cost, 248
 - imaging, 248
 - imaging protocols, 255–257
 - mechanisms, 247
 - Medline/EDLINE/PubMed search, 248
 - motor vehicle accidents, 248
 - thorax
 - ATLS, 249
 - description, 248–251
 - hand-gliding accident, 255, 257
 - NEXUS Chest trial, 249, 250
 - portable chest radiography, 253
 - risk factors, mortality, 249
 - ultrasound examinations, 253
 - Blunt traumatic extracranial cerebrovascular injury, 80
 - Bone marrow transplant (BMT), 198
 - Bowel necrosis, 570
 - Bowel obstruction, 403, 404
 - Brain ischemia
 - CT, 527
 - DWI, 527, 528
 - perfusion CT and MRI, 528, 529
 - PET and SPECT, 529
 - stroke centers, 526
 - Brain stem glioma, 120
 - Breast cancer, 179
 - British Thoracic Society (BTS), 267, 268

- C**
- Caecal volvulus, 331
 - Calyceal blunting, 617
 - Canadian Assessment of Tomography for Childhood Head injury (CATCH), 67, 69
 - Canadian C-spine rule, 151, 155
 - Canadian C-spine study, 154
 - Canadian CT Head Rule (CCTH), 51, 53
 - Carotid artery disease
 - high prevalence, 8
 - low prevalence, 8
 - Carotid injury, 85
 - Carotid ultrasound, 532, 533
 - Catastrophic volvulus, 584
 - Catheter angiography, 222
 - acute lower GI bleeding, 359–360
 - Cavernous angioma, 120
 - CDS-CPOE system, 44
 - Centers for Disease Control (CDC), 568
 - Centers for Disease Control and Prevention (CDC), 584
 - Central nervous system (CNS), 115, 121, 122
 - Cerebral venous thrombosis, 526
 - Cervical spine
 - after blunt trauma
 - Canadian cervical spine prediction rule, 154, 155
 - NEXUS, 153
 - nexus prediction rule, 153, 154
 - blunt trauma
 - accuracy studies, 155, 156
 - children, 159, 160
 - cost-effectiveness analysis, 155
 - elderly patients, 159
 - high- and moderate-risk patients, 156
 - low-risk patients, 156–158
 - obtunded patients, 158, 159
 - harborview high-risk, 158
 - in low-risk patients, 157
 - Chest pain, coronary CTA
 - catheter angiography, 222
 - coronary artery disease, 225
 - cost savings and length of stay, 224
 - cost-effectiveness, 220
 - definition, 219
 - emergency department (ED), 220
 - ICA, 221
 - imaging, 220
 - low-risk patients, 221, 222
 - moderate- to high-risk patients, 222
 - myocardial ischemic/infarction events, 220
 - normal ECG and negative troponin, 225
 - performance characteristics, 221, 224
 - radiation exposure, 223
 - randomized controlled trials, 221
 - ROMICAT -II, ACRIN-PA, and CT-STAT trials, 224
 - SPECT, 223
 - triple rule out CT examination, 223
 - utilization, 221
 - Chest radiograph (CXR), 266–267
 - bacterial differentiation and viral pneumonia, 268–270
 - CAP (*see* Community-acquired pneumonia (CAP))
 - diagnosis, 267
 - frontal and lateral, 265
 - ionizing radiation, 266
 - Child abuse
 - abdominal injuries, 549
 - abdominal trauma, 551–552
 - bowel injuries, 549
 - costs, 546
 - CT and MRI scans, 549, 550
 - dating fractures, 549
 - definition, 545
 - emergency department, 546
 - head trauma, 550
 - imaging goals, 546
 - liver injuries, 549
 - long bone fractures, 548–549
 - Medline search, 546
 - meta-analysis, 546
 - metaphyseal corner fractures, 548
 - mortality rate, 546
 - rib fractures, 548
 - SDH and AHT, 546, 547
 - skeletal survey, 550, 551
 - skull fractures, 547, 548
 - Children
 - headache disorder, 121
 - Children's Head injury Algorithm for the prediction of Important Clinical Events (CHALICE), 67, 69, 70
 - Cholelithiasis
 - computed tomography (CT), 318
 - ERCP, 319
 - EUS, 319
 - imaging strategy, 314, 319, 320
 - MRCP, 319
 - ultrasonography (US), 318
 - Chronic pancreatitis, 300
 - Chronic renal failure (CRF), 618
 - Chronic sinusitis
 - anatomical variations, 196
 - CT scan, 196
 - definition, 195
 - diagnosis, 185, 199
 - headache, 196
 - medical management, 196
 - quality of life, 196
 - surgery, 196
 - vague symptoms, 195
 - Chronic Sinusitis Survey and Rhinosinusitis Disability
 - Index scales, 196
 - Clinical decision support (CDS) systems, 28–33, 43
 - Clinical decision support systems, 37, 43
 - Clinically important TBI (ciTBI), 67, 68
 - Closed loop obstruction, 330, 336, 337, 340, 349
 - Cluster headache, 114, 125, 126
 - Collateral flow-based methods, 106
 - Colloid cyst, 120, 128
 - Colony-forming units (CFUs), 616
 - Common bile duct (CBD), 317

Community-acquired pneumonia (CAP)

- AAFP, 267
- airspace disease, 264
- airspace opacification, 266
- bronchiolitis, 268
- BTS, 267
- CT, 271
- IDSA and BTS, 268
- indications, 266
- patients with ARI, 266
- radiographic evidence, 266
- respiratory syncytial virus, 269
- societal guidelines, 266, 267
- test characteristics, 267

Complex febrile seizures (CFS), 136

Computed tomographic angiography (CTA), 101

Computed tomography (CT)

- acute intestinal ischemia
 - acute mesenteric ischemia, 376
 - conventional angiography, 376
 - diagnosis, 376
 - MDCT, 376
 - meta-analysis, 376
 - MRA, 377
 - ultrasound, 377
- LOB, 341–343
- SBO
 - associated bowel ischemia/strangulation, 340
 - detection of level and cause, 336, 337
 - emergency department, 332–333
 - nonoperative/operative management, 338

Computed tomography angiography (CTA), 209

Computerized physician order entry (CPOE), 30, 31

Congenital vaginal obstruction, 428, 429

CONSORT studies, 19

Conventional CT angiography (CTA), 113, 124, 125

Cooperative Study of Sickle Cell Disease (CSSCD), 522

Cost–benefit analysis, 10

Cost-effectiveness analysis (CEA), 9, 10, 127, 194

Cost-minimization analysis, 10

Cost–utility analysis, 10

Cranio cervical region, 79

Critical assessment, 19

CT angiography (CTA), 80–82

- acute lower GI bleeding, 357–358
- acute upper GI bleeding, 361–362
- descriptive analysis, 83
- protocols, 89

CT dose index (CTDI), 39

CT myelography, 173

CT perfusion imaging (CTP), 95, 96, 100, 104–106

CT pulmonary angiography, 228–230, 232–234, 236, 237

Current Procedural Terminology (CPT) code, 14

D

D-dimer, 230–232

Delayed repeat enema (DRE), 576

Detorsion, 584

Diabetic pedal osteomyelitis, 503

Diagnostic imaging process, 29

Differential reference standard bias, 23

Diffusion tensor imaging (DTI), 49, 53–55, 59–61, 70–72, 76, 126

Digital subtraction angiography (DSA), 82, 124

Digital subtraction catheter-directed angiography (DSA), 100

Disk herniation, 167, 168, 173, 176, 178

Distal intestinal obstruction syndrome, 330

Distal intestinal obstruction syndrome (DIOS), 336

Doppler ultrasound, 22

Dose length product (DLP), 39

Dose reduction techniques, 386

Down's syndrome, 584, 585

Duodenal–jejunal junction (DJJ), 587–589, 592, 593

Duodenum inversum, 587

Duplex sonography, 82–83

Duplex ultrasound, 234, 235, 242

E

Eastern Association for the Surgery or Trauma (EAST), 169

Echocardiography, 235, 236

Economic analysis

- assessment
 - cost, 11
 - direct cost, 11
 - fixed costs, 11
 - indirect cost, 11
 - outcomes, 11
- cost–benefit analysis, 10
- cost-effectiveness analysis, 10
- cost-minimization analysis, 10
- cost–utility analysis, 10
- well-defined types, 10

Ectopic pregnancy

- abdominal ultrasound, 422
- adnexal mass, 420
- costs, treatment, 419
- definition, 416
- intrauterine gestational sac, 422
- maternal death, 417
- MRI, 422
- transvaginal US and serial quantitative β -hCG, 422
- ultrasound, 422

ED Setting

- biological effects, 40
- diagnostic imaging, 38, 39
- inappropriate use, 43, 44
- medical imaging, 40
- radiation mechanisms of effects, 39–40
- radiation terminology, 39
- time cost, 42

Elbow dislocation

- CT, 449
- imaging algorithm, 453, 456
- MRI, 449
- ossification centers, 454

- Elbow dislocation (*cont.*)
 proximal radius and ulna, 448
 radiographs, 448, 449
 surgical management, 449
 vascular injury, 449, 450
- Electroencephalographic (EEG) abnormalities, 134
- Electronic health record (EHR), 27
- Electronic medical record (EMR), 27, 31
- Embolization, 356, 360–363, 365
- Emergency imaging
 abdominal pain, 294–307
 acute abdomen, pregnancy, 400
 acute musculoskeletal infections, 497–515
 acute pelvic pain, 431
 acute scrotal pain, 439
 acute sinusitis, 184
 appendicitis, 281–291
 ARI, 262–277
 blunt trauma, 247–259
 chest pain, coronary CTA, 219–225
 child abuse, 545–552
 definitions, 37
 epidemiology, 38
 excess radiation, 41
 goals of, 38
 IHPS, 555–565
 inappropriate use, 38
 incidental findings, 37, 42, 43
 lower extremity trauma, 477–494
 medical radiation, 37
 methods, 38
 monetary costs, 41
 pathophysiology, 37
 PE (*see* Pulmonary embolism (PE))
 pelvic fractures, 464
 research, 44
 stroke and SCD, 519–537
 upper extremity injuries, 459
 urinary tract stones, 384
- Emergency radiologists, 31–33
- Emergent imaging, 138, 140
- Emerging advanced imaging, 92, 105, 106
- Endometriosis
 CT and MRI, 428
 definition, 417
 laparoscopy, 427
 prevalence, 418, 419
 serum biomarkers, 428
 transvaginal US, 428
- Endoscopic retrograde cholangiopancreatography (ERCP)
 bile duct obstruction, 317
 choledocholithiasis, 319
- Endoscopic ultrasonography (EUS)
 bile duct obstruction, 317
 choledocholithiasis, 319
- Endovascular revascularization
 acute mesenteric venous thrombosis, 378
 bowel infarction, 377
 mesenteric embolus, 378
 pharmacologic/mechanical thrombectomy, 378
 preoperative clinical evaluation, 377
 Swedish vascular registry, 378
 systemic anticoagulation, 378
 transarterial thrombolysis, 378
- Endovascular thrombectomy (EVT), 91–93, 99–101, 103–107, 109
- Endovascular Treatment for Small Core and Anterior Circulation Proximal Occlusion with Emphasis (ESCAPE), 99
- End-stage renal disease (ESRD), 621
- Epilepsy imaging
 adult refractory, 135
 diagnosis of, 134
 ILAE, 140
 incidence of, 135
 medically refractory, 134
 MESS, 144
 nonfebrile seizures, 146
 subsequent refractory, 134
 syndrome, 134
 temporal lobe, 137
 temporal lobe epilepsy (TLE), 141
 TLE, 141
- Error
 medical literature, 19
 random, 19–23, 26
 systematic, 19
 systemic, 20
- Escherichia coli*, 616, 627
- European Cooperative Acute Stroke Study (ECASS) III trial, 94
- European-wide Multicenter Epilepsy and Single Seizure (MESS), 144
- Evaluation
 evidence classification, 25–26
- Evidence-based imaging (EBI)
 abdominal pain, 307
 acute abdomen, pregnancy, 400–412
 acute musculoskeletal infections, 515
 acute pelvic pain, 416–432
 acute scrotal pain, 435–439
 acute sinusitis, 184–201
 appendicitis, 291
 ARI, 276
 blunt trauma, 257
 chest pain, coronary CTA, 225
 child abuse, 552
 IHPS, 565
 lower extremity trauma, 494
 PE (*see* Pulmonary embolism (PE))
 pelvic fractures, 463–474
 Stroke and SCD and SCD, 537
 upper extremity injuries, 446–459
 urinary tract stones, 383–395
- Evidence-based imaging (EBI) process
 analytic/observational studies, 6
 Bayes' theorem, 14–16
 carotid artery disease, 8
 case-control studies, 6

challenges, 16
 clinical integration, 4
 cost-effectiveness and cost-utility studies, 9, 10
 descriptive studies, 6
 diagnostic tests, 7, 9
 EBI paradigm, 3
 economic analyses in medicine, 10, 11
 evidence, 13–16
 experimental studies, 6
 formulation of clinical question, 5
 GDP, 3
 goal of, 12
 health-care costs, 4
 imaging case studies, 16
 medical literature, 5
 meta-analysis, 11, 12
 patient's expectations, 4
 patient-centered clinical research, 4
 practice, 12
 prospective studies, 7
 qualitative analysis, 11
 quantitative analysis, 11
 readers, 16
 retrospective and prospective, 6
 RVU, 13, 14
 steps, 4
 take-home tables, 16, 17
 type of clinical study, 6
 US health-care system, 4
 Extending the Time for Thrombolysis in Emergency
 Neurological Deficits (EXTEND-IA), 99
 External hernias, 330

F

FAST, *see* Focused assessment with sonography for
 trauma (FAST)
 Femoral diaphyseal fractures, 485
¹⁸F-fluorodeoxyglucose (¹⁸F-FDG), 502
 Fluid attenuation inversion recovery (FLAIR), 54, 55, 59,
 60, 97, 106, 107, 109, 110, 116
 Fluoroscopy, 606, 607, 609, 610
 Fluoroscopy vs. sonography guidance, 576
 Focused assessment with sonography for trauma (FAST)
 CT examinations, 252
 ED evaluation, 252
 emergency ultrasound, 251
 ICU, 252
 mean diagnostic delay, 253
 non-invasive test, 251
 radiation doses, 252
 true-negative findings, 253
 Foreign body aspiration
 AP, 611–613
 chest radiographs, 605
 choking accident, eating, 610
 choking, feeding, 610
 coronal reformats, 612
 definitions, 605, 606
 diagnosis of, 608, 611

epidemiology, 606
 goal, 606
 ingestion of disk battery, 610
 inpatient assessment, 606
 methods, 606
 modality, 610
 pathophysiology, 605, 606
 radiological abnormality, 610
 radiological findings, 608, 609
 research, 610
 suspicion, 610
 Formulating clinical question, 5
 Functional MRI (fMRI), 70, 72, 74, 75, 125

G

Gastrointestinal (GI) bleeding
 acute lower, 357–361
 acute upper, 361–364
 cost to society, 356
 definitions and pathophysiology, 356
 diagnostic performance of imaging modalities, 364
 epidemiology, 356
 imaging, 356
 methodology, 356
 protocol CT angiogram, 365
 Glasgow Coma Scale (GCS), 50–52, 57–60, 66, 67, 69,
 70, 73, 74, 85, 154, 169
 Glenohumeral dislocation
 CT arthrography, 451
 imaging algorithm, 453, 456–458
 MRA, 451
 MRI, 450, 451
 prevalence, 450
 radiographs, 450
 Graphs of conditional probability (GCTs), 332

H

Half-Fourier acquisition single-shot turbo spin-echo
 (HASTE) MRI, 333
 Health information exchange (HIE), 27
 Health information technology (HIT)
 American healthcare system, 28
 application, 27
 ARRA, 28
 CDS, 27
 CPOE, 27
 diagnostic inquiry, 30, 31
 diagnostic procedure, 31
 ED imaging, 29
 EHR, 27
 emergency imaging, 29
 EMRs, 27
 HIE, 27
 HITECH Act, 28
 medical imaging, 29
 methods, 29–30
 NLP and CTRM, 33, 34
 PACS, 27

Health information technology (HIT) (*cont.*)
 physician services, 28
 primary and secondary preventative care, 28
 radiologists, 32, 33
 US healthcare expenditures, 28
 VNAs, 28

Healthy volunteer bias, 24

Helsinki CT score, 59

Hemorrhagic/ruptured ovarian cyst
 appearance, 425
 definition, 417
 diagnostic accuracy of US, 425
 fibrin strands, 425
 indeterminate cysts, 426
 non-pregnant females, 425
 recommendations, 425
 surgical evaluation, 426
 symptoms, 424

Henoch–Schonlein Purpura (HSP), 568, 572, 578, 579

Heterotaxy syndrome, 585, 591, 592

Hirschsprung’s disease, 585, 586

Hydrocephalus, 114, 120, 122, 128

Hydronephrosis, 385–387, 389–395, 616, 622, 624, 625, 628, 629, 632, 635–637

Hyperacute ischemic stroke
 acute intracranial hemorrhage, 94, 95
 after intravenous thrombolysis, 97, 98
 applicability, 98, 106
 CT angiography, 110
 CTA, 91
 definitions, 92
 diagnostic performance, 107
 direct and indirect costs of, 92
 DWI, 91
 endovascular thrombectomy, 98–107
 epidemiology, 92
 hyperacute ischemic stroke, 95–97
 hyperacute stroke MRI, 110
 imaging, 93, 108
 intravenous thrombolysis, 92–98, 107
 methods, 93
 noncontrast head CT, 91, 110
 pathophysiology, 92
 patients, 106, 107
 research, 110
 salvageable tissue, 92

I

Identifying medical literature, 5

IFS, *see* Invasive fungal sinusitis (IFS)

IHPS, *see* Infantile hypertrophic pyloric stenosis (IHPS)

Imaging utilization, 37–39, 42, 44

IMS-3 trial, 100

Incidental findings, 37, 38, 42–44

Infant
 heterotaxy syndrome, 591
 Ladd procedures, 584
 prevalence of, 584
 UGI studies, 589
 volvulus, 584

Infantile hypertrophic pyloric stenosis (IHPS)
 costs to society, 557
 definition, 555
 etiologic factors, 556
 genetic heterogeneity, 556
 imaging, 557
 infants and children
 alkalotic and hypochloremic, 558
 bicarbonate levels, 558
 clinical examination, 558
 clinical palpation, 558, 559
 diagnostic protocols, 563, 564
 gastritis, 557
 imaging algorithm, diagnosis, 562, 563
 management, 561, 562
 non-bilious vomiting, 557, 558
 performance characteristics, diagnostic examinations, 562, 563
 radiography, 564
 renal mechanisms, 557
 serum electrolyte levels, 558
 UGI examination, 559, 560, 562, 564
 ultrasound, 560–564
 vomiting, sonogram, 562, 563
 male/female ratio, 556
 MEDLINE search, 557
 pathophysiology, 556
 pentagastrin infusion, 556
 SNPs, 556
 testosterone levels, 556

Infectious Disease Society of America (IDSA), 268

Infectious Diseases Society of America in conjunction with the American Thoracic Society (IDSA/ATS), 267

Informatics
 integration of, 29

Internal hernia, 330, 336, 340

International Association of Pancreatology and the American Pancreatic Association (IAP/APA), 298

International Headache Society (IHS) criteria, 122

International Mission on Prognosis and Analysis of Clinical Trials (IMPACT), 57, 59, 60

International Registry of Acute Aortic Dissection (IRAD) database, 209

Intestinal malrotation, 584

Intestinal volvulus, 583

Intracerebral hemorrhage (ICH)
 CT
 advantages, 523, 524
 angiography, 525
 blood-brain barrier, 525
 craniectomy, 525
 emergency management, 523
 fluid blood levels, 525
 hematoma volume, 524
 intracranial aneurysm, 524
 ischemic stroke, 524
 mortality rate, 524
 non-contrast, 524
 and ischemic stroke, 534, 535

- MRI**
 acute cerebral infarcts, 525
 angiography, 526
 cavernomas, 526
 deoxyhemoglobin, 526
 gradient-echo sequences, 525
 hyperintense signals, 526
 neuroradiological tool, 526
 randomized trial, 526
 secondary hemorrhages, 526
- Intraluminal obstruction**
 SBO, 331
- Intravenous pyelogram (IVP), 622**
- Intravenous thrombolysis, 91–93, 96–98, 105–108, 110**
- Intravenous urograms (IVUs), 387, 389**
- Intussusception**
 abdominal radiographs, 567
 air enema, 580
 barium, 567
 clinical predictors, 569, 570
 definition, 567–568
 diagnostic imaging, 571, 579
 enema reduction and perforation rates, 573, 574
 epidemiology, 568
 imaging, 569
 imaging studies, 570–572
 irritability, vomiting and intermittent crying, 579
 Lead Point Mass, 579
 methods, 569
 pathophysiology, 567–568
 recurrent cases, 577, 578
 research, 580
 rotavirus vaccine, 568–569
 small bowel, 578–579
 surveys, 569
 therapeutic enema, 572–577
 ultrasound, 579–580
 US, 567
- Invasive coronary angiography (ICA), 221**
- Invasive fungal sinusitis (IFS), 197, 198**
- Ischemic stroke, *see* Stroke, SCD**
- K**
- Knee injuries**
 complex, 484
 meniscal, ligament, chondral, and nondisplaced bone injuries, 484
 OKR, 482, 489
 patellar fractures, 484
 pediatric, 482–484
- L**
- Ladd procedures, 584, 585, 590**
- Large bowel obstruction (LOB)**
 acute colonic pseudo-obstruction, 331
 algorithms, 347
 area under the curve, 346–347
 caecal volvulus, 331
 cost to society, 331
 CT, 341–343
 definition, 331
 epidemiology, 331
 graph of conditional probability, 346–347
 imaging, 331, 332
 imaging modalities, 345, 346
 methodology, 332
 MRI, 343, 345
 radiography, 344, 345
 sigmoid and caecal volvulus, 331
 sigmoid volvulus, 331
 water-soluble contrast enema, 343, 344
- Lead time bias, 24**
- Legg-Calvé-Perthes (LCP) disease, 597–600**
- Length time bias, 25**
- Life expectancy at birth, the United States, 3**
- Liquid enema, 567, 570, 572–580**
- Long bone fractures, 484–486**
- Lower extremity trauma**
 ankle and foot injuries, 486–489
 anterior inferior iliac spine (AIIS), 490, 493
 buttock pain, 490, 491
 cost distribution, 478
 CT, 479
 data analysis, 478
 definition, 477
 hip, 480–482
 knee, 482–484
 long bone fractures, 484–486
 Medline search, 478
 MRI, 479–480
 radiographic views, 479
 radiography, 479
 Salter 1 fracture, 490, 493
 sports injury, 490, 492
 ultrasound, 480
- Low-grade astrocytoma, 120**
- Lumbar spine, 160, 161, 164**
See also Thoracic spine
- M**
- Magnetic resonance angiography (MRA), 377, 451**
- Magnetic resonance cholangiopancreatography (MRCP), 321**
 ACC, 312
 bile duct obstruction, 317
 choledocholithiasis, 319
- Magnetic resonance imaging (MRI)**
 LOB, 343, 345
 SBO
 associated bowel ischemia/strangulation, 340
 detection of level and cause, 337
 emergency department, 333–335
- Magnetic resonance urography (MRU), 387**
- Magnetic resonance voiding cystourethrography (MRVCUG), 623**
- Maisonneuve fracture, 485–486**

- Malrotation
 anomalies, 585
 definition, 583–584
 diagnosis/exclusion, 586–588
 different children, 593
 epidemiology, 584, 585
 goals, 585
 heterotaxy syndrome, 591
 imaging protocols, 592
 infants and children, 583
 Ladd procedure, 585
 methods, 585
 midgut volvulus, 591
 normal duodenal–jejunal junction, 592
 normal UGI series, 592
 older child at low risk, 590, 591
 pathophysiology, 583–584
 research, 592
 UGI series, 583, 588–590
 and volvulus, 586
- Manual detorsion, 437–439
- Marshall classification, 58, 66
- Mayer-Rokitansky-Kuster-Hauser (MRKH) syndrome, 428
- MDCT angiography, 80, 82, 83
- Meckel's diverticulum, 585
- Medical literature, 19, 22
- MEDLINE search, 29, 80, 116, 569
- MEDLINE/PubMed (National Library of Medicine, Bethesda, MD), 311
- Methicillin-resistant *Staphylococcus aureus* (MRSA), 498, 500
- Microhemorrhages, 73
- Migraine headaches, 114, 115, 127
- Mild TBI (mTBI), 49–54, 56, 57, 59–61, 66
- Modic changes, 174
- Motor vehicle crash
 cervical spine imaging, 163
 helmeted rider, 87
 initial cervical spine imaging, 163
 restrained driver, 88
 unrestrained passenger, 86
- MR angiography (MRA), 113, 124, 130, 211
- MR cholangiopancreatography (MRCP), 296, 299, 300
- MR spectroscopy (MRS), 70, 71
- MR Urography (MRU), 622–623
- MRCP, *see* MR cholangiopancreatography (MRCP)
- MRI with contrast, 167, 173, 174
- Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands (MR CLEAN) trial, 98
- Multi-detector row CT (MDCT), 113, 124, 376
- Multiplanar reformats (MPRs), 32
- Multiple-detector computed tomography (MDCT)
 assessing ischemia or strangulation in SBO, 341
 complete SBO, 339
 detection of LBO, 342
 detection of level of obstruction in SBO, 336
 detection of SBO, 334
 diagnosis of LBO, 342, 343
 diagnosis of SBO, 332, 333
 high spatial resolution, 336
 imaging modality, 340
 imaging of ischemia, 340
 LBO, 346
 nonoperative management of SBO, 338
 now-obsolete technology, 340
 SBO, 346
 scanners, 333, 337, 342
 sensitivity, 333
 technique, 351, 352
 transition zones, 336
- Murphy's sign, 312
- Mycoplasmal pneumonias, 269
- N**
- N-acetylaspartate to creatine (NAA/Cr) ratio, 126
- Nasopharynx, 184, 187
- National Emergency X-Radiography Utilization Study (NEXUS), 44, 168, 170, 178, 249, 250, 253
- National Health Service (NHS), 267
- National Inpatient Sample Database, 378
- National Institute of Clinical Excellence, 361
- National Institute of Neurological Disorders and Stroke (NINDS) trials, 94
- National Institute of Neurological Disorders and Stroke tPA trial, 98
- National Library of Medicine PubMed search engine, 4
- National Pediatric Database, 85
- Natural language processing (NLP), 28, 33, 34
- N*-butyl-2-cyanoacrylate, 361
- Necrotizing fasciitis (NF)
 CT, 507
 definition, 499
 diagnosis, 506
 diagnostic performance characteristics, 508
 left ankle, 511, 513
 malaise, fever, hypotension, and progressive cellulitis, 512, 514
 MRI, 506, 507
 radiography, 506
 respiratory distress and shock, cardiac arrest, 514
 risk factors, 499
 surgical evaluation, 499
 ultrasound, 506
- Negative likelihood ratio (NLR), 15
- Negative predictive value (NPV), 8, 15
- Negative-contrast CT cholangiopancreatography (nCTCP), 317
- Neonatal hypothyroidism, 406
- Neonatal osteomyelitis, 498
- Neoplasms, 114–117, 119, 122
- Neurocysticercosis, 139
- Newborn, 583, 584, 586
- NEXUS, *see* National Emergency X-Radiography Utilization Study (NEXUS)
- NF, *see* Necrotizing fasciitis (NF)

- Non-contrast head CT (NCCT), 51, 52, 55, 61, 91, 94–99, 101, 103, 107
- Non-migraine families, 115
- Nonocclusive mesenteric ischemia (NOMI), 374
- Non-strangulating/simple mechanical obstruction, 331
- Nontraumatic headache (NTH), 118
- Non-traumatic hip pain
- definitions, 597
 - epidemiology, 597–598
 - femoral epiphysis, 600
 - initial diagnostic imaging, 598, 599
 - initial investigation, 599
 - Legg-Calvé-Perthes disease, 600
 - methods, 598
 - MRI, 597, 599
 - neutral and frog leg lateral radiographs, 600
 - pathophysiology, 597
 - radiographs, 597
 - research, 600–602
 - society of, 598
 - transient synovitis, 598, 600
 - transient synovitis/septic arthritis, 600
 - ultrasound, 597
- Nuclear medicine, 623–624
- acute lower GI bleeding, 358–359
 - acute upper GI bleeding, 362–363
- O**
- OAR, *see* Ottawa ankle rules (OAR)
- Obstetric and gynecologic (OB/GYN), 400
- OKR, *see* Ottawa knee rules (OKR)
- Orchiopexy, 439
- Osteochondritis dissecans (OCD), 484
- Osteopenic fractures, 167
- Ottawa ankle rules (OAR), 483, 486, 489, 490
- Ottawa knee rules (OKR), 482, 483, 489
- Ovarian torsion, 407, 408
- P**
- Paraduodenal hernias, 330
- Pathologic lead point (PLP), 568, 571, 572, 575, 578, 579
- Patient, Intervention, Comparison, Outcome (PICO), 5
- PE, *see* Pulmonary embolism (PE)
- Pediatric accidental traumatic brain injury
- boy with TBI, 75
 - children with TBI, 70
 - CT 3D skull reconstruction, 75
 - CT brain, 75
 - definitions, 65, 66
 - epidemiology, 66
 - girl with TBI, 75
 - hospitalization of children, 66
 - methods, 67
 - motor vehicle accident and mental status, 75
 - MRI brain, 75–76
 - neuroimaging, 66
 - pathophysiology, 65, 66
 - PECARN decision rule, 65
 - pediatric TBI, 70–75
 - research, 76
 - after TBI, 67–70
- Pediatric Emergency Care Applied Research Network (PECARN) guideline, 65, 67, 68, 70, 73, 75, 76
- Pediatric emergency department, 597
- Pediatric hip injuries
- femoral neck fractures, 481
 - pelvic avulsion fractures, 480
 - posterior hip dislocation, 481
 - SCFE, 481
- Pediatric knee injuries
- distal femoral physeal fractures, 482
 - OCD, 484
 - tibial eminence fractures, 484
 - tibial tubercle fractures, 483
 - transverse supracondylar fractures, 482
- Pediatrics, 356
- Pelvic fractures
- anteroposterior pelvic radiograph, 471–473
 - classification systems, 465
 - complications, 464
 - CT cystography, 473
 - CT technique, 470–473
 - definition, 463
 - genitourinary injuries, 468
 - hemorrhage (*see* Pelvic hemorrhage)
 - imaging algorithm, 469, 470
 - mechanisms, 463
 - motor vehicle crash, 470, 474
 - prediction, arterial hemorrhage, 469
 - PubMed search, 464
 - radiography, 464
 - urethral and bladder injuries, 468, 469
- Pelvic hemorrhage
- clinic injury scores, 466
 - contrast extravasation, CT, 466, 467
 - embolization, 465, 466
 - emergency radiology, 466
 - size of, 467, 468
 - Young-Burgess system, 465, 466
- Pelvic inflammatory disease (PID)
- causes, 418
 - costs, treatment, 418
 - C. trachomatis* and *N. gonorrhoeae* infections, 419
 - CT scan, 424
 - definition, 416
 - diagnostic dilemma, 423
 - hepatic capsular enhancement, 424
 - MRI, 424
 - sensitivity rates, 423
 - sexual behavior, 418
 - transabdominal and transvaginal sonography, 423, 424
 - transabdominal US, 423
 - transvaginal US, 424
 - ultrasound features, 423

- Pelvic ring disruptions, 463–465, 470, 473
 Pelvic trauma, *see* Pelvic fractures
 Penetrating atherosclerotic ulcer (PAU), 207, 208, 215, 216
 Penetrating trauma, 81–82
 Penumbra-based methods, 104
 Picture archiving and communications systems (PACS), 27, 32, 33
 PID, *see* Pelvic inflammatory disease (PID)
 Pituitary macroadenoma, 120
 Placental abruption, 407
 Pneumonia
 airspace/alveolar disease, 262
 cause of death, 262
 computed tomography (CT)
 complications, 271, 272
 immunocompromised patients, 268, 272, 273, 275
 sensitivity, 270
 cost-effectiveness of diagnostics, 262
 CXR (*see* Chest radiograph (CXR))
 direct and indirect costs, 262
 HIV, fever, and dyspnea, 276
 influenza, 262
 PJP, angioinvasive aspergillosis, and CMV, 276
 radiography, 276
 respiratory syncytial virus, 276
 risk factors, 262
 tree-in-bud, 262
 ultrasound, 274–276
 Positive predictive value (PPV), 8
 Positron emission tomography (PET), 70, 72, 74, 75
 Posterior fossa papilloma, 120
 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA), 376
 Primary and secondary headache, 114
 Provoked seizures, 133
 Pseudokidney sign, 571
 PTFE-covered stents, 364
 PubMed search, 152
 Pulmonary embolism (PE)
 anticoagulation, 242–243
 antiphospholipid antibody syndrome, 243
 chest radiography, 232, 233, 240, 241, 243
 clinical assessment, 230
 clinical outcomes, 229
 cost to society, 229
 CT pulmonary angiography, 233, 234, 239, 240
 D-dimer, 230–232
 deep vein thrombosis, 242–244
 definition, 228
 duplex ultrasound, 234, 235
 echocardiography, 235, 236
 evaluation, diagnostic algorithm, 231
 imaging, 229
 imaging modalities, 240
 incidence, 229
 medial literature, 229
 pathophysiology, 228
 pediatric risk factors, 238, 239
 pediatric vs. adult patients, 239
 pregnant women, 236–238
 risk factors, 228
 venous thromboembolic disease, 228
 V/Q scan, 233
 Pulmonary Embolism Rule-out Criteria (PERC), 230
 Pyelonephritis, 616
- Q**
 Qualitative literature summary, 25–26
 Quality Assessment of Studies of Diagnostic Accuracy included in Systematic Reviews (QUADAS), 376
 Quality-adjusted life year (QALY), 10, 11, 127
- R**
 Radiation safety, 41
 Radiography
 cervical spine, 155, 156, 158
 LOB, 344, 345
 SBO
 emergency department, 333
 Radiology, 550
 Random error, 19, 20
 Randomized control trial (RCT), 25
 Rebleeding rates, 361
 Receiver operating characteristic (ROC) curve, 9
 Red blood cell scintigraphy, 357, 358
 Reducibility, 570
 Relative value unit (RVU), 13, 14
 Renal and bladder ultrasonography (RBUS), 615, 616, 624, 625, 627, 629, 633, 635
 Renal colic, 404, 405
 Retrospective study, 175
 Right lower quadrant (RLQ) pain, 402, 403
 Right upper quadrant (RUQ) pain, 403, 408
 Rotterdam score, 58, 59, 66
- S**
 Scaphoid fractures
 complications, 447
 CT, 448
 MRI, 447, 448
 radiographs, 447
 scintigraphy, 448
 standard practice, 447
 wrist imaging, 455
 SCFE, *see* Slipped capital femoral epiphysis (SCFE)
 Scintigraphy, 438
 Scottish Intercollegiate Guidelines Network, 361
 SDH, *see* Subdural hematoma (SDH)
 Seizures
 adult refractory epilepsy, 135
 cerebral pyogenic abscess, 145
 children with new-onset febrile, 136, 137
 definitions, 134, 135
 economic impact, 135
 emergency neuroimaging, 143, 144

- epidemiology, 135
- febrile, 133
- focal neurological deficit, 133
- imaging, 135
- imaging methods, 133
- lethargy and poor feeding, 145
- mental status changes, headache and fever, 145
- methods, 136
- MRI, 133, 146
- neonatal MRI protocol, 147–148
- noncontrast brain CT protocol, 146
- noncontrast CT, 133
- pathophysiology, 134, 135
- patients with first unprovoked, 137–141
- patients with new-onset seizures, 141–143
- pediatric population, 135
- research, 149
- temporal lobe origin, 145
- Septic arthritis (SA)
 - adults, 505, 506
 - definition, 498, 499
 - diagnostic performance characteristics, 508
 - hip ultrasound, 504, 505
 - imaging algorithm, 509
 - left septic hip, 511, 512
 - left shin cellulitis and abscess, 511, 513
 - MRI, 504, 505
 - pelvic radiography, 504
- Sickle cell disease (SCD)
 - African-American population, 521
 - anemia, O₂ transport, 520
 - childhood mortality and life expectancy, 522
 - clinical symptoms, 521
 - costs, 522, 523
 - definition, 519, 520
 - deoxyhemoglobin, 520
 - HbS allele, 520
 - hemolytic pathway, 520
 - imaging, 523
 - malaria parasite, 522
 - MEDLINE search, 523
 - prevalence, 521 (*see also* Stroke, SCD)
- Sigmoid volvulus, 331
- Simple nucleotide polymorphisms (SNP), 556
- Single exposure
 - tissue reactions, 40
- Single-photon emission computed tomography (SPECT), 51, 56, 70, 72, 74, 171, 223, 631
- Single-photon emission computed tomography (SPECT)-enhanced RBC scintigraphy, 358–359
- Sinus radiography
 - abnormal imaging, 189
 - air-fluid level, 190
 - computed tomography (CT), 189
 - low-dose screening sinus CT, 200
 - mild mucoperiosteal thickening, 189
 - MRI, 201
 - sinus puncture, 188
 - sinusitis evaluation, 200
 - ultrasound and clinical examination, 188
- Sinusitis
 - antibiotics, 185
 - classification, 184
 - costs, 185
 - CT scan, 198
 - definition, 184
 - immunocompromised state, IFS, 197, 198
 - maxillary and ethmoid, 185
 - MEDLINE search, 186
 - prevalence, 184
- Slipped capital femoral epiphysis (SCFE), 481
- Small bowel obstruction (SBO)
 - adhesions, 331
 - algorithms, 347
 - area under the curve, 346–347
 - associated bowel ischemia/strangulation
 - CT, 340, 341
 - MRI, 340, 342
 - closed loop obstruction, 330
 - cost to society, 331
 - CT, 334
 - detection of level and cause
 - CT, 336, 337
 - MRI, 337
 - emergency department
 - CT, 332–333
 - MRI, 333–335
 - radiography, 333
 - ultrasound, 333
 - epidemiology, 331
 - external hernias, 330
 - extrinsic causes, 330
 - graph of conditional probability, 346–347
 - imaging, 331, 332
 - imaging modalities, 345
 - incarceration, 330
 - internal hernia, 330
 - intraluminal obstruction, 331
 - intraluminal pathologies, 330
 - intrinsic causes, 330
 - methodology, 332
 - nonoperative/operative management
 - CT, 338, 339
 - water-soluble contrast follow-through, 337, 338
 - non-strangulating/simple mechanical obstruction, 331
 - nonsurgical treatment, 329
 - paraduodenal hernias, 330
 - single transition zone, 330
 - strangulating, 330
 - transition zone, 330
 - tumours, 331
- Solitaire FR Device versus Best Medical Therapy, 99
- Solitaire with the Intention for Thrombectomy as Primary Endovascular Treatment (SWIFT-PRIME), 99
- Spondylolysis/spondylolisthesis, 167
- SQUIRE studies, 19
- Staphylococcal pneumonias, 269
- Staphylococcus aureus*, 616
- STARD studies, 19

- Strangulating SBO, 330
 Streptococcal pneumonia, 269
 Stroke mimics, 93, 95
 Stroke, SCD
 brain ischemia, 526–529
 cerebrovascular imaging
 CT angiography (CTA), 531, 532
 digital subtraction angiography (DSA), 532, 533
 extracranial vasculature, 531
 large-vessel occlusion, 531
 MR angiography (MRA), 532
 ultrasound techniques, 533
 cumulative risk, 522
 definition, 521
 emergency management, 534
 ICH (*see* Intracerebral hemorrhage (ICH))
 imaging protocols
 anatomical MRI of brain, 535, 536
 diffusion brain MRI, 536
 MR angiography, 536
 perfusion brain MRI, 536, 537
 TCCD, 537
 TCD, 537
 mental status, 521
 prevalence, 522
 recurrent stroke, 529–531
 silent infarcts, 521, 535, 536
 symptoms, 521, 533
 Subarachnoid hemorrhage (SAH), 117
 Subdural hematoma (SDH), 546, 547
 Subdural hemorrhage, 120
 Superior mesenteric artery (SMA), 375
 Superselective microcoil embolization, 363
 Susceptibility weighted imaging (SWI), 72–76
 Suspected blunt spine trauma, 153
 Systematic error, 19, 20, 22, 26
- T**
- ^{99m}Tc-hexamethylpropyleneamine oxime (^{99m}Tc-HMPAO), 501
 Tc-99m dimercaptosuccinic acid (Tc-99m DMSA), 621–623, 628, 631, 633, 635
 Tc-99m mercaptoacetyl triglycine (Tc-99m MAG3), 623, 624, 628
 Technetium-99m-labeled RBC scintigraphy, 368
 Temporal lobe epilepsy (TLE), 141, 144
 Tenecteplase, 100
 Testicular torsion
 clinical presentation and physical examination, 437
 color Doppler ultrasound, 438
 costs, 436
 definition, 436
 manual detorsion, 438, 439
 MRI, 438
 power Doppler ultrasound, 438
 predictors, 437
 scintigraphy, 438
 ultrasound, 437–439
- Thoracic endovascular aortic repair (TEVAR), 213, 215, 216
 Thoracic spine
 after blunt trauma
 multicenter study, 161
 multi-trauma patients, 161
 observational studies, 161
 prediction rules, 160
 single-center study, 161
 thoracolumbar imaging, 161
 blunt trauma
 children, 162
 CT protocols, 161
 evidence trial, 162
 Thoracolumbar spine, 151, 160–162, 164
 Thoracolumbar spine imaging, 162
 Thrombolytic candidacy, 94, 95, 100
 Tibia fractures, 485
 Transcranial color-coded Doppler (TCCD), 533, 537
 Transcranial Doppler (TCD), 102, 533, 537
 Transesophageal echocardiography (TEE), 210
 Transient synovitis, 597–600, 602
 Transjugular intrahepatic portosystemic shunt (TIPSS), 363, 364
 Trans-mesenteric hernia, 330
 Traumatic axonal injury (TAI), 49, 51, 53, 55, 59, 60
 Traumatic brain injury (TBI)
 acute CT findings, 52
 acute imaging, 52
 acute intracranial lesion, 53
 advantages, CT and MRI, 60
 African-American ethnicity, 53
 brain MRI, 49
 CCTH, 52
 comprehensive predictive rules, 52
 contusion, 53
 cross-sectional imaging, 56
 definitions, 50
 DTI for individual patient, 54
 DWI and ADC, 55
 economic of, 50
 epidemiology, 50
 GRE and SWI, 55
 GRE sequence, 64
 imaging, 51
 intracranial abnormalities, 51
 methods, 51
 mild TBI, 51, 53
 MRI, 53
 MRI vs. CT, 54
 NCCT, 52
 NEXUS-II, 53
 NOC and CCTH, 52
 non-contrast head CT, 49
 pathophysiology, 50
 patients, 51
 postmortem MRI and CT, 54
 pseudonormalization, 54
 research, 60, 61
 skull radiographs, 56

- SPECT and PET imaging, 56
- surgical intervention, 56–60
- surgical lesions, 51
- SWI, 54
- TAI, 55
- traumatic axonal injury, 53
- Traumatic extracranial vascular injury in adults
 - BCVI, 86
 - blunt neck trauma, 82, 83
 - blunt trauma, 85
 - CT detectors, 84
 - CT screening, trauma, 80–82
 - CTA, 84, 85
 - definition, 79
 - digital subtraction angiography, 79
 - epidemiology, 79, 80
 - imaging, 80
 - MDCT, 79
 - methods, 80
 - motor vehicle accident, 86
 - pathophysiology, 79
 - pediatric patients, 85
 - penetrating neck trauma, 83, 84
 - penetrating trauma, 85
 - research, 86–89
 - screening, 79, 80
 - traumatic neck injury, 84
- Trigeminovascular innervation, 114

- U**
- Ultrasonography (US)
 - AAC, 313, 314
 - ACC, 311, 312
 - bile duct obstruction, 317
 - choledocholithiasis, 318
- Ultrasound
 - SBO, 333
- Underlying infectious disease, 167
- Unprovoked seizures, 134, 135, 137–139, 144, 149
- Upper extremity injuries
 - ACJ separation, 451–453, 456, 458, 460
 - cost of, 446
 - data analysis, 446
 - definition, 446
 - elbow dislocation, 448–450
 - glenohumeral dislocation, 450, 451
 - growth plate, 453
 - imaging, 446
 - imaging recommendations, 454
 - radiography, 452
 - Salter-Harris classification system, 453
 - scaphoid fractures, 447, 448
 - search strategy, 447
 - types, 446
 - wrist fractures, 446
- Upper gastrointestinal (UGI)
 - in children, 583
 - fluoroscopic study with barium, 587
 - imaging test of, 583
 - in infant, 592
 - malrotation, 586
 - normal series, 592
 - pediatric UGI guideline, 586
 - reveal malrotation, 588
 - subtle malrotation, 588
- Upper GI hemorrhage, 367
- Ureteropelvic junction, 622, 636
- Ureterovesicular junction, 622
- Urinary tract infections (UTIs)
 - AAP guideline, 615
 - abdominal radiographs, 622
 - acute episode, 615
 - adolescent boy, 629, 631
 - bilateral hydronephrosis, 635, 636
 - children, 619–620
 - contradictory study, 615
 - costs, 384
 - CT and MRI/MRU, 622–623
 - definitions, 616, 617
 - diagnosis and management, 624–626
 - diabetic male, 391, 393, 394
 - DMSA test, 616
 - epidemiology, 617
 - febrile UTI, 629
 - female infant/child, 628
 - fetal hydronephrosis, 628, 629
 - goals, 618
 - infants and children, 618–621
 - initial febrile UTI, 630
 - IVP, 622
 - lower
 - CT scan, 388
 - immunocompromised/diabetic patients, 387
 - MRI, 388
 - patient symptoms and urinalysis, 387
 - radiography, 387, 388
 - ultrasound, 388
 - male infant/child, 626–628
 - methods, 618
 - nuclear medicine, 623, 624
 - pathophysiology, 616, 617
 - postnatal ultrasound views, 637
 - prophylactic antibiotics, 629
 - pyelonephritis, flank pain, 393, 394
 - RBUS, 616
 - recurrent febrile UTI, 630
 - research, 635
 - risk factors, 384
 - setting, 621–624
 - sonography, 622
 - ultrasound, 616
 - upper
 - CT scan, 390, 391
 - MRI, 391
 - radiography, 389
 - ultrasound, 389, 390
 - upper pole hydroureteronephrosis, 633
 - urethral anatomy in boys, 632
 - VUR, 624, 635

Urinary tract stones

- computed tomography (CT), 386, 387
 - costs, 384
 - definition, 383
 - hydronephrosis, pregnancy, 391, 393
 - imaging protocols, 394, 395
 - MRI, 387
 - obstructive uropathy and pyelonephritis, 391, 392
 - Ovid and PubMed search engines, 385
 - prevalence, 384
 - radiography, 385
 - risk factors, 383
 - ultrasound, 385, 386
 - UTIs (*see* Urinary tract infections (UTIs))
- Uroepithelium, 616

V

- Vendor neutral archives (VNAs), 28, 32, 33
- Ventilation-perfusion scintigraphy (V/Q scan), 232–234, 242–244
- Vertebral injury, 85
- Vesicoureteral reflux (VUR), 615–629, 631–633, 635–637
- Voiding cystourethrography (VCUG), 616–618, 623–629, 631–633, 635–637
- Volvulus, 330, 331, 336, 338, 342–346, 351

W

- Water-soluble contrast enema
 - LOB, 343, 344
- Water-soluble contrast follow-through, 338
- SBO, 337, 338
- White blood cell (WBC) scintigraphy, 502