

Index

A

- Achilles tendon allograft, 290, 308, 321
- Acute delayed infections, 181
- Acute hematogenous infection, 181
- Acute postoperative infections, 181
- Anterior cruciate ligament (ACL), 10, 385
- Anterior inferior iliac spine (AIIS), 373, 374
- Anterior tibial artery perforator flaps, 405
- Anterolateral thigh (ALT) flap, 405
- Anteromedial thigh perforator flaps, 405

B

- Bicompartmental UKA (bi-UKA), 347

C

- Cement and screws technique, 387
- Cement dowels, 197, 198
- Chronic periprosthetic joint infections, 182
- Chronic suppressive antibiotic therapy, 183
- Circular hexapod external fixator (CHEF), 162
- Closed incision negative-pressure therapy (ciNPT), 395, 399
- Closed incision negative-pressure therapy dressing, 397
- Computer-assisted surgery (CAS), 157
- Constrained condylar knee (CCK), 206, 260
- Continuous passive motion (CPM), 231

D

- Deep medial collateral ligament (dmMCL), 8, 14
- Deep vein thrombosis (DVT), 122, 123
- Diabetes mellitus, 389
- Distal femoral periprosthetic fractures, 333
- Distal femoral prosthesis
 - clinical results, 281–282
 - patients history, 279
 - physical examination, 279
 - postoperative results, 280
 - radiographs and advanced imaging, 279
 - surgical approach, 279–280
- Distal quadricepsplasty, 373

E

- Extra-articular deformity, 149
 - femoral extra-articular deformity correction with implant
 - clinical results, 168
 - highly constrained TKA, 168
 - intraoperative techniques, 167, 168
 - patient's history, 166
 - physical examination, 166
 - postoperative results, 168
 - preoperative evaluation, 166, 167
 - radiographic evaluation, 166
 - femoral osteotomy and TKA
 - clinical results, 171–173
 - patient history, 169
 - physical examination, 169, 170
 - postoperative results, 171
 - surgical approach, 170–171
 - X-ray, 170
 - implant, correction with, 152–154
 - clinical results, 156, 157
 - full-length standing radiographs, 151, 152
 - implant choice, 154
 - patient's history, 150
 - physical examination, 151
 - postoperative results, 155, 156
 - preoperative planning, 154
 - procedure, 155
 - tricompartamental arthrosis, 151
 - valgus deformity, 151, 153
 - intra-articular correction, 150
 - tibial osteotomy and TKA
 - AP hip to ankle radiograph, 159, 160
 - circular hexapod external fixation, 161–163
 - clinical results, 164
 - knee and ankle radiographs, 159–161
 - patient history, 158
 - physical examination, 158–159
 - postoperative results, 163, 164
 - preoperative problem list, 159, 160

F

- Failed patellofemoral arthroplasty, management of
 - clinical results, 362–364
 - patient's history, 360
 - physical examination, 360
 - postoperative results, 362
 - radiographs and advanced imaging, 360
 - surgical approach, 360–363
- Failed unicondylar arthroplasty
 - lateral unicondylar arthroplasty
 - common failure mechanisms, 355
 - edge loading, 355
 - IT band pain, 356
 - lateral failure modes, 356
 - lateral UKAs, 355
 - maltracking of implants, 355
 - medial compartment osteoarthritis, 355
 - patient's history, 349, 350
 - physical examination, 350
 - popliteus tendonitis, 355
 - postoperative results, 354
 - radiograph and advanced imaging, 350
 - surgical approach, 350, 352, 353
 - tibial/femoral loosening, 355
 - UKAs, outcomes of, 355
 - unicompartmental candidates, 354
 - medial UNI
 - careful cementation, 346
 - clinical results, 347–349
 - femoral condyle preparation, 346
 - femoral drill guide, 346
 - femoral preparation, 346
 - final tibial preparation, 346
 - lateral parapatellar arthrotomy incision, 345
 - patient's history, 344
 - physical examination, 344
 - postoperative results, 347
 - radiographs and advanced imaging, 344
 - tibial plateau, 345
- Fasciocutaneous flaps, 384
- Femoral bone grafting
 - clinical results, 271, 273
 - patient history, 265
 - physical examination, 266
 - postoperative results, 270
 - radiographs and advanced imaging, 266
 - surgical technique
 - operative procedure, 268–270
 - preoperative considerations, 266–268
- Femoral component removal, 195
- Femoral loosening, 343, 355
- Fibular osteotomy, 161
- First-generation implants, 88
- Fixed flexion contracture, 67–68
 - cruciate-retaining femoral component
 - clinical results, 72
 - patient history, 68, 70
 - physical examination, 68, 70
 - postoperative care, 70, 72
 - radiographs and advanced imaging, 68, 70
 - surgical approach, 69, 70
 - posterior stabilized TKA
 - clinical results, 77, 78
 - patient history, 73
 - physical examination, 73
 - postoperative rehabilitation, 77
 - radiographs and advanced imaging, 74
 - surgical approach, 74–77
- Fixed flexion contracture (FFC), 77–79
- Fixed valgus knee
 - lateral femoral epicondylar osteotomy
 - clinical results, 38, 41–43
 - patient's history, 37
 - physical examination, 38
 - postoperative results, 38
 - radiograph and advanced imaging, 38
 - surgical approach, 38
 - lateral parapatellar approach (*see* Lateral parapatellar approach (LPA))
 - medial parapatellar approach
 - bone resection, 43–45
 - clinical results, 54–56
 - ligament assessment and balancing, 45–47, 50–54
 - patient's history, 43
 - physical examination, 43
 - postoperative results, 54
 - radiograph and advanced imaging, 43
 - pie-crust technique
 - clinical results, 34–37
 - patient's history, 27
 - physical examination, 27
 - postoperative photograph, 33
 - radiographs and advanced imaging, 28
 - surgical technique, 28–36
- Flexion instability
 - complete revision
 - clinical results, 212–213
 - diagnosis, 209–210
 - patient's history, 208
 - physical examination, 208, 209
 - postoperative results, 211
 - surgical approach, 210–211
 - etiology of, 203
 - management of, 203
 - tibial component revision
 - clinical results, 206–207
 - patient's history, 204
 - physical examination, 204, 205
 - postoperative results, 206
 - radiographs and advanced imaging, 205
 - surgical approach, 205–206
- Free flaps, 406, 407
- Freestyle perforator flaps, 405

G

- Gastrocnemius muscle flap, 403
- Gentamycin, 322

- Gerdy tubercle (GT) osteotomy, 57–60
- Gigli saw, 369, 370
- Global instability
- complete revision
 - clinical results, 224–225
 - implant choice, 223, 224
 - patinets history, 222
 - physical examination, 222
 - pneumatic tourniquet, 223
 - polyethylene insertion, 223
 - postoperative results, 224
 - radiographs, 222
 - tibial component revision
 - clinical results, 219–221
 - patients history, 215
 - physical examinations, 216
 - postoperative results, 218
 - radiographs and advanced imaging, 217
 - surgical approach, 217, 218
- Gracilis muscle flaps, 403
- Gullwing patellar osteotomy for osteolytic deformity
- clinical results, 301–304
 - everted patella, articular side of, 299, 300
 - patient's history, 298
 - physical examination, 299
 - radiograph and advanced imaging, 299
 - surgical technique
 - decision-making, 300
 - mediolateral stability, 301
 - osteotomy, 300
 - postoperative management, 300
 - tibial-first technique, 299
- H**
- Heterotopic ossification (HO), 382
- I**
- Iliotibial band (ITB), 30, 39, 45, 49, 50, 52, 55, 57, 356, 373
- Indirect reduction technique, 332
- Infected total knee arthroplasty (TKA)
- irrigation and debridement
 - clinical results, 181–183
 - dilute betadine and chlorhexidine gluconate, 177
 - equipment, 177
 - patient history, 175
 - physical examination, 176
 - postoperative rehabilitation protocol, 181
 - postoperative results, 181
 - surgical technique, 177–181
 - two-stage procedure, 177
 - vancomycin pending culture results, 177
 - X-rays, 176
 - one-stage revision arthroplasty
 - hardware, removal of, 184
 - patient's history, 183
 - physical examination, 183
 - postoperative course, 187
 - preoperative planning and surgical approach, 183
 - radical debridement, 184–187
 - reimplantation, 187
 - two-stage exchange arthroplasty (*see* Two-stage exchange arthroplasty)
- Interpositional arthroplasty, 289
- Intramedullary nailing fixation, 332, 333
- Isolated tibial polyethylene insert exchange (ITPIE), 217, 220, 221
- K**
- Knee sepsis. *See* Sepsis
- Knitted polypropylene graft
- clinical results, 321
 - patients history, 316
 - physical examination, 316
 - postoperative results, 320
 - radiograph and advanced imaging, 317
 - surgical approach, 317–320
- L**
- Lateral collateral ligament (LCL) complex, 30, 45, 47, 49, 51, 54, 56, 205, 206, 222–224
- Lateral distal quadricepsplasty, 373
- Lateral gastrocnemius rotational flap, 384
- Lateral parapatellar approach (LPA)
- clinical results, 62, 63
 - patient's history, 56
 - physical examination, 56
 - postoperative results, 62
 - radiographs, 56
 - surgical approach
 - arthrotomy and wound closure, 61, 62
 - distal femur and proximal tibia cuts, 60, 61
 - femoral rotation, 61
 - flexion gap balancing, 61
 - GT osteotomy and lateral retinaculum, Z-plasty of, 57–60
 - preoperative planning, 57
- Lateral posterior capsule (LPC), 46, 52
- Lateral unicompartmental knee arthroplasty (UKA), 344, 350, 355, 356
- Latissimus flap, 406
- LCL. *See* Lateral collateral ligament (LCL) complex
- Local fasciocutaneous flaps, 404–406
- Local wound care
- distal femoral bone, 385, 386
 - final X-rays, 387, 390
 - multimodal periarticular injection, 387, 389
 - OrthAlign® device, 385
 - patella tracking, 387, 388
 - patient's history, 384
 - physical examination, 384
 - postoperative wound complications, 388
 - preoperative management

Local wound care (*cont.*)

- intraoperative, 391
- midline skin incision, 391
- minor superficial wound complications, 392
- NPWT, 392
- plastic surgery, 389
- polydioxanone bidirectional barbed suture wound closure, 391
- prophylactic flap coverage, 391
- subcutaneous flaps, 391
- transverse skin incisions, 391
- wound dressings, 392
- Prineo[®] skin closure system, 387, 389
- proximal tibial dissection, 387, 388
- radiographs and advanced imaging, 385
- spacer block and drop rod, 386, 387
- tibia component, 388
- tibial component, 387

Locked plating techniques, 333

M

- Macrostrain, 393
- Medial collateral ligament (MCL), 7, 8, 10, 13, 14, 205, 206, 281, 373, 387
- Medial compartment osteoarthritis, 355
- Medial gastrocnemius muscle flap, 401
- Medial gastrocnemius rotational flap, 384
- Medial parapatellar approach, 184, 195
 - bone resection, 43–45
 - clinical results, 54–56
 - ligament assessment and balancing, 45–47, 50–54
 - patient's history, 43
 - physical examination, 43
 - postoperative results, 54
 - radiograph and advanced imaging, 43
 - varus knee, 8
- Medial quadricepsplasty, 373
- Medial unicompartmental knee arthroplasty (UKA), 345, 356
- Metaphyseal bone loss, 259
- Methicillin-resistant *Staphylococcus aureus* (MRSA), 189
- Microstrain, 393
- Microvascular free flap, 406
- Mid-vastus approach, 16

N

- Negative-pressure dressing therapy (NPT)
 - macrostrain, 393
 - microstrain, 393
 - patient's history, 393
 - physical examination, 393, 394
 - postoperative radiographs, 397–399
 - radiographs and laboratory data, 394
 - surgical approach, 394–397
- Negative-pressure therapy with instillation (NPTi), 395, 398
- Negative-pressure wound therapy (NPWT), 383, 391, 392

O

- OA. *See* Osteoarthritis (OA)
- Obesity. *See* Super obese knee
- One-stage exchange arthroplasty, 142
- Open reduction internal fixation (ORIF)
 - clinical results, 332–334
 - distal femoral replacement, 339
 - patient's history, 329–330
 - physical examination, 330
 - postoperative results, 331
 - radiograph and advanced imaging, 330
 - surgical approach, 330, 331
- OrthAlign navigation system, 353
- Osteoarthritis (OA), 1, 344, 345, 347–349, 354, 356
- Osteolytic deformity, gullwing patellar osteotomy for articular side of everted patella, 299, 300
 - clinical results, 301–304
 - patient's history, 298
 - physical examination, 299
 - radiograph and advanced imaging, 299
 - surgical technique
 - decision-making, 300
 - mediolateral stability, 301
 - osteotomy, 300
 - postoperative management, 300
 - tibial-first technique, 299
- Oxford Knee Score (OKS), 344, 347

P

- Parallel to tibial cut technique, 4, 6
- Partial knee arthroplasty, 343, 344
- Patella bone loss, 285, 286
- Patella tendon ruptures
 - allograft survival rate, 326
 - AORI type 3 defect, 326
 - complication, 308
 - extensor mechanism allograft
 - allograft preparation, 311
 - clinical results, 314, 315
 - freeze-dried allografts, 310
 - host distal quadriceps preparation, 313, 314
 - knee, exposure of, 310, 311
 - patient's history, 308
 - physical examination, 308, 309
 - postoperative protocol and rehabilitation, 314
 - proximal tibial trough preparation, 311–313
 - radiographs and advanced imaging, 309, 310
 - specimen, 310
 - tensioning and closure, 313, 314
 - TKA component revision, 311
 - knitted polypropylene graft, extensor mechanism deficiency
 - clinical results, 321
 - patients history, 316
 - physical examination, 316
 - postoperative results, 320
 - radiograph and advanced imaging, 317
 - surgical approach, 317–320
 - structural allograft reconstruction, 326

- structural proximal tibia and patellar tendon
 - patient's history, 321, 322
 - physical examination, 321, 322
 - postoperative results, 324
 - radiograph and advanced imaging, 322
 - surgical exposure, 322–324
- tibial fixation, 326
- Patellar augments
 - prior patellectomy
 - clinical results, 297, 298
 - patient's history, 295
 - postoperative results, 296
 - radiographs and advanced imaging, 296
 - surgical approach, 296
 - TM implant
 - clinical results, 295
 - patient's history, 291
 - postoperative results, 295
 - radiographs and advanced imaging, 291
 - surgical approach, 291–293
- Patellar bone grafting
 - clinical results, 289, 290
 - patient's history, 286
 - physical examination, 287
 - postoperative results, 288
 - radiographs and advanced imaging, 287
 - surgical approach, 287, 288
- Patellar component removal, 196
- Patellar tracking, 87, 287
- Patellofemoral arthritis (PFOA), 362
 - failed metal patella replacement with erosion, 82, 83
- PFA
 - clinical results, 88–91
 - magnetic resonance imaging, 85
 - patient history, 83
 - physical examination, 83
 - postoperative results, 88
 - preoperative radiographs, 83
 - surgical approach, 85–88
 - surgical treatment, 81
 - for symptomatic patient, 81
- TKA
 - clinical results, 93–97
 - patient's history, 91
 - physical examination, 91
 - postoperative results, 93
 - radiograph, 91
 - surgical approach, 91–94
- Patellofemoral arthroplasty (PFA), 81, 82, 96, 97, 359, 363, 364
 - clinical results, 88–91
 - magnetic resonance imaging, 85
 - patient history, 83
 - physical examination, 83
 - postoperative results, 88
 - preoperative radiographs, 83
 - surgical approach, 85–88
- Patellofemoral instability, 219
- PCL. *See* Posterior cruciate ligament (PCL)
- Pedicle hemi-soleus muscle flap, 401
- Pedicle muscle flaps, 402–404
- Pedicle vastus lateralis myocutaneous flap, 404
- Perforator-based propeller flap, 405
- Periprosthetic femur fractures, 329
 - distal femoral replacement
 - clinical results, 337–339
 - patient's history, 335
 - physical examination, 335
 - postoperative results, 337
 - radiograph and advanced imaging, 335
 - surgical approach, 335–337
 - ORIF
 - clinical results, 332–334
 - patient, 330
 - physical examination, 330
 - postoperative results, 331
 - radiograph and advanced imaging, 330
 - surgical approach, 330, 331
- Periprosthetic infections, 181
- Periprosthetic joint infection (PJI), 142, 144, 145, 181–183, 194, 394
- Peroneal artery perforator flaps, 404
- PFA. *See* Patellofemoral arthroplasty (PFA)
- PFC SIGMA posterior-stabilized (PS) implant, 3
- Polyethylene (PE) exchange, 213
- Popliteus tendonitis, 355
- Posterior cruciate ligament (PCL), 10, 14, 16, 21, 51, 53, 68, 69, 209, 210, 212, 218, 231, 386
- Posterior stabilized (PS) implants, 207, 209, 212, 213
- Prineo® skin closure system, 387, 389, 391
- Prior knee fusion, management of
 - AHS, rectus femoris off of, 373
 - clinical results, 379, 380, 382
 - closure, 378
 - collateral ligaments repair, 378
 - distal quadricepsplasty, 373
 - general anesthesia, 368, 370, 374
 - incision, 368
 - insert drains, 378
 - ITB release, 373
 - knee replacement, 378
 - lateral collateral ligament, 378
 - lateral distal quadricepsplasty, 373
 - MCL release, 373
 - medial quadricepsplasty, 373
 - osteotomy of, 369
 - patella and patellar tendon, 378
 - patient's history, 367
 - physical examination, 367, 368
 - posterior release, 378
 - postoperative care, 378, 379
 - postoperative results, 379, 380
 - proximal medial quadricepsplasty, 378
 - proximal quadriceps, release of, 373
 - radiographs and advanced imaging, 368, 369
 - retinaculoplasty with ITB, 378
 - second incision, 373
- Proximal medial quadricepsplasty, 378

R

- Radical debridement, 184
- Reconstructive ladder, 401, 403
- Reduction osteotomy, 387, 388
- Retained hardware
 - complexed TKA, partial removal in
 - clinical results, 103–106
 - patient history, 102
 - physical examination, 102
 - postoperative results, 103
 - preoperative radiographs, 102
 - surgical approach, 102, 103
 - hardware removal
 - clinical results, 111–112
 - patient's history, 106, 107
 - physical examination, 107, 108
 - postoperative results, 111
 - surgical approach, 108–111
 - patient's history, 101
 - preoperative planning, 101
 - PSI, 101, 102
- Robotic technology, 344

S

- Sartorius muscle flaps, 403
- Second-generation patellofemoral arthroplasty implants, 89–90
- Segmental proximal tibial allograft, 326
- Semitendinosus tendon, 307
- Sepsis, 133
 - IV antibiotics, course of, 134
 - one-stage procedure
 - AP radiographs, 135, 136
 - clinical results, 140, 141
 - lateral radiographs, 135, 136
 - patient's history, 134
 - physical examination, 134
 - postoperative results, 140
 - preoperative planning, 135, 136
 - surgical approach, 136–140
 - two-stage revision arthroplasty
 - clinical results, 144–146
 - knee amputation, 143
 - modular components, exchange of, 142
 - one-stage exchange arthroplasty, 142
 - patient's history, 141
 - physical examination, 142
 - postoperative results, 144
 - radiograph and advanced imaging, 142
 - recurrent aspiration and antibiotic suppression, 142
 - stage 1 postoperative radiographs, 143, 144
 - stage 2 postoperative radiographs, 144, 145
 - two-stage exchange arthroplasty, 142
- Severe femoral bone loss
 - Anderson Orthopaedic Research Institute classification, 265, 266
 - distal femoral prosthesis (*see* Distal femoral prosthesis)

- femoral augments and cones
 - clinical results, 277–278
 - patient's history, 273
 - physical examination, 273
 - postoperative results, 276, 277
 - radiograph and advanced imaging, 273
 - surgical approach, 273–276
- femoral bone grafting (*see* Femoral bone grafting)
- sMCL. *See* Superficial medial collateral ligament (sMCL)
- Soft tissue defects management, 392
 - fasciocutaneous flaps, 384
 - flap reconstruction
 - free flaps, 406, 407
 - local fasciocutaneous flaps, 404–406
 - new flap options, 403
 - patient's history, 400
 - pedicled muscle flaps, 403, 404
 - physical examination, 400
 - postoperative results, 401, 402
 - radiographs and advanced imaging, 400
 - reconstructive escalator, 402
 - surgical approach, 400, 401
 - treatment algorithm, 401–403
 - lateral gastrocnemius rotational flap, 384
 - local wound care
 - distal femoral bone, 385, 386
 - final X-rays, 387, 390
 - multimodal periarticular injection, 387, 389
 - OrthAlign® device, 385
 - patella tracking, 387, 388
 - patient's history, 384
 - physical examination, 384
 - postoperative wound complications, 388
 - preoperative management, 389, 391, 392
 - Prineo® skin closure system, 387, 389
 - proximal tibial dissection, 387, 388
 - radiographs and advanced imaging, 385
 - spacer block and drop rod, 386, 387
 - tibia component, 388
 - tibial component, 387
 - medial gastrocnemius rotational flap, 384
 - NPT (*see* Negative-pressure dressing therapy (NPT))
 - proximal wound dehiscence and delayed healing, 383, 384
 - split-thickness skin grafts, 384
- Split-thickness skin grafts, 384, 405
- Stiff knees, 238
 - dynamic splinting, 228
 - ROM, 227, 228
 - stiff total knee arthroplasty
 - clinical results, 236, 238
 - patient's history, 233, 234
 - physical examination, 234
 - postoperative results, 236
 - radiographs and advanced imaging, 235
 - surgical approach, 235, 236
 - tibial component revision
 - AP radiograph, 228
 - clinical results, 232, 233

- lateral radiograph, 228
 - merchant patellar radiographs, 229
 - patient's history, 228
 - physical examination, 228
 - postoperative radiographs, 232
 - surgical approach, 229–232
 - Structural allograft reconstruction, 244
 - Subcutaneous flaps, 391
 - Sub-meniscal arthrotomy, 107
 - Super obese knee
 - conventional instrumentation, surgical technique with
 - AP radiograph of left knee, 126
 - clinical results, 129–130
 - left knee, lateral radiograph of, 126
 - patient's history, 124–125
 - physical examination, 125
 - postoperative radiographs, 129, 130
 - surgical technique, 126–130
 - specific instrumentation, surgical technique with
 - clinical results, 119–124
 - magnetic resonance image, 116
 - patient's history, 115
 - patient-specific instrumentation, 117–119
 - physical examination, 116
 - postoperative results, 119, 122
 - preoperative plan, 117
 - surgical technique, 117–121
 - X-ray, 116, 117
 - Superficial medial collateral ligament (sMCL), 3, 5, 7, 11, 14
 - Superior lateral genicular artery (SLGA) perforator flap, 405
 - Supracondylar periprosthetic fractures, 332, 333
- T**
- Thoracodorsal artery perforator (TDAP) flap, 406
 - Tibial bone loss, management of
 - Anderson Orthopaedic Research Institute
 - classification, 241, 242
 - proximal tibial prosthesis
 - clinical results, 258–260
 - patient's history, 255
 - physical examination, 256
 - postoperative results, 258
 - radiograph, 256
 - surgical approach, 256–258
 - tibial augments and cones
 - clinical outcomes, 251
 - clinical results, 252–255
 - physical examination, 250
 - postoperative results, 251, 252
 - radiograph and advanced imaging, 250
 - surgical approach, 251
 - tibial bone graft
 - clinical results, 247–250
 - CT scan, 242
 - patient's history, 242
 - physical examination, 242
 - postoperative results, 246, 247
 - radiographs, 242
 - surgical approach, 242–246
 - Tibial component revision
 - for flexion instability
 - clinical results, 206–207
 - patient's history, 204
 - physical examination, 204, 205
 - postoperative results, 206
 - radiographs and advanced imaging, 205
 - surgical approach, 205–206
 - global instability
 - clinical results, 219–221
 - patient's history, 215
 - physical examinations, 216
 - postoperative results, 218
 - radiographs and advanced imaging, 217
 - surgical approach, 217, 218
 - stiff knees
 - AP radiograph, 228
 - clinical results, 232, 233
 - lateral radiograph, 228
 - merchant patellar radiographs, 229
 - patient's history, 228
 - physical examination, 228
 - postoperative radiographs, 232
 - surgical approach, 229–232
 - Tibial loosening, 355
 - Tibial polyethylene trial, 197
 - Tibial tubercle avulsion, 307
 - Tibial tubercle osteotomy (TTO), 62, 63
 - Tobramycin, 197
 - Total knee replacement (TKR), 142, 204, 353
 - Total knee revisions (TKRs), 389
 - Trabecular metal (TM) cone, 258, 292, 296, 323
 - clinical results, 295
 - components, 297
 - patients history, 291
 - patellar components, 291, 293–295
 - patellar reconstruction, 297
 - postoperative results, 295
 - radiographs and advanced imaging, 291
 - surgical approach, 291–293
 - Two-stage exchange arthroplasty, 142, 187, 188
 - with mobile spacer
 - bone loss and articulating spacer construction, 196, 197
 - cement dowels, 197
 - clinical results, 199, 200
 - closure, 199
 - debridement, 196
 - diagnostic evaluation, 193
 - femoral component removal, 195
 - implantation, 197
 - median parapatellar approach, 195
 - patellar component removal, 196
 - patient's history, 193
 - physical examination, 193
 - polyethylene insert removal, 195
 - postoperative protocols and rehabilitation, 199
 - radiograph, 193–195
 - tibial component removal, 196

Two-stage exchange arthroplasty (*cont.*)

- sepsis, 146
- static spacer
 - clinical results, 191–192
 - first-stage spacer insertion, 190–191
 - patient's history, 189
 - physical examination, 189
 - postoperative outcomes, 191
 - radiographs and advanced imaging, 189

U

Ultra-high-molecular-weight polyethylene (UHMWPE)
component, 292, 293

Unicompartmental knee arthroplasty (UKA), 249, 250,
256, 344, 347, 348, 355

V

Vacuum-assisted closure (VAC) therapy, 402

Valgus deformity, 67, 151, 153, 159

Vancomycin, 197, 322

Varus knee

- flexion contracture to native alignment
 - clinical results, 23
 - patient's history, 14, 15
 - physical examination, 15
 - postoperative results, 21, 23
 - radiograph and imaging, 15
 - surgical approach, 16–21
- medial collateral ligament pie crusting
 - clinical results, 5–7
 - electrocautery, 3
 - extension gap, 3, 4

- intramedullary femoral alignment guide, 3
- parallel to tibial cut technique, 4, 6
- patient's history, 2
- pes anserinus insertion, 3
- PFC SIGMA posterior-stabilized implant, 3
- physical examination, 2
- postoperative results, 5
- radiographs, 2
- Ran-sall maneuver, 3
- sMCL, 3, 5
- spinal anesthesia, 3
- thick tibial cut, 3
- medial release of
 - clinical results, 12–14
 - curved osteotome, 8
 - distal femur, 10
 - extramedullary guide, 10
 - final implants, cementing of, 12, 13
 - gap-balance device, 11
 - medial parapatellar approach, 8
 - patient's history, 8
 - physical examination, 8
 - posterior Meckel retractor, 10
 - postoperative results, 12
 - radiographs and imaging, 8
 - spacer block, 10–12
 - trial implants, 12
- Varus-valgus constrained (VVC) trial, 224
- Vastus sparing approach, 369

W

Wound complications, 383, 384