

Index

A

- Aaron's sign, 12
- Abdominal compartment syndrome (ACS), 553
 - etiology pathway, 774
 - hemodynamic shifts, 773
 - pathophysiology, 772
- Abdominal distention, 117, 169, 174, 185, 188, 219, 228, 235, 273, 279, 288, 296, 300, 304, 328, 348, 367, 392, 399, 508, 517, 576, 640, 642, 677, 780
- Abdominal packing, 466, 511
- Abdominal pregnancy
 - classification, 608
 - diagnosis, 609
 - laboratory tests, 610
 - MRI, 610
 - transabdominal ultrasound, 610
 - fetal outcome, 615
 - historical perspective, 608
 - incidence, 608
 - laparotomy, 612
 - management, 611
 - maternal outcome, 615
 - non-tubal ectopic pregnancy, 590
 - risk factors, 609
 - surgical treatment
 - operative procedure, 612–614
 - perioperative embolization, 612
 - postoperative management, 614–615
- Abdominal trauma, 413, 473
 - blunt (*see* Blunt abdominal trauma)
 - cardiopulmonary resuscitation, 420–421
 - external risk factors, 416
 - fetal physiology, 419
 - hemodynamic changes, 420
 - incidence, 413–414
 - major hemorrhage, 421–422
 - maternal risk factors, 414–416
 - maternal physiology, 416–419
 - penetrating trauma (*see* Penetrating trauma)
 - pregnancy on trauma mortality, 420
 - prehospital management, 422–424
 - prevention, 416
 - uteroplacental interface, 417
- Abdominal wall endometrioma, 229, 230
- N-acetylcysteine, 807
- Acute appendicitis
 - abdominal CT and MRI, 20–23
 - age, 5
 - Alvarado score, 17
 - anatomic/physiological changes, 7, 8
 - appendiceal carcinoid, 36, 37
 - appendiceal endometriosis/deciduosis
 - clinical presentation, 35, 36
 - diagnosis, 36
 - historical consideration, 35
 - incidence, 35
 - prognosis, 36
 - risk factors, 35
 - third trimester of pregnancy, 29
 - assisted reproductive technique
 - IVF-ET techniques, 33
 - OHSS, 34
 - ruptured HP/EP, 33
 - unruptured HP/EP, 34
 - conservative treatment, 24, 37, 38
 - diagnosis, 16
 - diagnostic algorithm, 23
 - differential diagnosis
 - Fitz-Hugh-Curtis syndrome, 14
 - Meckel's Diverticulitis, 13
 - metritis, 14
 - pelvic thrombophlebitis/ovarian vein syndrome, 14–15
 - round ligament pain, 12–13
 - stump appendicitis, 13
 - urolithiasis/urinary tract infection, 13
 - vomiting of pregnancy, 13–14
 - ectopic/heterotopic pregnancies, 32, 33
 - female sex hormones, 8
 - fetal outcome
 - fetal morbidity, 41
 - fetal mortality, 40
 - historical perspective, 40
 - laparoscopic to open approach, 43
 - long term outcome, 43
 - medical intervention, 40
 - negative appendectomy, 42
 - open vs. laparoscopic approach, 42, 43
 - historical perspective, 3, 4
 - immunologic changes, 7

- Acute appendicitis (*cont.*)
- incidence, 4
 - incidental Meckel's Diverticulum, 32
 - laboratory findings
 - erythrocyte sedimentation rate, 17
 - leukocytosis, 16
 - leukocyturia and bacteriuria, 17
 - neutrophil granulocytosis with left shift, 16
 - raised C-reactive protein, 17
 - laparoscopic appendectomy
 - pneumoperitoneum, 27
 - trimester, 26, 27
 - trocar positions, 28
 - maternal morbidity, 39, 40
 - maternal mortality, 38–39
 - Meckel's diverticulum, 13
 - medical history
 - anorexia, 10
 - constant abdominal pain, 9
 - fundus height, 9
 - location change, 10
 - nausea, 10
 - postnatal shrinking of uterus, 9
 - metritis, 14
 - modified Alvarado score, 17, 18
 - multiple pregnancy, 5
 - normal appendix
 - appendectomy, 31
 - incidence, 30, 31
 - open appendectomy
 - lower midline vertical incision, 25, 26
 - muscle splitting incision, 25
 - right transrectal/pararectal/paramedian incision, 26
 - pathohistological examination, 29, 30
 - pelvic thrombophlebitis, 14, 15
 - perforation rate, 38
 - perityphlitic abscess, 37
 - physical examination, 11–12
 - postoperative course, 30
 - preterm labor, 28
 - puerperium, 37
 - recurrent chronic appendicitis, 8
 - risk factors, 5
 - sickle cell disease
 - clinical presentation, 35
 - incidence, 34–35
 - prognosis, 35
 - transabdominal ultrasound, 18
 - transvaginal ultrasound, 18–20
 - treatment algorithm, 24
 - trimester, 5–6
 - utero-induced radiation effects, 22
- Acute cholangitis, 78, 83, 84, 86, 89
- Acute cholecystitis/biliary colic
- biliary tract perforations
 - common bile duct, 73
 - diagnosis, 73
 - gallbladder perforations, 72
 - pathogenesis, 73
 - treatment, 73
 - cholelithiasis in diabetic pregnant patient, 72
 - cholelithiasis in IVF pregnancy, 72
 - combined/consecutive operations during same pregnancy, 76
 - conservative treatment
 - antibiotics, 68
 - anticholinergic antispasmodics, 68
 - diet, 68
 - pain management, 68
 - total parenteral nutrition, 68
 - ursodeoxycholic acid, 68
 - costal margin pain, 63
 - endoscopic ultrasound, 64
 - fetal outcome, 77
 - historical perspective, 55
 - hyperemesis gravidarum, 62
 - incidence
 - general population, 55, 56
 - pregnancy and puerperium, 56
 - laboratory findings, 63
 - laproscopic cholecystectomy, 74
 - advantage, 74
 - gasless laparoscopy, 75
 - hospitalization, 75
 - local anesthesia, 75
 - open (Hasson) technique, 74, 75
 - SILS, 75
 - third trimester, 74
 - maternal morbidity, 77
 - maternal mortality, 77
 - medical history, 61
 - MRCP, 64
 - open cholecystectomy, 73
 - operative treatment
 - general population, 68, 69
 - laparoscopic surgery, 72
 - open vs. laparoscopic cholecystectomy, 70–72
 - pregnant population, 69, 70
 - pathogenesis
 - bile acid, 61
 - estrogen, 58–60
 - insulin, 60, 61
 - obesity, 61
 - progesterone, 58
 - percutaneous biliary drainage, 68
 - perihepatitis, 62
 - perioperative management, 75
 - physical examination, 61
 - postpartum period
 - gallbladder disease, 76
 - pneumoperitoneum, 76
 - puerperium, 75
 - risk factors
 - age, 58
 - biliary sludge, 56
 - diabetes mellitus, 57
 - gall bladder volume and function, 58

- multiparity, 56, 57
 - obesity, 57
 - oral contraceptives, 57, 58
- treatment, historical perspective, 65–67
- transabdominal ultrasound, 63, 64
- Acute diaphragmatic fatigue, 248
- Acute fatty liver of pregnancy (AFLP), 79, 80, 102, 105, 109, 110, 118, 124, 128, 139, 145
- Acute inflammatory bowel disease, 345, 364
 - CD (*see* Crohn's disease (CD))
 - UC (*see* Ulcerative colitis (UC))
- Acute neonatal hypocalcemia, 149
- Acute pancreatitis (AP)
 - abdominal CT, 128, 129
 - abdominal MRI and MRCP, 129, 130
 - acute pancreatitis per se, 130, 131
 - AFLP, 105, 109, 110, 139, 145
 - age, 102
 - alcohol abuse, 103, 104, 114, 115
 - fetal outcome, 150
 - maternal outcome, 150
 - preeclampsia/eclampsia, 150
 - amylase/lipase, 126
 - antibiotics, 132
 - biliary, 106, 144
 - causes, 105
 - CRRT, 132, 133
 - diagnosis, 119, 122–124
 - endoscopic retrograde cholangiopancreatography, 129
 - endoscopic ultrasound, 129
 - fetal outcome, 146–148
 - gallstones, 145
 - HELLP syndrome, 145
 - heparin, 136
 - historical perspective, 101, 102
 - hypercalcemia, 131
 - hyperlipidemia/dyslipidemia, 103, 144, 145
 - estrogen therapy, 112, 113
 - hereditary/physiologic, 110
 - hypothyroidism, 114
 - medications, 114
 - hyperlipidemic pancreatitis, 134
 - hypertriglyceridemia
 - fetal outcome, 149, 150
 - maternal outcome, 149
 - incidence, 102
 - insulin, 136
 - intensive care unit, 134
 - lipid-lowering diet, 134, 135
 - lipid-lowering medications, 138
 - liver and pancreatic enzymes, 119–121, 124, 125
 - L-thyroxine, 136
 - maternal outcome, 146
 - medical history
 - AFLP, 118
 - hypertriglyceridemia, 118
 - PHPT, 117, 118
 - medications, 115, 150
 - mode of delivery, 145
 - pancreatic cyst fluid analysis, 130
 - pancreatic neoplasms, 105, 115, 116
 - pancreatic pseudocyst, 104, 140–143
 - pancreatic tumors, 141
 - parathyroid glands imaging, 130
 - parathyroid hormone, 127
 - PHPT, 104
 - physical examination
 - Cullen's sign, 119
 - hyperlipidemia, 120
 - intrapertitoneal fluid, 119
 - lipemia retinalis, 119, 121
 - plasmapheresis, 137
 - post-ERCP, 115
 - preeclampsia/eclampsia, 104, 116, 145
 - preterm labor, 145
 - primary hyperparathyroidism, 139, 141, 144
 - calcium-induced, 109
 - calcium-PTH metabolism in pregnancy, 106, 108, 109
 - fetal outcome, 148, 149
 - hypercalcemia, 109
 - maternal outcome, 148
 - puerperium, 116, 117
 - race, 103
 - Ranson criteria, 127
 - serum calcium and parathormone, 125, 126
 - serum lipids, 124, 125
 - serum markers, 126
 - supportive measures, 131, 132
 - therapeutic delivery, 144
 - total parenteral nutrition, 135, 136
 - transabdominal ultrasound, 127, 128
 - triglycerides, 126, 127
 - trimester, 102
 - Acute placental abruption, 444
 - Acute puerperal salpingitis, 690, 696
 - Acute pyelitis, 718
 - Acute pyelonephritis
 - clinical presentation, 718
 - complications, 718
 - diagnosis, 718
 - differential diagnosis, 718
 - etiopathogenesis, 717–718
 - fetal outcome, 719
 - incidence, 717
 - maternal outcome, 719
 - symptoms, 718
 - treatment, 718–719
 - Acute salpingitis, 689, 692
 - Acute urinary retention (AUR)
 - causes, 721, 722
 - diagnosis, 721
 - incidence, 719
 - pelvic examination, 721
 - postpartum, 721
 - recurrence, 724
 - retroverted, 720

- Adenomyoma of the round ligament, 210
- Adhesion formation, 292
- Adhesions
- chemical agents, 294
 - classification, 292
 - conservative treatment, 295
 - definition, 292
 - diagnostic workup, 295, 296
 - incidence, 292
 - liquid/solid mechanical barriers, 294
 - pathophysiology
 - postoperative adhesion formation, 292–293
 - previous appendectomy, 294
 - uterine enlargement, 293
 - surgical technique, 294
 - surgical treatment, 295
- Adhesive obstruction, 270, 292, 294
- Adjuvant chemotherapy, 584
- Adnexal masses, 545
- classification, 573, 574
 - diagnosis, 576
 - malignancy of, 577
 - optimal management, 579
 - robotic resection, 582
- Adnexal ring sign, 578
- Adnexal torsion (AT)
- abdominal CT, 550
 - abdominal pain, 546
 - abortion rates, 558
 - acute pelvic/abdominal pain, 547
 - adhesion formation, 557
 - adnexal preservation/resection, 551
 - adnexectomy, 552, 555
 - anatomic variations, 545
 - anesthetic and perioperative management, 557
 - and concomitant contralateral ectopic pregnancy, 553
 - definition, 543
 - detorsion/unwinding
 - cardiotocography, 554
 - laparoscopic control, 554
 - single-incision laparoscopic surgery, 554
 - Doppler mapping, 549
 - Doppler studies, 557
 - ectopic pregnancy, 546
 - electrosurgery, 555
 - hemorrhagic infarction, 548
 - hormonal pregnancy support, 557
 - hyperstimulation/pregnancy, 547
 - infundibulopelvic ligament, 546
 - intra-abdominal pressure, 545, 553
 - laboratory indicators, 547, 548
 - laparoscopic detorsion, 552
 - laparoscopic fixation, 558
 - laparoscopy, 551
 - laparotomy, 551
 - Leukocytosis, 548
 - McBurney's gridiron incision, 555
 - MRI, 550
 - normal/pathological hyperfunction, 545
 - obstetric indications, 558
 - oophoropexy, 554–556
 - ovariopexy, 558
 - pathological findings, 557
 - patient evaluation, 543
 - peritoneal irritation, 546
 - preterm labor, 557
 - recurrence, 558
 - recurrent torsion, 546
 - reproductive capability, 543
 - right-sided and left-sided, 547
 - sonographic findings, 548
 - surgical management, 543
 - tachycardia, 547
 - transvaginal ultrasound, 548, 549
 - trocar insertion, 553
 - ultrasound scans, 557
- Advanced Trauma Life Support (ATLS) guidelines, 422
- Airway management, 752–756
- Alder's sign, 11
- Alvarado score, 17
- American Congress of Obstetricians and Gynecologists (ACOG), 599, 628
- Aminosalicylates (FDA B), 352
- Amniotic cavity, 792
- Amniotic fluid embolism, 454, 455
- Anesthetic management
- airway management
 - abdominal trauma, 755–756
 - acute appendicitis, 756
 - complications, 752–753
 - diaphragmatic hernia, 754–755
 - anesthetic drugs
 - classification, 752
 - placental transfer characteristics, 753
 - sedation, 752
 - antibiotics
 - acute appendicitis, 757
 - acute cholecystitis, 757
 - ECMO, 756
 - goals, 757
 - intraoperative CO₂ monitoring, 756
 - pain management, 757–758
 - thromboprophylaxis
 - Cesarean section, 762
 - elective cholecystectomy, 761
 - emergent cholecystectomy, 761–762
 - IBD patients, 762
 - low-risk surgery, 761
 - risk factors for thrombosis, 759
 - torsion of the gravid uterus, 762
 - VTE, 762
- Angiomyolipoma (AML), 726
- Antenatal indomethacin, 807
- Antepartum hemorrhage (APH), 642
- Anthraquinone laxatives, 272
- Antibiotics, 757, 803
- Anticoagulation
- ovarian stimulation, 519
 - PVT, 521
- Appendectomy, 761

Appendiceal carcinoid, 36, 37
 Argon beam electrocoagulation, liver rupture, 512
 Assisted reproductive technology (ART), 517, 545

B

Benign cystic teratoma (BCT), 575
 Beta human chorionic gonadotropin (β HCG) level, 576
 Betamimetics, 809
 Bethesda criteria, 193
 Bicornuate, 664
 Bilateral salpingo-oophorectomy, 309
 Bilateral torsion, 544
 Biliary sludge/microlithiasis, 56
 Bladder injury, 733, 735
 Bleeding, 579
 Blood pressure, 642
 Blunt abdominal trauma, 733

- abdominal X-ray, 456
- acute deceleration injury, 445
- airbag use, 472
- airbag, safety system, 440, 441
- anatomic and physiological changes, 417
- annual crash rate, 426
- automobile crashes, 425
- belted *vs.* unbelted pregnant population, 436–439
- blood loss, 463
- cardiotocographic monitoring, 467
- cardiotocography, 461
- CT scan, 450, 461, 462
- disseminated intravascular coagulation, 470
- Doppler ultrasound, 461
- fatal injuries in seat belt wearers, 431
- fetal death, 419
- fetal evaluation, 463
- fetal loss and placental abruption, 425, 469
- fetal mortality *vs.* infant mortality, 428
- fetal sonography, 468
- focused abdominal sonography for trauma, 457
- gestational age, 470
- hemoglobin and hematocrit levels, 456
- in-hospital mortality rate, 471
- laboratory values, 419
- lap belt approaches, 433
- management algorithm, 464
- maternal hemorrhagic shock, 469
- maternal mortality, 469, 472
- maternal plasma bicarbonate level, 471
- maternal shock, 470
- motor vehicle accident with spleen laceration, 459
- nonstress test, 468
- obstetric complications
 - deceleration injury, 444
 - direct fetal injuries and fractures, 444, 447
 - indirect injury of the fetal viscera, 448
 - nonfatal fetal crash outcomes, 444
 - preterm labor, 447
 - uterine rupture, 448–451
- obstetric sonogram, 467
- pelvic free fluid, 457, 458

perinatal mortality rate, 471
 population-based studies, 428
 postpartum care

- placental abruption, 468, 469
- preterm labor, 469

 pregnancy loss, 428
 prenatal care, 469
 preterm labor, 428
 radiologic procedures, 462, 463
 ruptured uterus, 449
 seat belt

- in automotive use, 431, 433
- in pregnancy, 439
- usage, 438, 472

 severe fetal injury, 428
 social/domestic violence

- fetal loss, 444
- medical attention, 442
- sexual/physical abuse, 443
- subclinical chronic placental disruption, 444
- types, 442
- warning signs, 443

 spleen and liver, 460
 splenic laceration, 459, 460
 stress reactions, 428
 transabdominal sonography, 458
 uterine rupture, 425
 Bochdalek's hernia, 248, 258, 262
 Boerhaave syndrome, 184
 Borderline ovarian tumors, 579
 Braxton-Hicks contractions, 7
 Breastfeeding, 364
 Bryan's sign, 12
 Burst abdomen, 226

- definition, 779
- management, 783

C

Canal of Nuck, 211–214
 Capnography, 756
 Cardiopulmonary resuscitation (CPR), 420
 Carnett's sign, 63
 Cecal volvulus

- clinical presentation, 319
- diagnosis, 319–320
- differential diagnosis, 319
- distal ascending colon, 318
- historical perspective, 318
- incidence, 319
- maternal and fetal outcome, 321
- partial obstruction, 318
- risk factor, 319
- treatment, 320

Certolizumab pegol, 356
 Cervical ripening, 631
 Cervical remodeling, 791
 Cesarean myomectomy, 681
 Cesarean scar endometriosis, 229
 Cesarean scar hernias, 230

- Chilaiditi sign, 255
- Childhood cancers, 477
- Chlamydia trachomatis* infection, 686
- Cholecystitis, 764
- Choledochal cyst rupture, 90
- Choledochoscopy, 83, 84
- Chorioamnionitis, 810
- Cimetidine and ranitidine, 177
- Cirrhosis, 403
- Cisplatin, 307
- Classic cesarean delivery, 628
- Cocaine abuse, 609
- Coffee bean sign, 315
- Colic angulation, 331
- Colon and rectum carcinoma
 - age, 303
 - COX-2 enzymes, 303
 - diagnosis
 - abdominal MRI, 306
 - laboratory findings, 305
 - plain abdominal X-ray, 305
 - transabdominal ultrasound, 305–306
 - differential diagnosis, 304–305
 - elective presentation, 304
 - emergent presentation, 304
 - historical perspective, 302
 - hormonal receptors, 303
 - incidence, 302–303
 - predisposing factors, 303
 - prognosis
 - fetal outcomes, 310
 - maternal outcome, 308–310
 - treatment
 - chemotherapy, 307–308
 - conservative and bridge therapy, 307
 - mode of delivery, 307
 - radiotherapy, 308
 - surgical, 306–307
- Colonic dilation, 366
- Colorectal carcinomas (CRC)
 - acute abdomen, 190
 - elective presentation, 190
 - fetal outcome, 192
 - genetic counseling, 192, 193
 - gestational age, 190
 - historical perspective, 189
 - incidence, 189
 - lateral ovarian transposition, 192
 - maternal outcome, 192
 - medial ovarian transposition, 192
 - one-stage/two-stage procedure, 190, 191
 - ovarian transposition, 191, 192
 - pathogenesis, 189
 - perforation, 192
 - signs and symptoms, 189, 190
- Colostomy, 237, 238
- Colpotomy approach, 688
- Common bile duct (CBD) stones
 - AFLP, 79, 80
 - choledochoscopy, 83, 84
 - choledochotomy, 85
 - clinical presentation, 78
 - complications, 82, 83
 - contrast agents, 82
 - ERCP, 81, 82
 - EUS, 83
 - fetal outcome, 86
 - ICP, 78, 79
 - incidence, 77, 78
 - IOC, 85
 - laboratory findings, 80
 - mandatory cholecystectomy, 84
 - maternal outcome, 86
 - MRCP, 80, 81
 - puerperium, 85, 86
 - radiation, 81
 - risk factors, 78
 - sedation, 82
 - selective (wait-and-see) approach, 84
 - therapeutic delivery, 85
 - transabdominal ultrasound, 80
 - transcystic approach, 85
- Complete uterine rupture, 623
- Component separation technique, 232
- Congenital defects
 - falciform ligament, 326
 - sigmoid mesocolon hernia
 - abdominal MRI, 326
 - classification, 325
 - clinical features, 325
 - CT, 325
 - etiology, 325
 - incidence, 325
 - plain abdominal X-ray, 325
 - prognosis, 326
 - treatment, 326
 - transomental hernia, 326
- Congenital intestinal malrotation
 - clinical presentation, 298
 - diagnosis, 298
 - etiopathogenesis, 298
 - incidence, 298
 - prognosis, 299
 - treatment, 298–299
- Conservative and bridge therapy, 307
- Constipation
 - adverse events, 272
 - causes, 271
 - functional, 271
 - laxatives, 272
 - PEG, 272
 - Rome IV criteria, 271
 - urgency, 271
- Constriction-ring dystocia, pathognomonic signs, 640
- Continuation of pregnancy, pathophysiologic mechanisms, 584
- Continuous renal replacement therapy (CRRT), 132, 133
- Continuous venovenous hemofiltration (CVVH), 132

- Contralateral tubal ectopic pregnancy, 553
 Controlled ovarian hyperstimulation, 545
 Corpus luteum cyst, 574, 576
 Corpus luteum cystectomy, 583
 Corticosteroids, 353
 Costal margin pain, 63
 Crohn's disease (CD), 186
 - clinical presentation, 348
 - diagnosis
 - endoscopy, 350
 - imaging modalities, 349–350
 - laboratory findings, 348–349
 - differential diagnosis, 350
 - fetal outcome
 - breastfeeding, 364
 - inheritance, 362
 - perinatal outcome, 362–364
 - pregnancy, 364
 - historical perspective, 345
 - incidence, 345
 - maternal outcome
 - fertility and sexual health, 361–362
 - maternal morbidity and mortality, 360–361
 - medical treatment
 - antibiotics, 351–352
 - antidiarrheal agent, 357
 - corticosteroids, 353
 - cyclosporine, 354, 355
 - electrolytes, 352
 - enteral nutrition, 357
 - immunomodulators, 355–356
 - medications, 351
 - mesalamine, 352
 - methotrexate, 356
 - olsalazine, 352
 - probiotics, 357
 - sulfasalazine, 352
 - supplements, 352
 - thalidomide, 356–357
 - thiopurines, 353–354
 - topical 5-ASA, 352
 - total parenteral nutrition, 357
 - pathophysiology, 347
 - pregnancy, effect of
 - classification, 346
 - disease activity, 346
 - fibrous tissue, 346
 - higher remission rate, 346
 - maternal immune system, 345–346
 - Montreal classification, 347
 - parity and surgical resections, 347
 - surgical treatment
 - anesthetic and perioperative management, 359
 - elective cesarean section, 359, 360
 - elective surgery, 358
 - emergency surgery, 358–359
 - emergent cesarean section, 360
 - mode of delivery, 359–360
 - preterm labor, 359
- Culdocentesis, 597
- Cullen's sign, 119, 593
 Cyclosporine, 354–355
 Cystic teratoma, 553
 Cytokine-suppressive anti-inflammatory drugs (CSAIDs), 808
- D**
- Damage control surgery, 466, 467
 De novo adhesion formation, 292
 Decidualization, 575
 Deep bowel endometriosis, 186
 Deep venous thrombosis (DVT), 517, 758
 Defensive injuries, 443
 Dermoid cyst, 575, 583
 Diaphragmatic eventration, 248
 Diaphragmatic hernia (DH)
 - abdominal organ operations, 264
 - acute diaphragmatic fatigue, 248
 - asymptomatic patients, 259
 - blunt trauma, 249
 - CDH, 248, 249, 254
 - chest X-ray with peroral contrast, 255–257
 - conservative treatment
 - intravenous fluid, 259
 - nasogastric suction, 259
 - oxygenation, 259
 - diagnosis, 255
 - diaphragmatic defect closure, 263, 264
 - diaphragmatic eventration, 248
 - differential diagnosis, 254, 255
 - emergent presentation, 260
 - fetal outcome, 264
 - hiatal hernias
 - clinical presentation, 251, 252
 - type I (sliding), 251
 - type II (paraesophageal), 251
 - type III, 251
 - type IV, 251
 - historical perspective, 247
 - iatrogenic, 249, 250
 - laparoscopic repair, 262, 263
 - maternal mortality, 264
 - obstetric management, 260
 - open surgery
 - thoracoabdominal approach, 262
 - transabdominal approach, 261, 262
 - transthoracic approach, 260, 261
 - penetrating trauma, 249
 - perioperative management, 260
 - plain chest X-ray, 255, 256
 - quiescent phase, 252
 - recurrence rate, 264
 - spontaneous rupture during labor, 250, 251
 - strangulation, 253
 - symptomatic patients, 259, 260
 - thoracic CT, 257, 258
 - thoracic MRI, 258
 - thoracic ultrasound, 256
 - trauma, 252

- Diastasis recti abdominis (DRA)
 anatomy, 239
 classification, 240
 conservation treatment, 240
 incidence, 240
 surgical treatment, 240
- Diverticulectomy, 32
- Double-circle sign, 282
- Double-filtration technique, 137
- E**
- Eclamptic syndrome, 515
- Ectopic decidual, 35
- Ectopic pregnancy
 causes, 589
 classification, 590–591
 clinical criteria, 591–593
 clinical presentation, 330, 593–594
 CT, 596
 culdocentesis, 597
 definition, 589
 diagnosis, 330, 596, 597, 599
 differential diagnosis, 330, 594
 gynecologic treatment, 330
 hemodynamically unstable, 600
 incidence, 329–330
 MRI, 597
 prevalence, 590
 prognosis, 602
 risk factors, 590
 stable patients, 601
 surgical treatment, 330
 transvaginal ultrasound, 595
 treatment, 599–602
- Elective laparotomy, 581
- Emergency laparotomy, 580
- Emergency medical services (EMS), 423
- Endoluminal decompression, 315–317
- Endoluminal uterine instrumentation, 531
- Endometriomas, 576
- Endometriosis, 693, 700
 bowel perforation sites, 186
 clinical presentation, 187, 188
 definition, 186
 diagnosis, 188
 differential diagnosis, 188
 incidence, 186
 pathophysiology, 186–187
 treatment, 188
- Endosalpingiosis
 anesthetic and perioperative management, 224
 clinical presentation, 223
 definition, 222
 diagnosis, 223
 early postoperative complications, 224
 gravid uterus, 226
 mesh repair, 224
 obstetric management, 224
 pathogenesis, 222, 223
 plain abdominal X-ray, 223
 prognosis, 227
 transabdominal X-ray, 223
 treatment, 223
 umbilical hernioplasty during Cesarean section, 224, 225
 uterine fibroids, 225
- Endoscopic ultrasound (EUS), 83
- Endothelial dysfunction, 775
- Entero-amniotic fistula, 531
- Enterouterine fistula, 530, 531
- Epigastric hernia
 clinical presentation, 227
 diagnosis, 227
 differential diagnosis, 227
 incidence, 227
 management algorithm, 227, 228
 prognosis, 227
- Epilepsy, 811
- Erythrocyte sedimentation rate, 17
- Estrogen:progesterone ratio, 110
- Estrogen receptors (ER), 303
- European Society of Urogenital Radiology (ESUR)
 guidelines, 64, 65
- Extracorporeal membranous oxygenation (ECMO), 756
- Extraluminal stomal compression
 diagnosis, 311–313
 incidence, 311
 prognosis, 313
 treatment, 313
- Extrauterine/abdominal pregnancy, 531, 592
- F**
- Falciform ligament hernia, 326–327
- Fallopian tube pregnancy, 395, 531
- Fallopian tube-sparing surgery, 599
- Falls, nonfatal injury, 442
- Family planning, 442
- Famotidine and nizatidine, 177
- Fecal calprotectin, 349
- Fetal assessment, 419
- Fetal bacteremia, 809
- Fetal death in trauma, 442
- Fetal distress, 394, 641, 784
- Fetal gunshot wounds, 483
- Fetal head injury, 418
- Fetal heart rate monitoring, 763
- Fetal heart tones (FHTs), 465
- Fetal hypoxia, 653
- Fetal inflammatory response syndrome (FIRS), 809
- Fetal physiology, 419
- Fetal trauma exposure, 426
- Fetal-maternal hemorrhage (FMH), 419, 451–454
- Fetoplacental circulation, 419
- Fibroid red degeneration, 677
- Fibronectins, 791
- Fish oil supplements, 352
- Fitz-Hugh-Curtis syndrome, 14, 62
- Flow cytometry, 452

- 5-Fluorouracil (5-FU), 307
 Focal hepatic necrosis, 510
 Folate supplements, 352
 Follicular cysts, 574
 Free intestinal perforation
 pathophysiology, 347
 ulcerative colitis, 366
 Frimann-Dahl's sign, 315
 Functional constipation, 271
 Functional cyst, 576
- G**
- Gallbladder perforations, 72
 Gamma-glutamyl transpeptidase (GGT) levels, 124
 Gasless laparoscopic surgery (GLS), 27, 75, 551, 582
 Gastric banding, 174, 185
 Gastric cancer in pregnancy
 diagnosis, 181, 182
 fetal outcome, 183
 incidence, 180
 Krukenberg tumor, 180, 181
 maternal outcome, 182, 183
 pathology, 180
 pathophysiology, 180, 181
 perforated malignant ulcer, 181
 treatment
 cesarean section, 182
 hormonal therapy, 182
 metastatic tumor, 182
 primary gastric tumor, 182
 Gastric hemorrhage, 505
 Gastric outlet obstruction
 clinical presentation, 327
 diagnosis, 327–328
 embryology, 327
 incidence, 327
 treatment, 328
 Gastrodynia, 505
 Gastrointestinal–genital communications, 529, 530
 abdominal pregnancy, 533
 hemorrhagic shock, 533
 hysterosalpingography, 532
 infected fistula and inflammatory process, 533
 instrumental abortion, 532
 ipsilateral salpingo-oophorectomy, 532
 laparotomy and minimal surgery, 533
 maternal survival, 534
 proctoscopic examination, 532
 rectal bleeding, 532
 uterine fundus, 534
 Gastrointestinal inflammation, 531
 Gastrointestinal perforations, 163, 179, 180, 183, 185
 gastric cancer (*see* Gastric cancer in pregnancy)
 intestine (*see* Intestinal perforation)
 nonpeptic ulcer (*see* Nonpeptic ulcer perforations/
 ruptures)
 peptic ulcer (*see* Peptic ulcer disease (PUD))
 perforated gastric cancer (*see* Perforated gastric
 cancer in pregnancy)
- Genital anomaly, 694
 Gestational sac, 595
 Gravid uterus, 219, 226
 Groin hernia
 causes, 207
 clinical presentation, 206, 207
 etiopathogenesis, 206
 hernia recurrence, 217
 hydrocele of canal of Nuck, 213, 214
 incidence, 206
 inguinal endometriosis
 canal of Nuck, 210, 211
 imaging, 211–213
 incidence, 210
 symptoms, 210
 inguinal herniorrhaphy during cesarean section,
 215–217
 perioperative management, 215
 plain abdominal X-ray, 214
 postherniorrhaphy pain, 217
 pregnancy, 206
 risk factors, 206
 RLVs (*see* Round ligament varicosities (RLVs))
 round ligament stretch/pain, 209, 210
 surgical site/mesh infection, 217
 transabdominal ultrasound, 214
 treatment, 214, 215
- Gynecologic causes
 ectopic pregnancy
 clinical presentation, 330
 diagnosis, 330
 differential diagnosis, 330
 gynecologic treatment, 330
 incidence, 329–330
 surgical treatment, 330
 normal pregnancy
 diagnosis, 332–333
 historical perspective, 330–331
 incidence, 331
 pathophysiology, 331, 332
 treatment, 333
 ovarian teratoma
 clinical presentation, 328–329
 diagnosis, 329
 incidence, 328
 prognosis, 329
 treatment, 329
- H**
- Helicobacter pylori* infection, 170, 171
 HELLP syndrome, 145, 467, 506–508, 776
 Hemorrhage, 506, 592
 Hemorrhagic corpus luteum cyst, 579, 581
 Hemorrhagic ovarian cyst, 578
 Hemorrhagic shock, 653
 Hemothorax, 480
 Hepatic function test, 509
 Hernia of Bochdalek, 248, 249
 Herniation, 321

- Herniorrhaphy, 215
- Heterotopic pancreas (HP)
 - clinical presentation, 327
 - diagnosis, 327–328
 - embryology, 327
 - incidence, 327
 - treatment, 328
- Heterotopic pregnancy, 590
 - diagnosis, 595
 - transabdominal ultrasonography, 595
- High-density lipoprotein (HDL), 59, 60
- Histamine₂ receptor antagonists (H₂RA), 177, 178
- Hofstatter's sign, 594
- Hooking maneuver, 63
- Hyaline degeneration, 676
- Hydronephrosis, 725, 731
- Hyperemesis gravidarum, 62
- Hyperlipidemia/dyslipidemia, 120, 144, 145
- Hypocapnia, 418
- Hypochlorhydria, 165
- Hypovolemia, 420
- Hypoxic–ischemic encephalopathy, 654
- Hysterectomy, 671
- Hysterotomy, 670–671
- I**
- Idiopathic perforation of colon, 193
- Ileocecal intussusception, 279
- Ileosigmoid knotting (ISK)
 - abdominal CT, 301, 302
 - abdominal MRI, 301
 - clinical presentation, 300–301
 - definition, 299
 - diagnosis, 301
 - etiopathogenesis, 300
 - incidence, 300
 - plain abdominal X-rays, 301
 - prognosis, 302
 - risk factors, 300
 - transabdominal ultrasound, 301
 - treatment, 301–302
 - types, 299
- Ileostomy, 237, 238
- In vitro fertilization (IVF), 33, 545
- Incarcerated internal hernia
 - congenital defects (*see* Congenital defects)
 - post-bariatric surgery
 - abdominal CT, 323
 - abdominal MRI, 323, 324
 - classification, 321
 - clinical presentation, 322–323
 - fetal outcome, 325
 - incidence, 321
 - maternal Outcome, 325
 - morbid obesity, 321
 - treatment, 323, 325
- Incomplete (partial) uterine rupture, 622
- Index of wrong diagnosis, 31
- Induced abortion, 444
- Inferior epigastric artery (IEA) bleeding, 525
 - anticoagulant medications, 523
 - blunt trauma, 522
 - Carnett's sign, 524, 525
 - direct injuries, 522
 - early relaparotomy, 522
 - emergency CS, 523
 - external iliac artery, 521
 - hematoma, 523
 - inguinal hernia repair, 521
 - penetrating abdominal wall trauma, 522, 523
 - peritoneal irritation, 525
 - puerperium, 523
 - right-sided hematomas, 523
 - RSH (*see* Rectus sheath hematoma (RSH))
- Infliximab (IFX), 355, 356
- Injury severity score, 469
- Instrumental abortion
 - incidence, 194, 195
 - prognosis, 195, 196
 - treatment, 195
- Intentional injuries, 492
- Intercalary/peg cells, 223
- Internal hernia, 321, 322
- Internal papillary excrescence, 577
- International classification of diseases (ICD-10), 420
- Intestinal obstruction
 - causes, 270
 - clinical presentation, 271
 - constipation
 - adverse events, 272
 - causes, 271
 - laxatives, 272
 - PEG, 272
 - Rome IV criteria, 271
 - urgency, 271
 - fetal outcome, 276
 - historical perspective, 269–270
 - incidence, 270
 - irritable bowel syndrome, 272
 - maternal outcome, 276
 - Ogilvie's syndrome
 - clinical presentation, 273
 - conservative treatment, 273–274
 - diagnosis, 273, 274
 - incidence, 273
 - pathophysiology, 273
 - risk factors, 273
 - surgical treatment, 274
 - refeeding syndrome
 - initial management, 275–276
 - metabolic/physiological changes, 275
 - monitoring, 276
 - pathophysiology, 275
 - total parenteral nutrition, 275
- Intestinal perforation, 185, 188, 193, 194
 - CRC (*see* Colorectal carcinomas (CRC) in pregnancy)
 - endometriosis (*see* Endometriosis)
 - instrumental abortion (*see* Instrumental abortion)

- spontaneous perforation of colon (*see* Spontaneous colorectal perforation)
- Intestinal stomal obstruction
- causes, 310
 - extraluminal stomal compression
 - diagnosis, 311–313
 - incidence, 311
 - prognosis, 313
 - treatment, 313
 - intussusception and prolapse
 - clinical presentation, 311
 - diagnosis, 311
 - incidence, 310
 - prognosis, 311
 - risk factor, 310
 - treatment, 311
 - ostomy, 310
- Intra-abdominal adhesions, 270
- Intra-abdominal bleeding, 288, 390, 404, 424, 456, 577, 677, 679–681
- Intra-abdominal hypertension (IAH), 770, 775
- etiology pathway, 774
 - hemodynamic shifts, 773
 - maternal care, 773
 - pathophysiology, 772
- Intra-abdominal pressure (IAP), 771
- Intra-abdominal sepsis, 780, 781
- Intracytoplasmic sperm injection, 545
- Intrahepatic cholestasis of pregnancy (ICP), 78, 79
- Intramural fibroids, 675
- Intraoperative cholangiography (IOC), 85
- Intraperitoneal bleeding, paracentesis, 391
- Intraperitoneal rupture of the subcapsular hematoma, 510
- Intrauterine inflammation, 793
- Intrauterine pregnancy, 549
- Intrauterine pressure catheter (IUPC), 630
- Intussusception
- age at presentation, 277
 - bariatric Surgery, 284
 - classification, 277
 - clinical presentation, 279, 280, 311
 - clinicopathological characteristics, 277
 - conservative treatment, 283
 - diagnosis, 311
 - abdominal CT, 282
 - abdominal MRI, 282–283
 - anoscopy/rectoscopy, 283
 - dynamic pelvic MRI, 283
 - plain abdominal x-ray, 281
 - real-time dynamic transperineal ultrasound, 283
 - transabdominal ultrasound, 281–282
 - differential diagnosis, 280
 - etiopathogenesis
 - bariatric surgery, 279
 - benign neurilemmoma, 278
 - colon, 279
 - hamartomatous polyp, 278
 - ileocecal, 279
 - lipoma, 278
 - Meckel's diverticulum, 279
 - outer intussusciptens and the central intussusceptum, 278
 - pregnancy Per Se, 278
 - primary non-Hodgkin's lymphoma, 278–279
 - historical perspective, 277–278
 - incidence, 310
 - male-to-female ratio, 277
 - pregnancy, 278
 - prognosis, 284, 311
 - puerperium, 278
 - risk factor, 310
 - surgical treatment, 283–284
 - treatment, 311
- Ipsilateral Fallopian tube, 544
- Irinotecan, 308
- Irritable bowel syndrome (IBS), 272
- ISK, *see* Ileosigmoid knotting (ISK)
- Isolated Fallopian tube torsion (IFTT)
- abdominal tenderness, 565
 - anesthetic and perioperative management, 570
 - cellular damage, 564
 - classification of causes and theories, 564
 - CT findings, 566
 - description, 563
 - erythrocyte sedimentation rate, 565
 - extrinsic factors, 564
 - fetal and maternal outcome, 570
 - hypoperfusion, 564
 - intrinsic factors, 564
 - laparoscopic access, 568
 - laparotomy, 568
 - mechanical factors, 564
 - MRI, 566
 - operative detorsion, 565
 - pain, 565
 - preoperative diagnosis, 565
 - salpingectomy, 569, 571
 - sequential mechanical event, 564
 - sonographic findings, 566
 - spectral Doppler analysis, 566
 - surgical approach, 568
 - torsion of hydatids of Morgagni, 563
 - transabdominal and transvaginal ultrasound, 566, 567
 - tubal detorsion, 569
 - untwisting (detorsion), 569
 - urinalysis, 565
 - varicose veins, mesosalpinx, 565
 - whirlpool sign, 566
 - white blood cell count, 565
- J**
- Jenkin's rule, 232
- K**
- Kehr's sign, 642
- Kleihauer–Betke test, 447, 452, 453, 473, 477
- Krukenberg tumor, 180, 181

L

Ladd's procedure, 302
 Lansoprazole, 178
 Laparoscopic cystectomy, 582
 Laparoscopy, 584, 761
 advantages, 582
 ectopic pregnancy, 597
 rudimentary horn rupture, 605
 Laparotomy, 584
 abdominal pregnancy, 612
 ectopic pregnancy, 600
 Leiomyomas, 675
 Leukocytosis, 16, 349, 547, 548
 Leukocyturia, 17
 Lipemia retinalis, 119, 121
 Liver packing, 511, 512
 Liver rupture, *see* Spontaneous liver rupture
 Liver transplantation, 513
 Local splenic disorders, 387
 Long-chain 3-hydroxyacyl-CoA dehydrogenase (LCHAD), 109
 Low-density lipoprotein (LDL), 59–61
 Low-molecular-weight heparin (LMWH), 762
 L-thyroxine, 136

M

Magnesium sulfate infusion, 139
 Magnetic resonance cholangiopancreatography (MRCP), 64
 Malignant ovarian tumors, 576
 Mallory-Weiss syndrome, 184
 Marginal ulcer, 185
 Maternal anticonvulsant therapy, 440
 Maternal cardiac arrest, 487–490
 algorithm, 421
 management, 463
 Maternal entry gunshot wound healing, 477
 Maternal injuries and birth defects, 492
 Maternal mean arterial blood pressure (MAP), 418
 Maternal pelvic fractures, 473
 Maternal pelvic trauma, 441, 442
 Maternal respiratory acidosis, 779
 Maternal respiratory support, 463
 Maternal uterine trauma, 426, 427
 Mathews and Mitchell's statement, 270
 Matrix metalloproteinases (MMPs), 791
 Mature cystic teratoma, 575
 McBurney's point, 10, 12, 25
 Mean maximal axillary temperature, 12
 Meckel's diverticulum/diverticulitis (MD), 13, 32, 279
 Medical expulsive therapy (MET), 716
 Mesenteric ischemia
 MVT, 516–520
 PVT, 520, 521
 Mesenteric vein thrombosis (MVT), 520
 abdominal CT scan, 518
 acute, 517
 anticoagulation with heparin, 519

 arterial thrombosis, 520
 autopsy, 516
 clinical classification, 519
 clinical symptoms, 517
 contrast-enhanced abdominal CT, 518
 definitive diagnosis, 518
 explorative laparotomy, 518
 features, 518
 heavy smoking, 517
 high BMI, 517
 inferior mesenteric vein, 516
 intestinal infarction, 516, 518
 leukocytosis, 518
 obstetric management
 mode of delivery, 520
 pregnancy continuation/termination, 520
 occurrence, 516
 oral contraception, 518
 peripheral/regional thrombolysis, 519
 primary (idiopathic), 517
 regional heparinization, 519
 secondary, 517
 segmental gangrenous segments, 518
 subacute/chronic, 518
 surgical thrombectomy, 519
 thrombolytic agents, 518, 519
 Mesh webbing, 434
 Methotrexate (MTX), 356, 615
 Metritis, 14
 Microangiopathic hemolytic anemia, 507
 Microbial invasion of the amniotic cavity (MIAC), 794
 Microsatellite instability, 309
 Military antishock trousers (MAST), 424
 Mind-altering substances, 415
 Mirizzi syndrome, 64, 65, 80
 Modified Alvarado score, 17, 18
 Motor vehicle accidents (MVAs)
 and domestic violence, 416
 maternal injury, 413, 416
 maternal trauma, 414
 non-penetrating abdominal wounds, 413
 penetrating injuries, 414
 Müllerian anomaly, 603, 640
 Mussey, Robert Daniel, 172
 Myomas, 675
 Myometrial contractility, 631, 790
 Myometrial phenotype programming, 790

N

Natalizumab, 356
 National Automotive Sampling System/Crashworthiness Data System (NASS/CDS), 426
 National Trauma Data Bank, 420
 Necrosectomy, 140, 148
 Neonatal hemorrhage, 440
 Neonatal hypocalcemia, 148
 Neural tube defects (NTDs), 303
 Neutrophil granulocytosis with left shift, 349

Neutrophils, 798
 Nitrous oxide, 778
 Noncatastrophic abdominal trauma, 473
 Nonpeptic ulcer perforations/ruptures, 183, 185
 postoperative gastric perforation/rupture (*see*
 Postoperative gastric perforation/rupture)
 spontaneous rupture (*see* Spontaneous gastric
 rupture)
 Non-tubal ectopic pregnancy, 590
 Normal pregnancy
 diagnosis, 332–333
 historical perspective, 330–331
 incidence, 331
 pathophysiology, 331, 332
 treatment, 333
 Northern exposure sign, 315
 Nonsteroidal anti-inflammatory drugs (NSAIDs), 167,
 170, 757, 758, 807
 Nutrition, 781

O

Obstetrical outcomes, 309
 Obturator sign, 11
 Occasional uterine contractions, 467
 Occult rupture, 641
 Ogilvie's syndrome (OS)
 clinical presentation, 273
 conservative treatment, 273–274
 diagnosis, 273, 274
 incidence, 273
 pathophysiology, 273
 risk factors, 273
 surgical treatment, 274
 Omental infarction, 527–529
 Omeprazole, 178
 Omphalitis/periumbilical abscess, 219, 220
 Oophorectomy, 584
 Osmotic laxative, 272
 Osteoporosis, 349
 Ostomies, 237, 310
 Ovarian abscess (OA), 689
 causes, 695
 clinical presentation, 698
 definition, 695
 etiology, 695
 incidence, 689
 non-gynecologic conditions, 689
 Ovarian autoamputation, 544
 Ovarian clear cell carcinoma, 575
 Ovarian cysts
 complications, 575
 diagnosis, 579
 endometriomas, 576
 MRI, 578
 operative treatment, 580
 prevalence, 574
 surgical management, 579
 ultrasound-guided aspiration, 580

Ovarian endometriomas, 580
 Ovarian hyperstimulation syndrome (OHSS), 34, 545,
 546, 773, 776
 Ovarian pregnancy, 590
 Ovarian preservation, 550
 Ovarian teratoma, 581
 clinical presentation, 328–329
 diagnosis, 329
 incidence, 328
 prognosis, 329
 treatment, 329, 581
 Ovarian torsion, 544, 550, 557
 Ovarian transposition, 191, 192
 Ovarian tumors
 incidence, 573
 markers, 577
 sonography, 577
 Ovarian vein thrombosis, *see* Thrombophlebitis
 Ovariectomy, 583
 Ovary dictates, 601

P

Palpable femoral pulses, 420
 Pancreatic cyst fluid analysis, 130
 Pancreatic pseudocysts, 104, 140–143
 Pancreatic tumors, 141
 Paracentesis, 781
 Paracetamol, 758
 Paraovarian cyst during early pregnancy, 544
 Parastomal hernia
 classification, 237
 conservative treatment, 239
 definition, 237
 diagnosis, 238, 239
 emergent operation, 239
 incidence, 238
 medical history, 238
 ostomy types, 237
 physical examination, 238
 Parenchymal ruptures, 726
 Parturition, 165, 790, 792
 Pelvic abscess formation, 693
 Pelvic actinomycosis, 694
 Pelvic and acetabular fracture surgery in pregnancy, 465,
 466
 Pelvic arterial hemorrhage, 525
 Pelvic fractures
 during pregnancy, 468
 in late gestation, 418
 Pelvic inflammatory disease (PID)
 antibiotic therapy, 688
 characteristics, 685
 conservative therapy, 688
 diagnosis, 687
 differential diagnosis, 698
 history, 697
 OA (*see* Ovarian abscess (OA))
 TOA (*see* Tubo-ovarian abscess (TOA))

- Pelvic ligamentous laxity, 414
- Penetrating trauma
- blood tests, 477
 - CT, 478
 - diagnostic peritoneal lavage, 478
 - Doppler flow measurements, 478
 - emergency Cesarean section, 489
 - fetal outcome, 491
 - fetal radiation exposure, 477
 - fetal thoracoabdominal gunshot wounds, 492
 - gunshot wounds
 - anatomical alterations, 482
 - hazards of prematurity, 482
 - management, 480
 - maternal hemodynamic stability, 481
 - nonoperative management, 481
 - peritoneal irritation, 482
 - selective laparotomy, 481
 - selective surgical management, 481
 - surgical exploration, 481
 - tocolytics, 481
 - uterine damage/uncontrollable hemorrhage, 474, 482
 - intrauterine exposure, 477
 - intravenous iodinated contrast, 477
 - maternal mortality rate
 - from gunshot wounds, 491
 - stab injuries, 491
 - maternal visceral injury, 473
 - neurogenic shock, 474
 - obstetric management, 484
 - perimortem Cesarean section, 487, 489, 490
 - postmortem Cesarean section, 488
 - Roentgenogram, 478
 - splenic laceration, 479
 - stab wounds, 485, 486
 - superficial wounds, 484
 - tissue perfusion and fetal oxygenation, 479
 - toxicology screening, 477
 - treatment algorithm, 485, 486
 - ultrasonography, 478
 - uterine stab wounds, 474–476
- Penicillins, 757
- Peptic ulcer bleeding
- acid suppression medications, 515
 - conservative management, 515
 - dyspeptic symptoms, 513
 - esophagogastroduodenoscopy, 515
 - fetal mortality, 514, 516
 - gastroduodenostomy, 515
 - hematemesis, 514
 - hemorrhage, 515
 - maternal mortality, 515
 - progesterone treatment, abortion, 515
 - radiological examination, 515
 - surgical procedure, 515
 - surgical treatment for hemorrhage, 515
- Peptic ulcer disease (PUD)
- clinical presentation, 168
 - decade dependency, 163–164
 - diagnosis, 168, 169
 - fertility rate, 166, 167
 - gastric outlet obstruction, 172
 - gastric position changes, 165
 - H. pylori* infection, 170, 171
 - historical perspective, 163
 - hormones, 165, 166
 - hypochlorhydria, 165
 - incidence in pregnancy, 164
 - parturition, 165
 - perforated ulcer in pregnancy
 - antacids, 178
 - clinical presentation, 174
 - diagnosis, 176, 177
 - gastric acid suppression during lactation, 178, 179
 - general population, 172, 173
 - H₂RAs, 177–179
 - historical perspective, 172–174
 - incidence, 174
 - PPIs, 178, 179
 - prognosis, 179
 - puerperium, 174–176
 - risk factors, 174
 - surgical treatment, 177
 - ulcer types, 174, 175
 - prognosis, 169
 - risk factors
 - causes, 168
 - fasting, 168
 - Helicobacter pylori* infection, 167
 - hypersecretory states, 167, 168
 - lifestyle factors, 167
 - medications, 167
 - physiological stress, 167
 - treatment, 169
 - women vs. men, 163
 - ZES, 170
- Percutaneous biliary drainage, 68
- Percutaneous nephrostomy, 716
- Perforated gastric cancer
- incidence, 183
 - primary gastric tumor, 183
 - prognosis, 183
- Perianal disease activity index, 360
- Periappendicitis decidualis, 35
- Perihepatic packing, 511
- Perihepatitis, *see* Fitz-Hugh-Curtis syndrome
- Peripherally inserted central catheters (PICC), 68
- Perityphlitic abscess, 37
- Periumbilical cellulitis, 594
- Periumbilical ecchymosis, 594
- Persistent ectopic pregnancy, 591
- Pfannenstiel incision, 215, 216, 228, 229
- Placental abruption, 460–462, 465
- incidence of, 799
 - infection-induced, 799
 - pathophysiologic mechanisms, 799
 - principal mechanisms, 798
 - trauma-induced, 799
- Placental injury
- abdominal injury/deceleration, 446
 - countercoup mechanism, 446, 447

- direct injury, 446, 447
 - traumatic laceration, 446
 - Placental insufficiency, 444, 478
 - Placental pathology, 801
 - Placenta percreta, 650
 - Plasmapheresis, 137, 138
 - Pneumoperitoneum, 27, 764, 778
 - Pneumothorax, 480
 - Porro procedure, 509
 - Portal vein thrombosis (PVT), 520, 521
 - Post-bariatric surgery
 - abdominal CT, 323
 - abdominal MRI, 323, 324
 - classification, 321
 - clinical presentation, 322–323
 - fetal outcome, 325
 - incidence, 321
 - maternal Outcome, 325
 - morbid obesity, 321
 - treatment, 323–325
 - Post-gastric banding, 185
 - Postoperative (incisional) hernia
 - abdominal wall endometrioma
 - clinical presentation, 229, 230
 - diagnosis, 230, 231
 - incidence, 229
 - management, 235
 - prognosis, 237
 - component separation technique, 232
 - general population, 227
 - gravid uterus
 - clinical presentation, 229, 230
 - complications, 234
 - diagnosis, 230
 - incidence, 228
 - management, 234, 235
 - laparoscopic repair, 233
 - mesh repair before pregnancy, 236
 - open repair technique, 232, 235, 236
 - postoperative technique, 233
 - pregnant population, 228
 - preoperative technique, 233
 - Postoperative gastric perforation/rupture
 - diaphragmatic hernia, 185
 - post-gastric banding, 185
 - Postpartum bleeding, 641
 - Postpartum splenic rupture, 389, 390, 393
 - Preeclampsia, 774
 - Preeclampsia-eclampsia syndrome, 510
 - Pregnancy Risk Assessment Monitoring System (PRAMS), 435
 - Pregnancy-associated adnexal masses, 544
 - Prematurity, 811
 - Prenatal care, 416
 - Prenatal care provider counseling for seat belt use, 435, 436
 - Preterm labor (PTL)
 - definition, 789
 - etiopathogenesis, 792–800
 - fetal morbidity, 799
 - inflammation-induced, 802, 803
 - intrauterine infection, 801
 - maternal stress, 795
 - physiologic activation, 790
 - prevention and treatment, 570
 - proinflammatory cytokine, 795
 - tocolytic therapy, 804
 - transvaginal sonography, 801
 - Preterm premature rupture of the membrane (PPROM), 789–791, 794, 795, 798–801, 803, 804, 806, 809, 810
 - Primary hyperparathyroidism, 104
 - Probiotics, 357
 - Progesterone, 806
 - Progesterone receptors (PgR), 303
 - Prolapse
 - clinical presentation, 311
 - diagnosis, 311
 - incidence, 310
 - prognosis, 311
 - risk factor, 310
 - treatment, 311
 - Prophylactic plasmapheresis, 137
 - Prostaglandin, 791, 800
 - Psoas sign (Obraztsova's sign), 11
 - Puerperium, 694
 - Pyosalpinx, 578
- Q**
- Quiet rupture, 641
- R**
- Ranson scoring system, 127
 - Rebound tenderness, 11, 301, 367, 547
 - Rectal hemorrhage/melena, 530
 - Rectal/pelvic tenderness, 11
 - Rectal wall, 187
 - Rectus sheath hematoma (RSH), 522, 524
 - abdominal ultrasonography, 525
 - aortoiliac arteriogram, 527
 - CT, 525
 - Doppler sonography, 525
 - epigastric embolization, 526
 - Grey-Turner's sign, 524
 - hemodynamics, 526
 - MRI, 525, 526
 - opioids, 526
 - peroral NSAIDs, 526
 - skin changes, 525
 - transarterial embolization, 526
 - types, 525
 - Refeeding syndrome, 275
 - initial management, 275–276
 - monitoring, 276
 - pathophysiology, 275
 - Repeated blunt abdominal trauma, 473
 - Retroplacental hemorrhage, 799
 - Rho(D) immune globulin, 479
 - Right transrectal/pararectal/paramedian incision, 26
 - Rosette test, 452

- Round ligament pain/syndrome (RLP), 12, 13
- Round ligament stretch/pain, 209, 210
- Round ligament varicosities (RLVs)
- clinical presentation, 207, 208
 - diagnosis, 208, 209
 - incidence, 207
 - pathophysiology, 207
 - treatment, 208
- Rovsing's sign, 11
- Rudimentary horn rupture, 605
- anesthetic and perioperative management, 608
 - CT, 605
 - definition, 602
 - diagnostic criteria, 604
 - endometriosis incidence, 603
 - history, 603
 - hysterosalpingography, 605
 - microsurgical fallopian tube transposition, 608
 - mortality rate, 608
 - MRI, 605
 - noncommunication, 602
 - preclinical and preoperative detection, 604
 - treatment, 605
 - ultrasound, 604
 - uterus bicornis, 603
 - with ipsilateral salpingectomy, 606
- Ruptured splenic pregnancy
- amenorrhea, 395
 - anesthetic and perioperative management, 396
 - classification, 394
 - CT scan, 395
 - hemorrhagic lesion, 395
 - intraperitoneal bleeding, 395
 - laparoscopic treatment, 396
 - placental attachment, 394
 - preterm labor, 396
 - primary, 394
 - transvaginal ultrasonography, 395
- S**
- Safe motherhood initiatives, 442
- Salpingectomy, 600
- Salvageable infant survival, 470
- Salvageable infants, 490, 491
- Scar endometrioma, 229
- Seat belt syndrome, 434, 435
- Sedation, 752
- Segmental liver resection, 512
- Selective (wait-and-see) approach, 84
- Sepsis/septicemia, 719
- Septate uterus, 640, 664
- Sexually transmitted organisms, 686
- Shoelace technique, 232
- Sigmoid colon, 186, 188
- Sigmoid mesocolon hernia
- abdominal MRI, 326
 - classification, 325
 - clinical features, 325
 - CT, 325
 - etiology, 325
 - incidence, 325
 - plain abdominal X-ray, 325
 - prognosis, 326
 - treatment, 326
- Sigmoid volvulus
- abdominal MRI, 315, 316
 - clinical presentation, 314–315
 - incidence, 313, 314
 - laboratory findings, 315
 - pathophysiology, 313–314
 - plain abdominal X-ray, 315
 - prognosis, 318
 - risk factors, 313–314
 - surgical treatment, 317–318
- Silent antepartum UR, 641
- Silent rupture, 641
- Single incision laparoscopic surgery (SILS), 75
- Small bowel volvulus
- abdominal CT, 297
 - clinical presentation, 296
 - congenital intestinal malrotation
 - clinical presentation, 298
 - diagnosis, 298
 - etiopathogenesis, 298
 - incidence, 298
 - prognosis, 299
 - treatment, 298–299 - incidence, 296
- ISK
- abdominal CT, 301, 302
 - abdominal MRI, 301
 - clinical presentation, 300–301
 - definition, 299
 - diagnosis, 301
 - etiopathogenesis, 300
 - incidence, 300
 - plain abdominal X-rays, 301
 - prognosis, 302
 - risk factors, 300
 - transabdominal ultrasound, 301
 - treatment, 301–302
 - types, 299
 - pathophysiology, 296
 - prognosis, 298
 - transabdominal ultrasound, 297
 - treatment, 297–298
- Smead-Jones mass closure, 232
- Spiegelberg criteria, 590
- Spinal cord injury, 479, 480, 485, 756
- Splenic artery aneurysm (SAA)
- clinical presentation
 - ruptured splenic artery (aneurysm), 399
 - symptomatic splenic artery aneurysm, 399 - cross-sectional imaging techniques, 397
 - definition, 396
 - diagnosis
 - abdominal sonography, 400
 - contrast-enhanced abdominal CT, 400
 - 3D reconstruction of abdominal MSCT, 400

- differential diagnosis, 399
- pathophysiology, 398, 399
- postmortem examination, 396
- prognosis, 401, 402
- ruptured saccular aneurysm, 397
- spontaneous rupture, normal splenic artery (vein), 398
- spontaneous splenic artery aneurysm rupture, 398
- treatment
 - anesthetic and perioperative management, 401
 - preterm labor prevention, 401
 - ruptured SAA, 401
 - symptomatic SAA, 401
- types, 397, 398
- visceral artery aneurysms, 396
- Splenic rupture
 - anesthetic and perioperative management, 393
 - arteriography, 392
 - autopsies, 385
 - Ballance's sign, 390
 - Cesarean section, 389
 - classification, 386
 - direct trauma, 386
 - spontaneous rupture, 387
 - toxemia of pregnancy, 387
 - CT scan, 392
 - ectopic pregnancy, 390
 - epigastric pain, 390
 - hemodynamic monitoring, 392
 - intraabdominal bleeding, 389, 391, 393
 - intrasplenic aneurysm, 388, 389
 - Kehr's sign, 390
 - laparoscopic splenectomy, 393
 - laparotomy, 392
 - maternal death, 393
 - mechanisms of rupture, 388
 - normal or physiological leukocytosis, 385
 - pregnancy-related causes, 386
 - preterm labor, 393
 - puerperium, 390
 - radiographic examination, 391
 - rapid plasma expansion, 389
 - splenectomy, 392
 - splenic artery angiography, 393
 - spontaneous rupture, 388
 - transabdominal ultrasound, 391
 - uterine tetany, 391
 - uteroplacental perfusion, 394
- Splenic torsion
 - anesthetic and perioperative management, 406
 - clinical presentation, 405, 406
 - diagnosis, 406
 - differential diagnosis, 406
 - etiopathogenesis, 405
 - historical perspective, 405
 - incidence, 405
 - preterm labor prevention, 406
 - prevention, 406
 - prognosis, 407
 - surgical treatment, 406
- Splenic vein aneurysm (SVA), 402–404
- Spontaneous abdominal wall endometrioma, 229
- Spontaneous colorectal perforation, 193, 194
 - idiopathic perforation, 193
 - intestinal perforation (*see* Spontaneous intestinal perforation)
 - stercoral perforation (*see* Stercoral perforation)
- Spontaneous gastric rupture
 - etiopathogenesis, 184, 185
 - historical perspective, 183
 - incidence, 183
 - prognosis, 185
- Spontaneous hepatic hematomas associated with pregnancy (SHHP), 507, 508, 510, 511
- Spontaneous intestinal perforation
 - incidence, 194
 - prevention, 194
 - risk factor, 194
 - treatment, 194
- Spontaneous liver rupture
 - anticoagulation, 510
 - antihypertensive agents, 510
 - argon beam electrocoagulation, 512
 - blood volume replacement, 510
 - cardiovascular instability, 508
 - causes, 506, 508
 - cicatrizization, 507
 - CT scan, 509
 - eclampsia, 506
 - etiopathogenesis, 507
 - fetal heart sounds, 508
 - fetal mortality, 513
 - HELLP syndrome, 506
 - hemodynamic monitoring, 510
 - hemodynamic status, patient, 508
 - hemolysis, 507
 - hepatic ischemic stage, 507
 - hypertension, 510
 - hypovolemic shock, 507
 - intraoperative hepatic artery occlusion, 511
 - large hepatic hematoma, 507
 - liver packing, 511
 - liver transplantation, 513
 - magnesium sulfate administration, 510
 - maternal mortality, 513
 - median laparotomy, 513
 - muscular activity, 506
 - muscular effort, 506
 - nonoperative management, 510
 - obstetric management, 513
 - pathology, 508
 - percutaneous angiographic embolization, 510, 511
 - perforation of subcapsular hepatic hematoma, 507
 - peritonitis, 508
 - Pfannenstiel incision, 513
 - preeclampsia, 506
 - pregnancy complications, 505
 - pulmonary embolism, 508
 - segmental liver resections, 512
 - sudden death during eclampsia, 506
 - surgical modalities, 511

- Spontaneous liver rupture (*cont.*)
 toxemia, 506
 transabdominal ultrasound, 509
 transient hypertension, 506
 traumatic, 506
 treatment algorithm, 512
- Spontaneous rectus sheath hematoma, 523, 524, 526–527
- Spontaneous rupture, 725
- Spontaneous splenic vein (aneurysm) rupture
 anesthetic and perioperative management, 405
 cirrhosis, 403
 clinical presentation, 404
 diagnosis, 404
 differential diagnosis, 404
 etiology, 402
 fetal outcome, 405
 hemodynamic and hormonal changes in pregnancy, 403, 404
 historical perspective, 402
 incidence, 402
 maternal outcome, 405
 mechanical factors, 403
 preterm labor treatment, 405
 risk factors, 403
 surgical treatment, 405
- SpyGlass Direct Visualization System, 83
- Stercoral perforation
 diagnostic criteria, 193
 incidence, 193
 prognosis, 193
 risk factor, 193
 treatment, 193
- Stimulant laxatives, 272
- Stump appendicitis, 13
- Subcapsular liver hematoma, 509, 510
- Submucosal fibroids, 675
- Subserosal fibroids, 675–676
- Subtotal hysterectomy, 650
- Sulfasalazine, 807
- Supine hypotensive syndrome, 423, 770
- Symptomatic choledochal cysts
 acute cholangitis, 89
 asymptomatic patients, 89
 choledochal cyst rupture, 89, 90
 clinical presentation, 87
 diagnosis, 87, 88
 differential diagnosis, 87
 fetal outcome, 90
 historical perspective, 86, 87
 maternal outcome, 90
 obstetric management, 88
 pathophysiology, 87
- Symptomatic cholelithiasis, 764
- T**
- Temporary abdominal closure (TAC) method, 781, 782
- Teratogenicity, 751
- Tetanus prophylaxis, 479
- Thalidomide, 356–357
- Thiopurines, 353–354
- Thrombophlebitis, 14, 15
- Thromboprophylaxis guidelines, 762
- Thrombosis, 759
- TLR4 antagonists, 807
- Tocolytics, 804–809
- Torsion of hydatids of Morgagni, 544
- Total abdominal hysterectomy, 649
- Total hysterectomy, 651
- Total parenteral nutrition, 275
- Toxemia, 774
- Toxic megacolon
 pregnancy outcomes, 376
 ulcerative colitis, 366
- Transmesosigmoid hernia, 325, 326
- Transomental hernia, 326
- Transvaginal instrumental uterine perforation
 clinical presentation, 287–288
 conservative treatment, 290
 diagnosis
 abdominal CT, 289, 290
 abdominal MRI, 289
 abdominal ultrasound, 288–290
 laboratory findings, 288
 plain abdominal x-ray, 288
 gynecologic/obstetric procedures, 291
 incidence, 284–285
 prognosis, 291–292
 risk factors, 285
 small bowel obstruction, mechanisms, 285–287
 surgical treatment, 290–291
- Transvaginal sonography (TVS), 698
- Transvaginal technique, 596
- Trauma victim, 755
- Traumatic placental abruption, 444–446
- Traumatic rectus sheath hematoma, 442, 524–525
- Traumatic renal rupture, 726
- Tubal pregnancy, 590
- Tuberculous salpingitis, 694
- Tubo-ovarian abscess (TOA), 686, 703
 antibiotic therapy, 688
 clinical features, 691
 diagnosis, 687
 incidence, 685, 694
 intraperitoneal rupture, 687
 mechanisms, 692
 ruptured and unruptured, 697
 surgical therapy, 701
 treatment, 687
 ultrasonography, 687
- Twisted ovarian mass, 555
- Twisted ovaries in pregnancy, 548
- U**
- Ulcerative colitis (UC)
 classification, 366
 clinical presentation
 elective, 366–367
 emergent, 367

- diagnosis
 - abdominal CT, 368, 369
 - colonoscopy, 368
 - flexible and rigid sigmoidoscopy, 368
 - plain abdominal X-ray, 368
 - proctoscopic examination, 368
 - supine X-ray, 368
 - toxic dilatation, 369
 - ultrasound, 368
- differential diagnosis, 367
- incidence, 364
- medical treatment, 370
- pathophysiology, 366
- pregnancy outcomes
 - completed pregnancies, 374
 - elective surgery, 375
 - emergency surgery, 375, 376
 - low birth weight and postmaturity, 375
 - maternal mortality rate, 374
 - predictors, 375
 - toxic megacolon, 376
 - uncomplicated vaginal delivery, 375
- pregnancy, effect of, 364–366
- surgical treatment
 - anesthetic and perioperative management, 372
 - Cesarean section, 372
 - fertility and sexual health, 373–374
 - indications, 370
 - mode of delivery, 372–373
 - preterm labor, 372
 - subtotal colectomy, 372
 - total abdominal colectomy, 372
 - Turnbull “blowhole” procedure, 370–372
- Ultrasound-guided aspiration, 580
- Umbilical endometriosis (UEM)
 - clinical presentation, 220
 - diagnosis, 220, 221
 - incidence, 220
 - prognosis, 221, 222
 - treatment, 220–222
- Umbilical hernia
 - clinical presentation, 219
 - endosalpingiosis (*see* Endosalpingiosis)
 - gravid uterus, 219
 - incidence, 217, 218
 - laparoscopic/single port surgery, 219
 - omphalitis/periumbilical abscess, 219, 220
 - risk factors, 218
 - UEM (*see* Umbilical endometriosis (UEM))
 - uterine fibroids, 218
- Ureterorenoscopy, 716
- Ureteroscopy, 717
- Urinalysis, 727
- Urinary bladder injury, 732
 - incidence, 732
 - intraperitoneal, 733
- Urinary bladder rupture, 442
- Urinary stone disease, 712
- Urinary tract infections (UTIs)
 - anatomic and physiological changes, 717
 - types, 717
- Urinary tract stones
 - CT scan, 715
 - diagnosis, 714
 - diet therapy, 715, 716
 - differential diagnosis, 713
 - etiopathogenesis, 712
 - intravenous urography, 715
 - medical expulsive therapy, 716
 - medical management, 715
 - MR urography, 715
 - signs/symptoms, 713
 - surgical intervention, 716
 - ultrasound, 715
- Urine β HCG assay, 595
- Urolithiasis/urinary tract infection, 13, 712
- Uroperitoneum, 734
- Urostomy, 237
- Ursodeoxycholic acid, 68
- U-shaped distended sigmoid loops, 315
- Uterine fibroids, 218, 225
 - classification, 675
 - definition, 675
- Uterine hypertonia, 667
- Uterine isthmus, levels, 666
- Uterine leiomyomas
 - incidence, 676
 - myomectomy, 681
 - spontaneous hemoperitoneum, 677
- Uterine myoma
 - diagnosis, 681
 - MRI, 679
 - optimal treatment, 680
 - ultrasound, 678
- Uterine rupture (UR)
 - cause, 627
 - classification, 623
 - CT scan, 643
 - definition, 622
 - diagnosis, 642
 - fetal mortality, 654
 - history, 621
 - incidence, 626
 - management, 627
 - maternal morbidity, 652
 - mechanisms
 - atypical site rupture, 625
 - obstruction, 625
 - scar rupture, 624
 - MRI, 644
 - neonatal consequences, 653
 - neonatal morbidity, 653
 - physical examination, 642
 - prophylactic antibiotics, 647
 - risk factors
 - classic cesarean delivery, 628
 - clinical heterogeneity, 629
 - labor induction, 630
 - low-transverse cesarean, 629
 - low-vertical cesarean, 628

- Uterine rupture (UR) (*cont.*)
 - scarred, 627
 - subsequent trial of labor, 629
 - unknown scar, 628
 - surgical treatment, 648
 - complications, 652
 - hysterectomy, 649
 - management, 651
 - suture repair, 648, 649
 - total hysterectomy, 651
 - symptoms and signs, 640–642
 - transvaginal and transabdominal ultrasound, 643
 - types, 624
 - uterine tone monitoring, 645
- Uterine scar dehiscence, 623
- Uterine torsion (UT)
- causative factors, 666
 - clinical findings, 667
 - clinical presentation, 666, 667
 - CT scan, 668
 - detorsion, 669
 - differential diagnoses, 668
 - during puerperium, 666
 - history, 663
 - hysterotomy, 670
 - intrinsic intrapelvic pathology, 665
 - maternal mortality rate, 671
 - MRI, 668
 - nonsurgical methods, 669
 - obstetric decisions, 671
 - operation goals, 668
 - physiological dextrorotation, 664
 - preoperative diagnoses, 668
 - ultrasound, 668
- Uterocaval compression, 423
- Uteroplacental circulation, 418
- Uterosacral ligaments , bilateral plication, 670
- Uterus injury scale, 451
- V**
- Vacuum-assisted closure (VAC), 782
 - Vaginal birth after CS (VBAC), 627
 - Vaginal delivery, 712
 - Vasopressors, 755
 - Vasospasm of hepatic arterial circulation, 507
 - Venous thromboembolism (VTE), 762
 - Veterinary obstetrics, 663
 - Villar nodule, 220
 - Vitamin D deficiency, 349
- Z**
- Zollinger-Ellison syndrome (ZES), 169, 170