
Index

A

Abbreviated mental test score (AMTS), 111
Activities of daily living (ADLs), 47, 129
Acute confusional state, 110
Acute delirium, 61
Advanced activities of daily living (AADLs), 47
Advanced healthcare directive (AHD), 141
Airway, 55
Alcohol, 6
Anti-resorption, 10

B

Baseline health, assessment of, 110
Basic activities of daily living (BADLs), 47
Bisphosphonates, 10
Blanching, 87
Blaylock Risk Assessment Screening Score (BRASS), 131
Body mass index (BMI), 6
Bone
 biology, 2
 bone mass, 1
Bone mineral density (BMD), 6–8
Breathing, 55

C

Calcium, 97
Calorie supplementation, 18, 19
Caregiver burden, 140, 141
Care integration, 133
CGA, *see* Comprehensive geriatric assessment
Circulation, preoperative care, 55
Cognition, 47

Cognitive syndromes, 109
Communication, 138
Competence, 152
Comprehensive geriatric assessment (CGA), 110
 assessment team, 49–51
 assessment tools, 48, 49
 cognitive impairment, 41
 concept, 42–45
 dimensions, 46–48
 malnutrition, 41
 nursing assessment, 42
 posthospital discharge, 49
 pressure ulcers, 41
 secondary prevention, 49
Confusion Assessment Method (CAM), 112, 113
Constipation
 fundamental nursing care, 61
 prevention of, 77
 stasis, complications of, 71, 72
Continuity of care, 132, 133
Coordination of care, 133

D

Deep vein thrombosis (DVT), 71
Deficit Accumulation Index, 18
Dehydration, 95, 98
 evidence-based interventions, 100, 101
 hydration and, 101–104
 prevalence, 98
 risk factors, 102
 screening and assessing patients, 99, 100
 symptoms, 98

- Delirium
 acute, 61
 acute confusional state, 110
 assessment, 110–112
 consequences, 110
 identifying delirium, 112
 management, 113, 114
 medication, 114, 115
 predisposing and precipitating factors, 111
 prevention, 112, 113
- Dementia
 advocacy, 117, 118
 capacity, 116, 117
 early stage, 115
 late stage, 115
 middle stage, 115
 two-stage assessment, for mental capacity, 117
- Depression, 28, 118, 119
 assessment, 119, 120
 interventions, 120
- Deprivation of Liberty Safeguard (DoLS), 118
- Disability, preoperative care, 55
- Discharge
 and post-hospital care, 129
 admission phase, 131
 continuity of care, 132
 discharge home, 130
 discharge phase, 131, 132
 hospitalisation phase, 131
 nurse's role in, 133
 setting, 130, 131
- Dual-energy X-ray absorptiometry (DXA), 7, 8, 10, 36
- DVT, *see* Deep vein thrombosis
- E**
- Eating disorders, 6
- Effective communication, 150
- End of life care, 141–143
- Ethnicity, 5
- Exercise, 6, 69, 70
 assessment, 74
 evidence-based interventions, 74, 75
 frailty, 18
 sarcopenia, 21
- Exposure, 55
- F**
- Falls, 29
 assessment, 114
 causes, 29
 CGA, 47
 extrinsic factors, 29
 frailty, sarcopenia and, 23
 holistic, person-centred assessment, 28
 intrinsic factors, 29
 prevention strategies, 30
 environment, 30
 exercise, 31
 fear of falling, 31, 32
 footwear and foot care, 31
 medication review, 31
 pathways and guidelines, 32
 visual impairment, 31
 screening and assessment, 29, 30
- Family partnerships, 138, 139
 assessment, 139, 140
 caregiver burden, 140, 141
 legal and ethical considerations, 141
- Fear of falling, 31, 32
- Femoral neck fracture, 56
- Financial support, CGA, 47, 48
 4AT, 111
- Fracture Liaison Coordinator, 33, 34, 37
- Fracture liaison services (FLS), 7, 28, 33, 34
- Fracture Risk Assessment Tool (FRAX®), 1, 8, 36
- Fragility fracture, 125, 144
 and osteoporosis, 3, 4
 case finding, 7
 diagnosis, 7
 risk evaluation, 7, 8
 risk factors, 5–7
 secondary osteoporosis, differential diagnosis of, 9
 treatment, 9–11
 secondary' prevention, 1
- Frailty, 16
 assessment, 17, 18
 frailty index, 18
 interventions, 18, 19
- Functional Independence Measure (FIM®), 131
- Functional status, 46, 47
- G**
- Genant classification, 8
- Geriatric syndrome, 41
- H**
- Healthcare-associated infection, 61
- Health education, 37
- Hip fracture, 4
 diagnosis and surgery, 56–58
 discharge and post-hospital care, 129
 DVT, 71
 intertrochanteric, 56
 pain management, 58, 59

- palliative and end of life care, 141, 142
 - perioperative care, 54
 - preoperative care, 54
 - rehabilitation, 126, 128
 - Hospital-acquired infections (HAIs), 71, 92
 - Hospital Anxiety and Depression Score (HADS), 119
 - Hospital Elder Life Program (HELP), 114
 - Hydration, 95, 101
 - assessment, 102, 103
 - evidence-based interventions, 103, 104
 - fundamental nursing care, 61
 - screening and assessing patients with dehydration, 102
 - Hypogonadism, primary/secondary, 5

 - I**
 - Incontinence-associated dermatitis (IAD), 86
 - Independent mental capacity advocate (IMCA), 117
 - Informal caregiving, 138
 - Instrumental/intermediate activities of daily living (IADLs), 47
 - Integrated care pathways (ICPs), 127
 - International Osteoporosis Foundation (IOF), 33
 - Intertrochanteric hip fracture, 56

 - L**
 - Long-term glucocorticoid therapy, 5
 - Low energy trauma, 3

 - M**
 - Malnutrition, 41, 95, 98
 - evidence-based interventions, 100, 101
 - hydration and, 101–104
 - prevalence, 98
 - risk factors, 102
 - screening and assessing patients, 99, 100
 - symptoms, 98
 - Mechanostat theory, 21
 - Menopause, 5
 - Mental Capacity Act, 116, 118
 - Mini Mental State Examination (MMSE), 112
 - Mobility, 68, 69
 - assessment of, 72, 73
 - evidence-based interventions, 73, 74
 - Modified falls efficacy scale, 30
 - Monoclonal antibody, 10
 - Montreal Cognitive Assessment (MoCA), 112
 - Mood, 47
 - Multidisciplinary team (MDT), 31, 73, 129, 144, 147, 151

 - Multimodal analgesia, 59

 - N**
 - Nutrition
 - calcium and vitamin D, 97
 - fundamental nursing care, 61
 - healthy diet, for older adults, 96, 97
 - malnutrition and dehydration, 98
 - evidence-based interventions, 100, 101
 - hydration and, 101–104
 - prevalence, 98
 - risk factors, 102
 - screening and assessing patients, 99, 100
 - symptoms, 98
 - pressure injury prevention, 91
 - sarcopenia, 21

 - O**
 - Oestrogen deficiency, 6
 - Opioid analgesia, 59
 - Orthogeriatric nursing, 43, 147, 148
 - care quality, 150
 - caring art and a science, 149
 - contribution, 151
 - fundamental nursing care, 60, 61
 - health improvement and health promotion, 151
 - hip fracture and
 - diagnosis and surgery, 56–58
 - pain management, 58, 59
 - International Council of Nurses (ICN)
 - definitions of nursing, 148
 - orthogeriatric care, education for, 152
 - orthogeriatric patient, 149
 - perioperative care, 54
 - postoperative care, 60
 - preoperative care, 54, 55
 - preparation for discharge, 62
 - resource nursing, 152
-
- Osteoblasts, 2
- Osteoclasts, 2
- Osteocytes, 2
- Osteopenia, 27
- Osteoporosis
 - epidemiology, 3
 - fragility fracture and, 3, 4
 - case finding, 7
 - diagnosis, 7
 - risk evaluation, 7, 8
 - risk factors, 5–7
 - secondary osteoporosis, differential diagnosis of, 9
 - treatment, 9–11
 - normal and osteoporotic bone, 2, 3

P

Pain, 4, 48, 58–60, 68, 70, 75
Palliative care, 137, 141–143
Parathyroid hormone, 10
Patient-centred medicine, 138
Patient Health Questionnaire 9 (PHQ-9), 120
Peak bone mass, 2
Perioperative care, 54
Physical Frailty Phenotype (PFP), 17
Polypharmacy, 19, 47, 48
Preoperative care, 54, 55
Pressure injuries, 86
 classification of, 87, 88
 pathophysiology and causes, 86, 87
 prevention, 88
 evidence-based preventive
 interventions, 89–91
 patient assessment, 89
Pressure injury prevention, 61
Previous fracture, 5
Primary/secondary hypogonadism, 5
Protein supplementation, 18, 19
Proximal femur fracture, 4
Public health, 3
Pulmonary infections, prevention of, 77
Pulmonary tract infection, 71

Q

Quality care, 150
Quality of life, 4

R

Rehabilitation
 mobility and exercise, 127, 128
 pathways, 126, 127
 patient motivation, 128, 129
 primary outcomes, 126
Remobilisation
 evidence-based interventions, 73, 74
 mobility and, 68, 69
 potential, 72, 73
Remodelling, 2
Rheumatoid arthritis, 5

S

Sarcopenia, 19
 clinical consequences, 21
 interventions, 21–23
 screening and assessment, 20

Secondary fracture prevention, 32, 33, 61
 fracture prevention services and guidelines,
 33–35
 models of care, 28
 morbidity, mortality and functional ability
 outcomes, 28
 typical patient journey, 35
 assessment, 36
 evaluation, 38
 FLS, first contact with people, 36
 health education, 37
 personal plan, establishing, 37
Selective oestrogen receptor modulators
 (SERM), 10
Smoking, 6
Social support, CGA, 47, 48
Stasis
 complications of, 70
 assessment, 75, 76
 constipation, 71, 72
 DVT, 71
 prevention, 76–79
 pulmonary and urinary tract
 infection, 71

T

Thirty-second chair stand, 30
Thromboembolism, stasis complications,
71, 76, 77
Timed Up and Go test (TUG), 30
Tinetti balance assessment tool, 30

U

Urinary catheters, 71
Urinary tract infections (UTI), stasis
 complications, 71, 77

V

Vertebral fracture, 4
Vertebral Fracture Assessment (VFA), 8
Vitamin D supplementation, 6, 19, 97

W

Weight loss, 19
Wound healing, 91
Wound management, 91, 92
 preventing wound infection, 92, 93
 studies, 93