

References

1. Alazard, T., Baldi, P.: Gravity capillary standing water waves. *Arch. Ration. Mech. Anal.* **217**(3), 741–830 (2015). <https://doi.org/10.1007/s00205-015-0842-5>
2. Alazard, T., Delort, J.-M.: Global solutions and asymptotic behavior for two dimensional gravity water waves. *Ann. Sci. Éc. Norm. Supér.* (4) **48**(5), 1149–1238 (2015)
3. Alazard, T., Delort, J.-M.: Sobolev estimates for two dimensional gravity water waves. *Astérisque* **374**, viii+241 (2015)
4. Alazard, T., Métivier, G.: Paralinearization of the Dirichlet to Neumann operator, and regularity of three-dimensional water waves. *Commun. Partial Differ. Equ.* **34**(10–12), 1632–1704 (2009). <https://doi.org/10.1080/03605300903296736>
5. Alazard, T., Burq, N., Zuily, C.: On the water-wave equations with surface tension. *Duke Math. J.* **158**(3), 413–499 (2011). <https://doi.org/10.1215/00127094-1345653>
6. Alazard, T., Burq, N., Zuily, C.: On the Cauchy problem for gravity water waves. *Invent. Math.* **198**(1), 71–163 (2014). <https://doi.org/10.1007/s00222-014-0498-z>
7. Alazard, T., Burq, N., Zuily, C.: Cauchy theory for the gravity water waves system with non-localized initial data. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **33**(2), 337–395 (2016). <https://doi.org/10.1016/j.anihpc.2014.10.004>
8. Alinhac, S.: Paracomposition et opérateurs paradifférentiels. *Commun. Partial Differ. Equ.* **11**(1), 87–121 (1986). <https://doi.org/10.1080/03605308608820419>
9. Alinhac, S.: Existence d’ondes de raréfaction pour des systèmes quasi-linéaires hyperboliques multidimensionnels. *Commun. Partial Differ. Equ.* **14**(2), 173–230 (1989). <https://doi.org/10.1080/03605308908820595>
10. Ambrose, D.M.: Well-posedness of vortex sheets with surface tension. *SIAM J. Math. Anal.* **35**(1), 211–244 (2003). <https://doi.org/10.1137/S0036141002403869>
11. Ambrose, D.M., Masmoudi, N.: The zero surface tension limit of two-dimensional water waves. *Commun. Pure Appl. Math.* **58**(10), 1287–1315 (2005). <https://doi.org/10.1002/cpa.20085>
12. Arnol’d, V.I.: Reversible systems. In: *Nonlinear and Turbulent Processes in Physics*, vol. 3 (Kiev, 1983), pp. 1161–1174. Harwood Academic, Chur (1984)
13. Baldi, P., Berti, M., Haus, E., Montalto, R.: Time quasi-periodic gravity water waves in finite depth. *Inv. Math.* (2018). <https://doi.org/10.1007/s00222-018-0812-2>
14. Bambusi, D., Grébert, B.: Birkhoff normal form for partial differential equations with tame modulus. *Duke Math. J.* **135**(3), 507–567 (2006). <https://doi.org/10.1215/S0012-7094-06-13534-2>

15. Bambusi, D., Delort, J.-M., Grébert, B., Szeftel, J.: Almost global existence for Hamiltonian semilinear Klein-Gordon equations with small Cauchy data on Zoll manifolds. *Commun. Pure Appl. Math.* **60**(11), 1665–1690 (2007). <https://doi.org/10.1002/cpa.20181>
16. Berti, M., Montalto, R.: Quasi-periodic standing wave solutions of gravity-capillary water waves. *Mem. Am. Math. Soc.*, Memo 891 (to appear)
17. Berti, M., Biasco, L., Procesi, M.: KAM for reversible derivative wave equations. *Arch. Ration. Mech. Anal.* **212**(3), 905–955 (2014). <https://doi.org/10.1007/s00205-014-0726-0>
18. Beyer, K., Günther, M.: On the Cauchy problem for a capillary drop. I. Irrotational motion. *Math. Methods Appl. Sci.* **21**(12), 1149–1183 (1998). [https://doi.org/10.1002/\(SICI\)1099-1476\(199808\)21:12<1149::AID-MMA990>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1099-1476(199808)21:12<1149::AID-MMA990>3.0.CO;2-C)
19. Boutet de Monvel, L.: Comportement d'un opérateur pseudo-différentiel sur une variété à bord. I. La propriété de transmission. *J. Anal. Math.* **17**, 241–253 (1966)
20. Boutet de Monvel, L.: Comportement d'un opérateur pseudo-différentiel sur une variété à bord. II. Pseudo-noyaux de Poisson. *J. Anal. Math.* **17**, 255–304 (1966)
21. Boutet de Monvel, L.: Boundary problems for pseudo-differential operators. *Acta Math.* **126**(1–2), 11–51 (1971)
22. Broer, H.W., Huitema, G.B., Sevryuk, M.B.: Quasi-Periodic Motions in Families of Dynamical Systems. *Lecture Notes in Mathematics*, vol. 1645. Springer, Berlin (1996). Order amidst chaos
23. Coutand, D., Shkoller, S.: Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J. Am. Math. Soc.* **20**(3), 829–930 (2007). <https://doi.org/10.1090/S0894-0347-07-00556-5>
24. Craig, W.: An existence theory for water waves and the Boussinesq and Korteweg-de Vries scaling limits. *Commun. Partial Differ. Equ.* **10**(8), 787–1003 (1985). <https://doi.org/10.1080/03605308508820396>
25. Craig, W., Nicholls, D.P.: Travelling two and three dimensional capillary gravity water waves. *SIAM J. Math. Anal.* **32**(2), 323–359 (2000). <https://doi.org/10.1137/S0036141099354181>
26. Craig, W., Sulem, C.: Numerical simulation of gravity waves. *J. Comput. Phys.* **108**(1), 73–83 (1993). <https://doi.org/10.1006/jcph.1993.1164>
27. Craig, W., Sulem, C.: Normal form transformations for capillary-gravity water waves. In: *Hamiltonian Partial Differential Equations and Applications*. Fields Institute Communications, vol. 75, pp. 73–110. Fields Institute for Research in Mathematical Sciences, Toronto (2015). https://doi.org/10.1007/978-1-4939-2950-4_3
28. Craig, W., Worfolk, P.A.: An integrable normal form for water waves in infinite depth. *Physica D* **84**(3–4), 513–531 (1995). [https://doi.org/10.1016/0167-2789\(95\)00067-E](https://doi.org/10.1016/0167-2789(95)00067-E)
29. Delort, J.-M.: A quasi-linear Birkhoff normal forms method. Application to the quasi-linear Klein-Gordon equation on S^1 . *Astérisque* **341**, vi+113 (2012)
30. Delort, J.-M.: Quasi-linear perturbations of Hamiltonian Klein-Gordon equations on spheres. *Mem. Am. Math. Soc.* **234**(1103), vi+80 (2015). <https://doi.org/10.1090/memo/1103>
31. Delort, J.-M., Szeftel, J.: Long-time existence for small data nonlinear Klein-Gordon equations on tori and spheres. *Int. Math. Res. Not.* **37**, 1897–1966 (2004). <https://doi.org/10.1155/S1073792804133321>
32. Deng, Y., Ionescu, A.D., Pausader, B., Pusateri, F.: Global solutions of the gravity-capillary water-wave system in three dimensions. *Acta Math.* **219**(2), 213–402 (2017). <https://doi.org/10.4310/ACTA.2017.v219.n2.a1>
33. Dimassi, M., Sjöstrand, J.: *Spectral Asymptotics in the Semi-classical Limit*. London Mathematical Society Lecture Note Series, vol. 268. Cambridge University Press, Cambridge (1999). <https://doi.org/10.1017/CBO9780511662195>
34. D'yachenko, A.I., Zakharov, V.E.: Is free-surface hydrodynamics an integrable system? *Phys. Lett. A* **190**(2), 144–148 (1994). [https://doi.org/10.1016/0375-9601\(94\)90067-1](https://doi.org/10.1016/0375-9601(94)90067-1)
35. Fang, D., Han, Z., Zhang, Q.: Almost global existence for the semi-linear Klein-Gordon equation on the circle. *J. Differ. Equ.* **262**(9), 4610–4634 (2017). <https://doi.org/10.1016/j.jde.2016.12.013>

36. Faou, E., Grébert, B.: Quasi-invariant modified Sobolev norms for semi linear reversible PDEs. *Nonlinearity* **23**(2), 429–443 (2010). <https://doi.org/10.1088/0951-7715/23/2/011>
37. Germain, P., Masmoudi, N., Shatah, J.: Global solutions for the gravity water waves equation in dimension 3. *Ann. Math. (2)* **175**(2), 691–754 (2012). <https://doi.org/10.4007/annals.2012.175.2.6>
38. Germain, P., Masmoudi, N., Shatah, J.: Global existence for capillary water waves. *Commun. Pure Appl. Math.* **68**(4), 625–687 (2015). <https://doi.org/10.1002/cpa.21535>
39. Giorgilli, A.: Rigorous results on the power expansions for the integrals of a Hamiltonian system near an elliptic equilibrium point. *Ann. Inst. H. Poincaré Phys. Théor.* **48**(4), 423–439 (1988). http://www.numdam.org/item?id=AIHPB_1988__48_4_423_0
40. Giorgilli, A., Posilicano, A.: Estimates for normal forms of differential equations near an equilibrium point. *Z. Angew. Math. Phys.* **39**(5), 713–732 (1988). <https://doi.org/10.1007/BF00948732>
41. Harrop-Griffiths, B., Ifrim, M., Tataru, D.: Finite depth gravity water waves in holomorphic coordinates. *Ann. PDE* **3**(1), Art. 4, 102 (2017). <https://doi.org/10.1007/s40818-017-0022-z>
42. Hunter, J.K., Ifrim, M., Tataru, D.: Two dimensional water waves in holomorphic coordinates. *Commun. Math. Phys.* **346**(2), 483–552 (2016). <https://doi.org/10.1007/s00220-016-2708-6>
43. Ifrim, M., Tataru, D.: Two dimensional gravity water waves with constant vorticity: I. Cubic lifespan. Preprint (2014). <https://arxiv.org/abs/1510.07732>
44. Ifrim, M., Tataru, D.: Two dimensional water waves in holomorphic coordinates II: Global solutions. *Bull. Soc. Math. Fr.* **144**(2), 369–394 (2016)
45. Ifrim, M., Tataru, D.: The lifespan of small data solutions in two dimensional capillary water waves. *Arch. Ration. Mech. Anal.* **225**(3), 1279–1346 (2017). <https://doi.org/10.1007/s00205-017-1126-z>
46. Ionescu, A.D., Pusateri, F.: Global regularity for 2D water waves with surface tension. Preprint (2014). <https://arxiv.org/abs/1408.4428>
47. Ionescu, A.D., Pusateri, F.: Global solutions for the gravity water waves system in 2D. *Invent. Math.* **199**(3), 653–804 (2015). <https://doi.org/10.1007/s00222-014-0521-4>
48. Ionescu, A.D., Pusateri, F.: Long-time existence for multi-dimensional periodic water waves. Preprint (2018). <https://arxiv.org/abs/1807.02932>
49. Iooss, G., Plotnikov, P.: Existence of multimodal standing gravity waves. *J. Math. Fluid Mech.* **7**(suppl. 3), S349–S364 (2005). <https://doi.org/10.1007/s00021-005-0164-8>
50. Iooss, G., Plotnikov, P.: Multimodal standing gravity waves: a completely resonant system. *J. Math. Fluid Mech.* **7**(suppl. 1), S110–S126 (2005). <https://doi.org/10.1007/s00021-004-0128-4>
51. Iooss, G., Plotnikov, P.I.: Small divisor problem in the theory of three-dimensional water gravity waves. *Mem. Am. Math. Soc.* **200**(940), viii+128 (2009). <https://doi.org/10.1090/memo/0940>
52. Iooss, G., Plotnikov, P.: Asymmetrical three-dimensional travelling gravity waves. *Arch. Ration. Mech. Anal.* **200**(3), 789–880 (2011). <https://doi.org/10.1007/s00205-010-0372-0>
53. Iooss, G., Plotnikov, P.I., Toland, J.F.: Standing waves on an infinitely deep perfect fluid under gravity. *Arch. Ration. Mech. Anal.* **177**(3), 367–478 (2005). <https://doi.org/10.1007/s00205-005-0381-6>
54. Lannes, D.: Well-posedness of the water-waves equations. *J. Am. Math. Soc.* **18**(3), 605–654 (electronic) (2005). <https://doi.org/10.1090/S0894-0347-05-00484-4>
55. Lannes, D.: *The Water Waves Problem. Mathematical Surveys and Monographs*, vol. 188. American Mathematical Society, Providence, RI (2013). <https://doi.org/10.1090/surv/188>. Mathematical analysis and asymptotics
56. Levi-Civita, T.: Détermination rigoureuse des ondes permanentes d'ampleur finie. *Math. Ann.* **93**(1), 264–314 (1925). <https://doi.org/10.1007/BF01449965>
57. Lindblad, H.: Well-posedness for the motion of an incompressible liquid with free surface boundary. *Ann. Math. (2)* **162**(1), 109–194 (2005). <https://doi.org/10.4007/annals.2005.162.109>

58. Ming, M., Zhang, Z.: Well-posedness of the water-wave problem with surface tension. *J. Math. Pures Appl.* (9) **92**(5), 429–455 (2009). <https://doi.org/10.1016/j.matpur.2009.05.005>
59. Moser, J.: Convergent series expansions for quasi-periodic motions. *Math. Ann.* **169**, 136–176 (1967)
60. Nalimov, V.I.: The Cauchy-Poisson problem. *Dinamika Splošn. Sredy* (Vyp. 18 Dinamika Zidkost. so Svobod. Granicami), 104–210, 254 (1974)
61. Plotnikov, P.I., Toland, J.F.: Nash-Moser theory for standing water waves. *Arch. Ration. Mech. Anal.* **159**(1), 1–83 (2001). <https://doi.org/10.1007/PL00004246>
62. Schweizer, B.: On the three-dimensional Euler equations with a free boundary subject to surface tension. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22**(6), 753–781 (2005). <https://doi.org/10.1016/j.anihpc.2004.11.001>
63. Wu, S.: Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.* **130**(1), 39–72 (1997). <https://doi.org/10.1007/s002220050177>
64. Wu, S.: Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Am. Math. Soc.* **12**(2), 445–495 (1999). <https://doi.org/10.1090/S0894-0347-99-00290-8>
65. Wu, S.: Almost global wellposedness of the 2-D full water wave problem. *Invent. Math.* **177**(1), 45–135 (2009). <https://doi.org/10.1007/s00222-009-0176-8>
66. Wu, S.: Global wellposedness of the 3-D full water wave problem. *Invent. Math.* **184**(1), 125–220 (2011). <https://doi.org/10.1007/s00222-010-0288-1>
67. Yosihara, H.: Gravity waves on the free surface of an incompressible perfect fluid of finite depth. *Publ. Res. Inst. Math. Sci.* **18**(1), 49–96 (1982). <https://doi.org/10.2977/prims/1195184016>
68. Zakharov, V.E.: Stability of periodic waves of finite amplitude on the surface of a deep fluid. *J. Appl. Mech. Tech. Phys.* **9**(2), 190–194 (1968)
69. Zhang, J., Gao, M., Yuan, X.: KAM tori for reversible partial differential equations. *Nonlinearity* **24**(4), 1189–1228 (2011). <https://doi.org/10.1088/0951-7715/24/4/010>

Index

- $A(U; t, x, \xi)^\vee$, 94
- $(a\#b)_\infty$ (Composition of symbols), 54, 55
- $a\#b$ (Composition of symbols), 56
- $(a\#b)_{\rho, N}$ (Composition of symbols), 64
- $\text{Ad}_{O\rho^{\text{BW}}(F(U))}$ (Commutator), 124

- $B_\sigma^k(I, r)$ (Ball in the space of functions of (t, x)), 34
- $B_{\sigma_0}^k(I, r, j)$ (Ball in the space of functions of (t, z, x)), 159

- $C^K(I, \dot{H}^\sigma(\mathbb{T}^1, \mathbb{C}^2))$ (Space of functions), 32
- $C_{*\mathbb{R}}^k(I, \dot{H}^\sigma(\mathbb{T}^1, \mathbb{C}^2))$ (Space of functions), 32
- Reality conditions, 94
- Parity preserving condition, 94
- Reversibility conditions, 94

- $\mathcal{E}_{K, K', 1}^m[r, N]$ (Space of matrices of symbols), 100
- E_j^s (Space of functions on a strip), 165

- Φ^* (Paracomposition operator), 88
- $\tilde{\mathcal{F}}_p$ (Space of homogeneous functions), 73
- $\mathcal{F}_{K, K', p}[r]$ (Space of non-homogeneous functions), 73
- $\Sigma \mathcal{F}_{K, K', p}[r, N]$ (Space of functions), 73
- $\Sigma \mathcal{F}_{K, K', p}^{\mathbb{R}}[r, N]$ (Space of real functions), 73
- $\Sigma \mathcal{F}_{K, 0, p}[r, N, \text{aut}]$ (Space of autonomous functions), 74
- F_j^s (Space of functions on a strip), 158

- $G(\eta)$ (Dirichlet-Neumann operator), 28
- $\tilde{\Gamma}_p^m$ (Space of homogeneous symbols), 33
- $\Gamma_{K, K', p}^m[r]$ (Space of non-homogeneous symbols), 34
- $\Gamma_{K, 0, p}^m[r, \text{aut}]$ (Space of autonomous symbols), 35
- $\Sigma \Gamma_{K, K', p}^m[r, N]$ (Space of symbols), 36
- $\Sigma \Gamma_{K, 0, p}^m[r, N, \text{aut}]$ (Space of autonomous symbols), 36
- $\tilde{\Gamma}_p^m[\mathcal{B}]$ (Space of homogeneous symbols in a strip), 158
- $\Gamma_{K, 0, j, p}^m[\mathcal{B}, r]$ (Space of non-homogeneous symbols in a strip), 159
- $\Sigma \Gamma_{K, 0, j, p}^m[\mathcal{B}, r, N]$ (Space of symbols in a strip), 159
- $\mathcal{G}_{K, p}^\sigma(U)$ (Norm at power p), 32
- $\mathcal{G}_{0, p}^\sigma(\mathcal{U})$ (Product of norms), 32
- $\mathcal{G}_{K, p}^{\sigma, j}(\Phi, t)$ (Power of norm of a function on a strip), 158
- $\mathcal{G}_{0, p}^{\sigma, j}(\Phi)$ (Product of norms of functions on a strip), 158
- $\tilde{\mathcal{G}}_{K, 1}^{\sigma, j}(U, t)$ (Norm on a strip), 169

- $\text{Im } A_q$ (Imaginary part of a matrix of symbols), 133
- $K_0(z, z', \xi)$ (Poisson kernel), 164

- $\tilde{\mathcal{L}}_{p, \pm}^{\sigma, -\rho}$ (Space of multilinear energy forms), 143

- $\widetilde{\mathcal{M}}_p^m$ (Space of homogeneous maps), 51
 $\mathcal{M}_{K,K',N}^m[r]$ (Space of non-homogeneous maps), 52
 $\mathcal{M}_{K,0,N}^m[r]$ (Space of autonomous maps), 52
 $\Sigma \mathcal{M}_{K,K',p}^m[r, N]$ (Space of maps), 52
 $\Sigma \mathcal{M}_{K,0,p}^m[r, N, \text{aut}]$ (Space of autonomous maps), 52
 $\widetilde{\mathcal{M}}_p$ (Space of homogeneous maps), 52
 $\mathcal{M}_{K,K',p}[r]$ (Space of non-homogeneous maps), 52
 $\Sigma \mathcal{M}_{K,K',p}[r, N]$ (Space of maps), 52
- \mathcal{N}_p^m (Space of homogeneous maps), 203
 $\mathcal{N}_{K,j,p}^m[r]$ (Space of non-homogeneous maps), 203
 $\Sigma \mathcal{N}_{K,j,p}^m[r, N]$ (Space of maps), 204
 $\widetilde{\mathcal{N}}_p^{\partial,m}$ (Space of maps), 205
 $\mathcal{N}_{K,j,N}^{\partial,m}[r]$ (Space of maps), 205
 $\Sigma \mathcal{N}_{K,j,p}^{\partial,m}[r, N]$ (Space of maps), 205
- Op_τ (τ -quantization), 37
 Op^{BW} (Bony-Weyl quantization), 43
 Op^{W} (Weyl quantization), 37
- $\widetilde{\mathcal{P}}_p^{m,\pm}$ (Poisson homogeneous symbols from the boundary to the interior), 162
 $\widetilde{\mathcal{P}}_p^{m,\text{int}}$ (Poisson homogeneous symbols from the interior to the interior), 162
 $\mathcal{P}_{K,0,p}^{m,\pm}[r]$ (Poisson non-homogeneous symbols from the boundary to the interior), 163
 $\mathcal{P}_{K,0,p}^{m,\text{int}}[r]$ (Poisson non-homogeneous symbols from the interior to the interior), 163
 $\Sigma \mathcal{P}_{K,0,p}^{m,\pm}[r, N]$ (Poisson symbols from the boundary to the interior), 163
 $\Sigma \mathcal{P}_{K,0,p}^{m,\text{int}}[r, N]$ (Poisson symbols from the interior to the interior), 163
- $\widetilde{\mathcal{R}}_p^{-\rho}$ (Homogeneous smoothing operators), 44
 $\mathcal{R}_{K,K',N}^{-\rho}[r]$ (Non-homogeneous smoothing operators), 45
 $\mathcal{R}_{K,0,N}^{-\rho}[r, \text{aut}]$ (Autonomous non-homogeneous smoothing operators), 45
 $\Sigma \mathcal{R}_{K,K',p}^{-\rho}[r, N]$ (Smoothing operators), 45
 $\Sigma \mathcal{R}_{K,0,p}^{-\rho}[r, N, \text{aut}]$ (Autonomous smoothing operators), 45
 $\widetilde{\mathcal{R}}_p^{-\rho}[\mathcal{B}]$ (Space of homogeneous smoothing operators in a strip), 160
 $\mathcal{R}_{K,0,j,N}^{-\rho}[\mathcal{B}, r]$ (Space of non-homogeneous smoothing operators in a strip), 161
 $\Sigma \mathcal{R}_{K,0,j,p}^{-\rho}[\mathcal{B}, r, N]$ (Space of smoothing operators in a strip), 161
 $\widetilde{\mathcal{R}}_p^{-\rho,\pm}$ (Space of homogeneous smoothing operators from the boundary to a strip), 169
 $\widetilde{\mathcal{R}}_p^{-\rho,\text{int}}$ (Space of homogeneous smoothing operators on a strip), 169
 $\mathcal{R}_{K,0,N}^{-\rho,\pm}[r]$ (Space of non-homogeneous smoothing operators from the boundary to a strip), 170
 $\mathcal{R}_{K,0,N}^{-\rho,\text{int}}[r]$ (Space of smoothing operators on a strip), 170
 $\Sigma \mathcal{R}_{K,0,p}^{-\rho,\pm}[r, N]$ (Space of smoothing operators from the boundary to a strip), 170
 $\Sigma \mathcal{R}_{K,0,p}^{-\rho,\text{int}}[r, N]$ (Space of smoothing operators on a strip), 170
- S (Reversibility operator in real form), 29
 S (Reversibility operator in complex form), 93
 $\sigma(D_x, D_\xi, D_y, D_\eta)$, 54
- U_S (Action of reversibility operator S on U), 93

Editor in Chief: *Ciro Ciliberto and Susanna Terracini*

Editorial Policy

1. The UMI Lecture Notes aim to report new developments in all areas of mathematics and their applications - quickly, informally and at a high level. Mathematical texts analysing new developments in modelling and numerical simulation are also welcome.
2. Manuscripts should be submitted to
Redazione Lecture Notes U.M.I.
umi@dm.unibo.it
and possibly to one of the editors of the Board informing, in this case, the Redazione about the submission. In general, manuscripts will be sent out to external referees for evaluation. If a decision cannot yet be reached on the basis of the first 2 reports, further referees may be contacted. The author will be informed of this. A final decision to publish can be made only on the basis of the complete manuscript, however a refereeing process leading to a preliminary decision can be based on a prefinal or incomplete manuscript. The strict minimum amount of material that will be considered should include a detailed outline describing the planned contents of each chapter, a bibliography and several sample chapters.
3. Manuscripts should in general be submitted in English. Final manuscripts should contain at least 100 pages of mathematical text and should always include
 - a table of contents;
 - an informative introduction, with adequate motivation and perhaps some historical remarks: it should be accessible to a reader not intimately familiar with the topic treated;
 - a subject index: as a rule this is genuinely helpful for the reader.
4. For evaluation purposes, please submit manuscripts in electronic form, preferably as pdf- or zipped ps-files. Authors are asked, if their manuscript is accepted for publication, to use the LaTeX2e style files available from Springer's web-server at
<ftp://ftp.springer.de/pub/tex/latex/svmonot1/> for monographs
and at
<ftp://ftp.springer.de/pub/tex/latex/svmult1/> for multi-authored volumes
5. Authors receive a total of 50 free copies of their volume, but no royalties. They are entitled to a discount of 33.3% on the price of Springer books purchased for their personal use, if ordering directly from Springer.
6. Commitment to publish is made by letter of intent rather than by signing a formal contract. Springer-Verlag secures the copyright for each volume. Authors are free to reuse material contained in their LNM volumes in later publications: A brief written (or e-mail) request for formal permission is sufficient.