

References

- [1] Aikman, D.P. and Anderson, W.P. (1971), *Ann. Bot.* **35**, 61–72.
- [2] Allaway, W.G. (1973), *Planta*, **110**, 63–70.
- [3] Anderson, R. and Cronshaw, J. (1969), *J. Ultrastruct. Res.*, **29**, 50–59.
- [4] Anderson, W.P. (ed) (1973), *Ion Transport in Plants*, Academic Press, London and New York.
- [5] Anderson, W.P. (1974), *Symp. Soc. Exptl. Biol.*, **28**, 63–85.
- [6] Anderson, W.P. (1975), in [11] p. 231.
- [7] Atkinson, M.R. *et al.* (1967), *Aust. J. Biol. Sci.*, **20**, 589–599.
- [8] Baker, D.A. (1968), *Planta*, **83**, 390–392.
- [9] Baker, D.A. (1973), *Planta*, **112**, 293–299.
- [10] Baker, D.A. and Hall, J.L. (1973), *New Phytol.*, **72**, 1281–1291.
- [11] Baker, D.A. and Hall, J.L. (eds) (1975), *Ion Transport in Plant Cells and Tissues*, North Holland, Netherlands.
- [12] Baker, D.A. and Weatherley, P.E. (1969), *J. Exptl. Botany*, **20**, 485–496.
- [13] Barry, W.H. (1968), *J. Cell Physiol.* **72**, 153–160.
- [14] Bell, C.W. and Biddulph, O. (1963), *Plant Physiol.* **38**, 610–614.
- [15] Ben Zioni, A., *et al.* (1971), *Physiologia Plantarum*, **24**, 288–290.
- [16] Biddulph, O., and Gory, R. (1960), *Plant Physiol.* **35**, 689–695.
- [17] Bollard, E.G. (1960), *Ann. Rev. Plant Physiol.* **11**, 141–166.
- [18] Bowen, M.R. and Wareing, P.F. (1969), *Planta*, **89**, 108–125.
- [19] Bowling, D.J.F. (1969), *Biochim. Biophys. Acta.* **183**, 230–232.
- [20] Bowling, D.J.F. (1973), *J. Exptl. Botany*, **24**, 1041–1045.
- [21] Bowling, D.J.F. (1973), in [4] p. 483.
- [22] Bowling, D.J.F. (1976), *Uptake of Ions by Plant Roots*. Chapman and Hall, London.
- [23] Bowling, D.J.F. (1976), *Nature*, Lond. **262**, 393–394.
- [24] Bowling, D.J.F. *et al.* (1966), *J. Exptl. Botany*, **17**, 410–416.
- [25] Canny, M.J. (1973), *Phloem Translocation*, Cambridge University Press, London.
- [26] Clarkson, D.T. (1974), *Ion Transport and Cell Structure in Plants*, McGraw-Hill, Maidenhead.
- [27] Clarkson, D.T. *et al.* (1971), *Planta*, **96**, 296–305.
- [28] Collins, J.C. and Kerrigan, A.P. (1973), in [4] p. 589.
- [29] Crafts, A.S. and Broyer, T.C. (1938), *Amer. J. Botany*, **25**, 529–535.
- [30] Crafts, A.S. and Crisp, C.E. (1971), *Phloem Transport in Plants*, Freeman, San Francisco.
- [31] Cronshaw, J. (1974), in Robards, A.W. (ed) *Dynamic Aspects of Plant Ultrastructure*. p. 391 McGraw-Hill, Maidenhead.
- [32] Cronshaw, J. and Esau, K. (1967), *J. Cell. Biol.* **34**, 801–816.
- [33] Dixon, H.H. and Ball, N.G. (1922), *Nature*, **109**, 236–237.
- [34] Dunlop, J. and Bowling, D.J.F. (1971), *J. Exptl. Botany*, **22**, 434–444.
- [35] Epstein, E. (1973), *Int. Rev. Cytol.* **34**, 123–168.
- [36] Epstein, E. (1973), *Experientia*, **29**, 133–134.
- [37] Eschrich, W. (1967), *Planta*, **73**, 37–49.

- [38] Eschrich, W. (1975), in [142] p. 245.
- [39] Eschrich, W. and Heyser, W. (1975), in [142] p. 101.
- [40] Fensom, D.S. (1957), *Can. J. Botany*, **35**, 573–582.
- [41] Fensom, D.S. (1972), *Can. J. Botany*, **50**, 479–497.
- [42] Fritz, E. (1973), *Planta*, **112**, 169–179.
- [43] Fujino, M. (1967), *Sci. Bull. Fac. Educat. Nagasaki Univ.* **18**, 1–47.
- [44] Geiger, D.R. and Sovonick, S.A. (1975), in [142] p. 256.
- [45] Gerson, D.F. and Poole, R.J. (1971), *Plant Physiol.* **48**, 509–511.
- [46] Giaquinta, R. and Geiger, D.R. (1977), *Plant Physiol.* **59**, 178–180.
- [47] Ginsburg, H. and Ginsburg, B.Z. (1970), *J. Exptl. Botany*, **21**, 593–604.
- [48] Gunning, B.E.S. and Pate, J.S. (1969), *Protoplasma*, **68**, 107–133.
- [49] Hall, J.L. and Baker, D.A. (1975), in [11] p. 39.
- [50] Hall, J.L. and Baker, D.A. (1977), *Membranes and Ion Transport*, Longman, London.
- [51] Hall, S.M. and Baker, D.A. (1972), *Planta*, **106**, 131–140.
- [52] Hall, S.M. and Medlow, G.C. (1974), *Planta*, **119**, 257–261.
- [53] Hartig, T. (1860), *Allgem. Forst-u-Jagdztg*, **36**, 257–261.
- [54] Hartt, C.E. (1965), *Plant Physiol.* **40**, 718–724.
- [55] Hartt, C.E. (1969), *Plant Physiol.* **44**, 1461–1469.
- [56] Hartt, C.E. (1970), *Plant Physiol.* **45**, 183–187.
- [57] Hartt, C.E. and Kortschak, H.P. (1967), *Plant Physiol.* **42**, 89–94.
- [58] Hay, R.K.M. (1974), in Kolek, J. (ed) *Structure and Function of Primary Root Tissues*, p. 329, Slovak Academy of Science.
- [59] Heine, H. (1885), *Ber. Deutsch. Bot. Ges.* **3**, 189–194.
- [60] Hejnowicz, Z. (1970), *Protoplasma*, **71**, 343–364.
- [61] Hoad, G.V. (1973), *Planta*, **113**, 367–372.
- [62] Hope, A.B. (1971), *Ion Transport and Membranes*, Butterworths, London.
- [63] House, C.R. and Findlay, N. (1966), *J. exptl. Botany*, **17**, 344–354.
- [64] Huber, B. (1956) in Ruhland, W. (ed) *Hanb. Pflanzenphysiol.* Vol. III, p. 511, Springer-Verlag, Berlin.
- [65] Humble, G.D. and Hsiao, T.C. (1969), *Plant Physiol.* **44**, 230–234.
- [66] Humble, G.D. and Raschke, K. (1971), *Plant Physiol.* **48**, 447–453.
- [67] Isermann, K. (1971), *Z. Pflanzenr Bodenkunde*, **128**, 195–207.
- [68] Jacoby, B. (1965), *Physiol. Plant*, **18**, 730–739.
- [69] Jaffe, M.J. (1970), *Plant. Physiol.* **46**, 768–777.
- [70] Jeschke, W.D. and Simonis, W. (1969), *Planta*, **88**, 157–171.
- [71] Koukkari, W.L. and Hillman, W.S. (1968), *Plant Physiol.* **43**, 698–704.
- [72] Kursanov, A.L. (1963), *Advan. Botan. Res.* **1**, 209–274.
- [73] Laties, G.G. (1969), *Ann. Rev. Plant Physiol.* **20**, 89–116.
- [74] Läuchli, A. (1972), *Ann. Rev. Plant. Physiol.* **23**, 197–218.
- [75] Lüttge, U. (1975), in [11] p. 335.
- [76] Lüttge, U. and Pitman, M.G. (eds) (1976), *Encyclopedia of Plant Physiology* N.S. Volume 2A, Springer-Verlag, Berlin, Heidelberg, New York.
- [77] Lüttge, U. and Pitman, M.G. (ed) (1976), *Encyclopedia of Plant Physiology* N.S. Volume 2B, Springer-Verlag, Berlin, Heidelberg, New York.
- [78] Macallum, A.B. (1905), *J. Physiol.* **32**, 95–118.
- [79] MacRobbie, E.A.C. (1971), *Biol. Rev.*, **46**, 429–481.
- [80] MacRobbie, E.A.C. (1971), *Ann. Rev. Plant Physiol.* **22**, 75–96.
- [81] Malek, F. and Baker, D.A. (1977), *Planta*, **135**, 297–299.
- [82] Mason, T.G. *et al.* (1936), *Ann. Bot.* **50**, 23–58.

- [83] Mees, G.C. and Weatherley, P.E. (1957), *Proc. Roy. Soc. B.* **147**, 381–391.
- [84] Milburn, J.A. and McLaughlin, M.E. (1974), *New Phytol.* **73**, 861–871.
- [85] Mullins, M.G. (1970), *Ann. Bot.* **34**, 897–909.
- [86] Münch, E. (1930), *Die Stoffbewegungen in der Pflanze*, Jena: Fischer.
- [87] Nelson, G.D. and Gorham, P.R. (1957), *Can. J. Bot.* **35**, 339–347.
- [88] Nelson, C.D. *et al.* (1961), *Plant Physiol.* **36**, 581–588.
- [89] Nissen, P. (1974), *Ann. Rev. Plant Physiol.* **25**, 53–79.
- [90] Parker, B.C. (1965), *J. Physiol.* **1**, 41–46.
- [91] Parthasarathy, M.V. and Mühlethaler, K. (1969), *Cytobiologie*, **1**, 7–36.
- [92] Pate, J.S. *et al.* (1974), *Planta*, **120**, 229–243.
- [93] Peel, A.J. and Weatherley, P.E. (1959), *Nature*, **184**, 1955–1956.
- [94] Penny, M.G. and Bowling, D.J.F. (1974), *Planta*, **119**, 17–25.
- [95] Philips, D.A. and Cleland, C.F. (1972), *Planta*, **102**, 173–178.
- [96] Pitman, M.G. (1972), *Aust. J. Biol. Sci.* **25**, 243–257.
- [97] Pitman, M.G. (1975), in [11] p. 267.
- [98] Postlethwait, S.N. and Rogers, B. (1958), *Am. J. Botany*, **45**, 753–757.
- [99] Preston, R.D. (1958) in Everett, D.H. and Stone, F.S. (eds) *The Structure and Properties of Porous Materials*, p. 366 Butterworths, London.
- [100] Qureshi, F.A. and Spanner, D.C. (1973), *J. Exptl. Botany*, **24**, 751–762.
- [101] Raschke, K. and Fellows, M.P. (1971), *Planta*, **101**, 296–316.
- [102] Raschke, K. and Humble, G.D. (1973), *Planta*, **115**, 47–57.
- [103] Raven, J.A. (1967), *J. Gen. Physiol.* **50**, 1607–25.
- [104] Raven, J.A. (1975), in [11] p. 125.
- [105] Robards, A.W. and Robb, M.E. (1972), *Science*, **178**, 980–982.
- [106] Salter, R.L. *et al.* (1970), *Am. J. Botany*, **57**, 916–926.
- [107] Schumacher, W. (1930), *Jahrb. Wiss. Bot.* **73**, 770–823.
- [108] Shone, M.G.T. *et al.* (1969), *Planta*, **86**, 301–314.
- [109] Siddiqui, A.W. and Spanner, D.C. (1970), *Planta*, **91**, 181–189.
- [110] Slatyer, R.O. (1967), *Plant-Water Relationships*, Academic Press, New York.
- [111] Smith, F.A. (1970), *New Phytol.* **69**, 903–917.
- [112] Smith, R.C. and Epstein, E. (1964), *Plant Physiol.* **39**, 338–341.
- [113] Spanner, D.C. (1958), *J. Exptl. Botany*, **9**, 332–342.
- [114] Spanner, D.C. and Jones, R.L. (1970), *Planta*, **92**, 64–72.
- [115] Tamulevich, S.R. and Evert, R.F. (1966), *Planta*, **69**, 319–337.
- [116] Thaine, R. (1962), *J. Exptl. Botany*, **13**, 152–160.
- [117] Thellier, M. (1973), in [4] p. 47.
- [118] Thomas, D.A. (1975), in [11] p. 377.
- [119] Thompson, R.G. and Thompson, A.D. (1973), *Can. J. Botany*, **51**, 933–936.
- [120] Trip, P. and Gorham, P.R. (1968), *Plant Physiol.* **43**, 877–882.
- [121] Tukey, H.B. Jr. (1970), *Ann. Rev. Plant Physiol.* **21**, 305–324.
- [122] Tyree, M.T. (1970), *J. Theor. Biol.* **26**, 181–214.
- [123] Tyree, M.T. and Fensom, D.S. (1970), *J. Exptl. Botany*, **21**, 304–324.
- [124] Van Die, J. and Tammes, P.M.L. (1975), in [142] p. 196.
- [125] Vernon, L.P. and Aronoff, S. (1952), *Arch. Biochem. Biophys.* **36**, 383–398.

- [126] Weatherley, P.E. and Johnson, R.P.C. (1968), *Int. Rev. Cytol.* **24**, 149–192.
- [127] Weatherley, P.E. *et al.* (1959), *J. Exptl. Botany*, **10**, 1–16.
- [128] Williams, S.E. and Spanswick, R.M. (1976), *J. comp. Physiol.* **108**, 211–223.
- [129] Williamson, R.E. (1972), *Planta*, **106**, 149–157.
- [130] Winter, K. and Lüttge, U. (1976), *Aust. J. Plant Physiol.* **3**, 653–664.
- [131] Wooding, F.P.B. (1971), *Phloem*, Oxford Biology Readers, Oxford University Press.
- [132] Wyn Jones, R.G. (1975), in [11] p. 193.
- [133] Yeo, A.R. *et al.* (1977), *J. Exptl. Botany*, **28**, 17–29.
- [134] Yu, G.H. and Kramer, P.J. (1969), *Plant Physiol.* **44**, 1095–1100.
- [135] Ziegler, H. (1974), *Symp. Soc. Exptl. Biol.* **28**, 43–62.
- [136] Ziegler, H. (1975), in [142] p. 59.
- [137] Ziegler, H. and Kluge, M. (1962), *Planta*, **58**, 144–153.
- [138] Zimmermann, M.H. (1958), *Plant Physiol.* **33**, 213–217.
- [139] Zimmermann, M.H. (1963), *Scient. Am.* **208**, March, 132–142.
- [140] Zimmermann, M.H. (1964), *Plant Physiol.* **39**, 568–572.
- [141] Zimmermann, M.H. (1969), in Wilkins, M.B. (ed) *Physiology of Plant Growth and Development*, p. 383, McGraw-Hill, London.
- [142] Zimmermann, M.H. and Milburn, J.A. (eds) (1975), *Encyclopedia of Plant Physiology* N.S. Volume 1, Springer-Verlag, Berlin, Heidelberg, New York.

Index

- Abscisic acid (ABA), 29, 30, 34, 47, 52
Actin, 70, 71
Active transport, 10–11, 15
Agialitis annulata, see mangrove
Ajugose, 47, 48
Albuminous cells, 42
Amino acids, 34, 47–49, 53, 54
Aphids, 40, 47, 57
Apparent free space (AFS), 23
Apoplast, 23–31, 38, 58, 59
Aristolochia siphon, 64
ATP (Adenosine triphosphate), 15–17, 47, 49, 53, 59
ATPase, 17, 49
Atragene alpina, 64
Auxin (IAA), 34, 47, 51, 52, 62
- Barley, 35
Beech, see *Fagus*
Benzyl-adenine, 62
Beta vulgaris, 55
Bidirectional movement, 56, 57, 68
Birch, 37
Boron, 61
Brush border, 69
- Calcium
 ethylenediaminetetra
 acetic acid (CaEDTA), 35
 fluxes, 11, 30
 levels in various
 materials, 8, 35, 37, 47, 50
 oxalate crystals, 36
 transport across root, 27, 28
Callose, 40, 41, 43, 70
Carbon dioxide, 16, 18–20, 40, 47, 49, 53, 57, 60, 61
Carbon monoxide, 12
Carbonylcyano *m* chlorophenyl
 hydrazone (CCCP), 17
Carriers, 12–15
Casparian band, 26–28
- Cavitation, 64, 66
Chemical coupling hypothesis, 16
Chemical potential, 9
Chemiosmotic hypothesis, 16
Chloride
 fluxes, 10, 11, 15–18, 36
 levels in various
 materials, 8, 19, 21, 35, 37, 47
 permeability coefficient, 11
 transport in symplasm, 28
Chloroplasts, 15, 17
Cohesion theory, 63–66
Commelina communis, 21
Companion cells, 30, 41, 42, 58
Conductance, 11
Conductivity, 54
Copper, 51
Co-transport of sugars, 59
Crassulacean acid metabolism (CAM), 20
Cucurbita melopepo torticollis, 55
Cucurbita pepo, 55
Cyclosis, see cytoplasmic streaming
Cytokinins, 34, 47, 52, 62
Cytoplasmic streaming (cyclosis), 24, 41, 58
- Dendrograph, 65
Dichlorophenyldimethyl urea (DCMU), 17
Dictyosomes, 41, 42
Diffusion potential, 10, 11
DNA, 53
Donnan free space (DFS), 23
Driving forces, 9–12, 35
- Electrical potential, 9, 25
Electrochemical potential, 9, 25
Electron probe microanalyser (EPM), 19–21, 24, 34
Electron transport system, 15
Electro-osmosis, 50, 68–70, 71
Endodermis, 24–28, 38
Endoplasmic reticulum (ER), 33, 41, 69

- Energy sources for active transport, 15–18
- Eosin, 40
- Epidermal cells, 25, 30, 38
- Fabaceae, 53
- Facilitated diffusion, 12
- Fagus*, 37, 51, 64
- Ferredoxin, 16
- Fluorescein, 56, 57
- Fructose, 47, 48
- Galactose, 47, 48
- Giant algal cells, 7, 11
- Gibberellins, 34, 47, 51, 52, 62
- Gland cells, 31
- Glucan, 42
- Glucose, 47, 48
- Glycine max*, *see* Soybean
- Goldman equation, 11
- Golgi apparatus, 33
- Gossypium hirsutum*, 55
- Grape, 43, 64
- Guard cells, 18–21
- Guttation, 26, 36
- Hagen-Poiseuille relationship, 63, 64, 66, 67
- Halophytes, 37–38
- Helianthus annuus*, 25, 64
- Heracleum mantegazzianum*, 55
- Hodgkin-Katz equation, 11
- Hydraulic conductivity (L_p), 26, 29, 30
- Hydrodictyon*, 15, 17
- Hydroxyl fluxes, 10, 18
- IAA (indole-3-acetic acid), *see* auxin
- Interdependent ion fluxes, 10
- Invertase, 48
- Iodine, 45
- Ion fluxes, 10–11, 15–18, 29, 36
- Ion-sensitive electrodes, 20, 24
- Iron, 51
- Kelps, 40
- Kinetic analyses, 29
- Kinetin, 29, 30
- Lateral transfer, 56–57
- Leaching, 36, 37
- Leaf nutrient content, 36–38
- Loading of translocates, 58
- Loop-path, 56, 57
- Macrocystis*, 48
- Magnesium
levels in various materials, 8, 35, 37, 47, 50
- Maize, *see* *Zea mays*
- Malate, 20, 21, 47, 52–54
- Mammalian erythrocyte, 8
plasma, 8
- Manganese, 51
- Mangrove, 37
- Mannitol, 28, 47, 48
- Mannose, 62
- Maple, 37
- Marrow, 28
- Mechanisms for phloem transport, 66–72
- Membrane potential, 9–12
- Mesophyll, 36, 38
- Metabolic inhibitors, 60
- Metasequoia glyptostroboides*, 55
- Michaelis-Menten equation, 13, 15
- Microautoradiography, 40
- Micro-peristaltic model, 70
- Mimosa pudica*, 18, 22
- Mitochondria, 15, 17, 30, 33, 41, 42
- Molybdenum, 51
- Nacreous thickening, 43
- Nernst equation, 9–10, 15
potential, 9, 35
- Nerves, 22
- Neuroid action potential, 22
- Neutral red, 41
- Nitella*, 8, 15, 16, 26, 50, 70
- Nitrate
in phloem sap, 47, 51
in xylem sap, 34, 35, 53
- Nitrogen, 34, 37
- Nucleoli, 44
- Nucleus, 41, 42, 44
- Oak, 37, 64
- Onion, 8, 30
- Organic acids, 47, 52
- Osmotic pressure, 19, 48, 54, 66
- Ouabain, 17
- Oxidative phosphorylation, 15
- Parallel model of ion uptake, 14, 15

- Partition coefficient, 10, 12
 Passage cells, 27
 Pericycle cells, 25, 30
 Permeability, 10, 12, 22
 pH
 in guard cells, 21
 in phloem saps, 54
Phaseolus vulgaris, 55
 Phloem fibres, 40
 loading, 31, 58, 59
 parenchyma, 30, 39, 41
 sap, 39, 40
 sap composition, 46–54
 un-loading, 58, 59
 Phosphates
 levels in phloem sap, 47, 50
 levels in xylem sap, 34
 Phosphoenol pyruvate (PEP), 20
 carboxylase, 21
 Phosphorus, 37
 Phytochrome, 22
 Photophosphorylation, 15
 cyclic, 16, 17
 Photoreceptor, 22
 Photosystems I and II (PS I and II), 16–17
 Pit membrane, 34, 66
 Plasma membrane (plasmalemma), 11, 14, 15, 27–30, 33
 Plasmalemma, *see* plasma membrane
 Plasmodesmata, 23, 26, 27, 34, 41, 42, 58
 Plasmolysis, 41, 45
 incipient, 19
 Plastids, 33, 41, 45, 58
 Potassium
 and electro-osmosis, 68, 69
 deficiency, 61
 fluxes, 10, 11, 15–18, 30, 59
 levels in various materials, 8, 19–22, 24, 25, 35, 37, 47, 50, 53
 partial ion conductance, 11
 permeability coefficient, 11
 sodium ratio, 15, 35
 transport across root, 27–29
 Potato, 39, 55
 P-protein (phloem protein), 41, 43–46, 67–72
 body, 44, 45
 P1, 41
 P2, 44
 Pressure-driven mass flow, 66–68, 71, 72
 Proteins, 48–50
 Proton (H^+)
 and electro-osmosis, 68
 fluxes, 10, 11, 18, 59
 gradient, 16, 18, 24, 59
 motive force, 16
 Protoplasts, 23
 Pulvinus, 21
 Pumps, 12, 15
 anion, 15–18
 cation, 15–18
 coupled, 17
 ‘Punch-through’, 12, 13
Quercus petraea, *see* Oak
 Radial barriers, 26–30
 Radial ion transport, 24
 Raffinose, 47–49, 62
 Reflection coefficient, 29
 Resorcin blue, 42
Rhizobium, 31
Rhododendron ferrugineum, 64
 hirsutum, 64
 Ribosomes, 41
Ricinus, 8, 34, 35, 46, 47, 49, 51, 52, 55, 59
 RNA, 53
 Root pressure exudation, 25, 26, 28, 34
Saccharum officinarum, *see* sugar cane
Salix, 51, 55, 67
 Salt glands, 36–38
 hairs, 37, 38
 Sea water, composition of, 8
 Semi-conductor, membrane as, 14
 Sensitive cells, 18–22
Sequoia, 67
 Series model of ion uptake, 14, 15
 Sieve, cells, 40
 element, 31, 39–62, 66–72
 element reticulum, 45
 plate, 41, 43, 45, 58, 68
 pores, 41–43, 45, 46, 71
 tubes, 40, 41, 56, 58, 68
 Sinks for translocates, 54, 57, 58, 60, 66, 67
 Sodium
 and stomatal movement, 20

- Sodium (*continued*)
 fluxes, 10, 15–18
 levels in various materials,
 8, 19, 24, 34, 35, 37, 47
 50
 permeability coefficient, 11
 transport in symplasm, 28
- Sorbitol, 47, 48
- Sources of translocates, 54, 57,
 58, 60, 66, 67
- Soybean, 52, 54
- Specific mass transfer (SMT),
 54–56
- Stachyose, 47–49, 62
- Stele, 25
- Stomata, 18–21
- Suberin, 27
- Sucrose, 47–49, 55, 61, 62, 68
- Sugar cane, 50, 55, 59–61
- Sulphate
 level in phloem sap, 47, 50
 level in xylem sap, 35
- Sulphur, 37
- Symplast, 23–31, 37, 38, 58
- System I and II ion uptake, 14,
 15
- Tensile strength of water, 64
- Tonoplast, 14, 15, 24, 41, 42,
 45
- Transcellular strands, 42, 43, 46
- Transfer cells, 30–32, 36
- Translocation rates, 54–56
- Transpiration stream, 26, 30,
 63–66
- Triticum aestivum*, 55
- Unloading of translocates, 58, 59
- Ussing-Teorell equation, 10, 15
- Verbascose, 47, 48
- Vicia faba*, 20
- Viscosity, 54, 63, 67
- Vitis vinifera*, *see* grape
- Volume flux, *see* water flux
- Water flux (volume flux),
 28–30, 63
- Water free space (WFS), 23
- Whale, 71
- White ash, 62
- Willow, *see* *Salix*
- Xanthium*, 52
- Xylem, fibres, 32, 33
 parenchyma, 25, 30, 32–34,
 38
 sap composition, 34, 35
 structure, 32–34
 tracheids, 32–33
 vessel elements, 32, 33
- Yucca*, 46, 51
- Zea mays* (Maize), 21, 24, 30, 34
- Zinc, 51