

Index

A

Atomic force microscopy, 156
Accessory olfactory system, 105
Acid-sensing ionic channels, 214
Action potentials, 226, 229, 233, 234, 236
Adenosine triphosphate, 200
Adenylyl cyclase, 107
Ageusia, 309
Amino-terminal domains, 242
Amplitude, 68, 71
Anosmia, 309
Application of electronic nose
 food evaluation, 30
 medical applications, 33
 public security, 32
Artificial-lipid sensors, 273

B

Beverages recognition and discrimination, 36
Binding pocket, 132, 133
Bioelectronic, 227
Bioelectronic nose, 46, 61, 72, 74
BioFET, 82
Bioengineered olfactory cell, 51
Biological nose, 21
 olfactory cortex neural network, 20
Biological olfaction, 168
 biomimetic smell sensor, 168
 electronic nose, 168
 sniffer dogs, 168
Biomimetic
 olfactory sensing, 319
 taste sensing, 319
Biomimetic membrane, 265
Biomimetic sensors, 265
Biosensor, 46, 47, 51, 52, 56, 57, 130, 133, 135

 cell-based biosensor, 45, 46, 49, 52, 53, 57
 smell biosensor, 45
 smell sensor, 45, 46, 51, 54, 56
Bitter, 226, 227, 232–235, 237
Brain computer interface (BCI), 310, 312
 BrainPort, 310, 311
Butanoic acid, 93

C

Caenorhabditis elegans, 10
Carbon nanotubes, 253, 258
Carbon screen-printed electrode, 208, 209
Carboxylated-polypyrrole nanotubes, 113
Central processing unit, 312
Chemical sensors, 20
 potentiometric sensors, 28
 potentiometry, 28
Chemical self-assembly, 135
Chemiresistor, 23
Chorda tympani nerve, 244
Cilia, 61–63, 66
Classification, 70, 71
Classification and prediction, 26
 linear discriminant analysis (LDA), 26
 partial least squares (PLS), 26
 regression and artificial neural nets
 (ANN), 26
CMOS, 23
Computed tomography, 11
Conducting polymer (CP) sensors, 21
Conducting polymers, 253
Conformation changes, 130, 131, 142
Cross-selectivity, 37
Cyborg, 316
 cyborg animals, 317
 cyborg nose, 315

cyborg taste, 316
 rat-cyborg, 315
 Cyclic AMP, 107
 Cyclic nucleotide-gated, 107
 Cyclic voltammetry, 251
 Cytosine, 146

D

Data processing, 276
 ANN, 277
 MRA, 277
 PCA, 277
 PLS, 277
 Deoxyribonucleic acid, 145
 Diacylglycerol, 244
 Dimension reduction, 24
 principal component analysis (PCA), 24
 Dimethyl methylphosphonate, 159
 Dinitrotoluene, 155, 158
 Duration, 68, 71

E

Electrical cell-substrate impedance sensing, 203
 Electroantennogram, 82
 Electroantennographic detectors, 97
 Electrochemical impedance spectroscopy (EIS),
 4, 113, 117, 130, 133, 140, 142, 251
 Electrochemical workstation, 136
 Electrolyte insulator [SiO₂] semiconductor
 [Si], 202
 Electroneurographic signals, 20
 Electronic nose, 2, 3, 5, 6, 9, 11, 13, 19, 46
 electronic nose systems, 20
 traditional electronic nose, 19
 Electronic tongue, 2, 5–7, 9, 11, 13, 225,
 236, 238
 Electrophysiological activity, 225, 230
 Electrophysiological recording, 227, 229–231
 Electrophysiological signals, 45, 48, 49, 53,
 66, 67, 69, 176
 β bursts, 178
 firing rate, 177
 local field potential, 176
 spikes, 176
 Epithelial Na channels (ENaCs), 226, 244
 Extracellular potentials, 64, 65, 68, 70, 72

F

Fast Fourier transformation, 229
 Field effect transistor, 4, 78, 112, 201, 248
 Firing mode, 69, 70

Firing rate, 71
 Flame ionization detectors, 97
 Flexible electrical circuit, 310
 Flow-injection analysis, 210
 Food taste measurement, 278
 Frequency spectrum analysis, 69
 Fungiform papillae, 226, 228, 230

G

Gas chromatography (GC)-mass spectrometry
 (MS), 2, 86, 122
 Gastrointestinal, 258
 Global selectivity, 310
 G-protein coupled receptor, 10, 110, 200,
 207, 215, 242
 Guanine, 146
 Gustatory epithelium, 225–238
 Gustatory receptor, 19, 34, 241, 242, 244–247,
 249, 250, 253, 255–258
 Gustatory receptor cells, 199, 200, 202, 206,
 212, 214, 220, 225, 226, 229, 230,
 232, 233, 237, 238
 Gustatory system, 27, 289–292, 296
 across-fiber pattern, 292
 BLA, 292
 brain-computer interface, 298
 chorda tympani, 291, 299
 conditioned taste aversion, 295
 greater superior petrosal, 291
 gustatory cortex (GC), 292, 301
 gustatory receptor cell, 291
 intraoral cannula, 295
 labeled line, 292
 lingual–tonsillar branch of the
 glossopharyngeal, the, 291
 local field potential, 298
 metric space analysis, 304
 PBN, 292
 rostral part of nucleus of solitary,
 the (rNST), 292
 spike, 298
 patch clamp, 296, 300
 taste bud, 291
 temporal coding, 300
 transgenic animal model, 293
 VPMpc, 292
 Gustatory tissue, 225, 237, 238

H

Hemolymph ringer solution, 86, 89
 High performance liquid chromatography,
 122, 219

- Hodgkin–Huxley, 204, 205, 214
Honeybee, 131, 140
Human embryonic kidney, 108
Hybrid sensor, 39
Hydrophobic molecules, 129
Hyperpolarization-activated cyclic nucleotide-gated cation channels, 214
- I**
Inositol triphosphate, 200, 244
Insect antenna-based smell sensors, 77, 78, 81–83, 85–87, 89–91, 93, 94, 96, 98, 99
Insect antennae, 77–82, 84–87, 89–91, 93, 94, 97, 99
Insects, 129–131, 137, 140
Integrated sensing and actuation system, 319
 actuation, 321
 biosensor chips, 320
 locomotion, 321
 myocytes, 321
 neural cells, 320
Inter digital transducer, 24, 114
Interdigitated electrodes, 133–136
Interspike interval histograms, 210
In vitro, 66, 68, 74
In vivo bioelectronic nose, 169
 craniotomy, 175
 electrical signals, 170
 main olfactory bulb, 170
 microwire array electrodes, 173
 multielectrode, 173
- L**
Langmuir–Blodgett, 117
Light-addressable potentiometric sensor, 8, 49, 51, 53, 54, 201
Limit of detection, 252
Linear discriminant analysis, 158
Lingual epithelium, 228
Lipid membranes, 266
- M**
Main olfactory bulb, 105
Main olfactory epithelium, 105
Main olfactory system, 105
MATLAB, 229
Messenger RNA, 109
Metal oxide semiconductor (MOS), 21
Microelectrode arrays (MEAs), 4, 63–68, 73, 74, 201, 227, 228, 230, 231, 238
Micro-electro-mechanical system (MEMs), 55, 227, 228, 230
Micro-fabrication, 62, 72, 73
Model, 64, 65
Molecular docking, 133, 139, 140
Monomolecular odors detection, 189
Mouse trace amine-associated receptor 5, 120
Multichannel, 65, 71, 230–232, 237, 238
Multivariate data analysis, 182
 feature extraction, 184
 K-nearest neighbor, 184
 PCA, 184
- N**
Natural odor detection, 189
 food freshness, 192
 GC/MS chromatogram, 190
 related odor pairs, 190
Nyquist plots, 137
- O**
Octadecyltrichlorosilane, 253
Odorant, 66–74
Odorant-binding proteins (OBPs), 79, 103, 129–142
Odorant detection, 178
 sensitivity, 178
 specificity, 178
 working life, 178
Odor discrimination, 69
Odors, 130
Olfaction, 45, 46, 52, 54, 57
Olfactory
 olfactory neuron, 45–47, 49, 54
 olfactory system, 46, 52, 56
Olfactory binding proteins, 4
Olfactory cell-based biosensor, 46–48, 52, 54, 56
Olfactory cell-based smell sensor, 46, 51, 53, 56, 57
 olfactory cell-based smell system, 46
Olfactory epithelium, 62–74
Olfactory receptor, 2, 4–6, 19, 20, 51–53, 55, 56, 62, 63, 68, 103–105, 107, 110, 116, 121, 124, 129, 131
 olfactory receptor cell, 46, 50, 51, 54
 olfactory receptor gene, 55
 olfactory receptor neuron, 46, 47, 51, 54, 57
 receptor cells, 61–64, 66, 68, 70
Olfactory sensory neurons, 2, 10, 103, 105
Optical sensors, 21

P

- Partial least squares, 158
- Parts per billion, 98
- PEG, 135, 136
- Peristimulus time histograms, 210
- Phenylthiocarbamide, 253
- Physical adsorption, 135
- Piezoelectric sensors, 21, 24
 - quartz crystal microbalance (QCM), 24
 - surface acoustic wave (SAW), 24
- Plasma enhanced chemical vapor deposition, 89
- Polycystic kidney disease-like, 214
- Poly(dimethylsiloxane), 208
- Poly-l-ornithine and laminin, 198
- Polymer polydiacetylene, 122
- Potentiometric sensors
 - ion-selective electrodes, 28
 - ion-sensitive field-effect transistors, 29
 - LAPS, 29
 - pH-sensitive glass electrode, 28
 - PVC membranes, 28
- Power spectrum analysis, 69, 70
- Principal component analysis, 158, 198
- Principal component regression, 158
- Process monitoring, 39
- Propylthiouracil, 255

Q

- Quality control, 278
- Quartz crystal microbalance, 4, 115, 247

R

- Randles circuit, 137

S

- Salt, 226, 227, 231–235, 237
- Seal, 64, 68, 72
- Self-assembled monolayers, 249
- Self-assembled multilayer, 116, 117
- Self-assembly, 266
- Semiochemicals, 136–140, 142
- Sensing techniques, 20
- Sensor system, 276
- Serotonin, 201, 217
- Seven helix TM domains, 242
- Shear horizontal surface acoustic wave, 314
- Signal shape, 68, 69
- Single-stranded DNA, 11, 146
- Signal-to-noise ratio, 72, 211
- Single-wall carbon nanotube, 11, 112, 122
- Single-walled carbon nanotube field effect transistors (swCN-FETs), 146, 149, 253

- Smell and taste hybrid system, 312
- Smell sensors, 61, 62, 65, 67, 68, 72, 74
- Sodium dodecyl sulfate, 247
- Solute transportation, 270
- Sonogram, 69, 70
- Sour, 226, 227, 232–235, 237
- Spatial domain, 229
- Spatio temporal analysis, 70
- Spatiotemporal pattern, 70, 71
- Specific and nonspecific molecular recognition, 19
- Spike firing, 231
- Sprague Dawley, 205
- Suppression effect evaluation, 281
- Surface acoustic wave, 4, 114
- Surface plasmon resonance, 4
- Sweet, 226, 227, 230, 232–237

T

- Tastant, 225–227, 229, 231, 232, 234–237
- Taste categories, 19
 - bitter, 19, 26
 - salty, 19, 26
 - sour, 19, 26
 - sweet, 19, 26
 - umami, 19, 26
- Taste sensors, 271
- Temporal domain, 229
- Theoretical models, 132
- Three electrodes system, 38
- Time-domain analysis, 68, 69
- Time-to-voltage conversion, 90, 91
- Transepithelial potential, 84
- Transient receptor potential cation subfamily V member 1, 244
- Trimethylamine, 120, 150

U

- Umami, 226, 227, 232–235, 237

V

- Venus flytrap, 242
- Vertebrates, 129, 140
- Volatile organic compounds, 9, 11, 22, 31, 33
- Voltammetric electronic tongue, 30, 39
- Voltammetric sensors, 29
 - voltammetry, 29
 - voltammograms, 30

W

- Water environment monitoring, 37