

SpringerBriefs in Mathematics

Series Editors

Krishnaswami Alladi

Nicola Bellomo

Michele Benzi

Tatsien Li

Matthias Neufang

Otmar Scherzer

Dierk Schleicher

Vladas Sidoravicius

Benjamin Steinberg

Yuri Tschinkel

Loring W. Tu

G. George Yin

Ping Zhang

SpringerBriefs in Mathematics showcases expositions in all areas of mathematics and applied mathematics. Manuscripts presenting new results or a single new result in a classical field, new field, or an emerging topic, applications, or bridges between new results and already published works, are encouraged. The series is intended for mathematicians and applied mathematicians.

For further volumes:

<http://www.springer.com/series/10030>

Matteo Colangeli

From Kinetic Models to Hydrodynamics

Some Novel Results

 Springer

Matteo Colangeli
Department of Mathematics
Politecnico di Torino
Turin
Italy

ISSN 2191-8198 ISSN 2191-8201 (electronic)
ISBN 978-1-4614-6305-4 ISBN 978-1-4614-6306-1 (eBook)
DOI 10.1007/978-1-4614-6306-1
Springer New York Heidelberg Dordrecht London

Library of Congress Control Number: 2013931839

Mathematics Subject Classification (2010): 76P05, 82C40, 82C70

© Matteo Colangeli 2013

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

*Cum lumen sequimur,
soror itineris tenebra*

Preface

The Boltzmann equation is a milestone of classical and quantum kinetic theory of gases [1–6]. It describes the evolution of a dilute gas, initially prepared in a nonequilibrium state, by means of a statistical approach which allows to disregard the detailed knowledge of the motion of the single particles. In the absence of external forces, the gas undergoes a process of relaxation to equilibrium, whose mathematical essence is captured by the celebrated H-Theorem, which also marked the first clear onset of irreversibility in classical particle systems. On the other hand, a different perspective in the behaviour of a fluid is offered by the laws of continuum physics, which found a self-contained settlement in the equations of hydrodynamics. This work is concerned with the ambitious and long-standing task of linking the two different levels of description, the kinetic and the hydrodynamic. The question arises as to whether this is actually an original ambition. The Grad’s moment method, for instance, which still spreads its influence on the modern theory of Extended Irreversible Thermodynamics [7], dates back to the late 1940s of the last century. Earlier attempts are represented by Hilbert’s procedure and by the Chapman-Enskog expansion, which constitute an important success in kinetic theory, as they allowed to derive the hydrodynamic laws from the Boltzmann equation and provided consistent expressions for the transport coefficients. However, almost a century of effort to extend the hydrodynamic description beyond the Navier–Stokes–Fourier approximation failed even in the case of small deviations around the equilibrium, due to the onset of instabilities which prevent the use of the hydrodynamic solutions [8, 9]. A different route, in kinetic theory, is represented by the recent Invariant Manifold method [10]. This technique, based on the computation of a slow invariant manifold in the space of distribution functions, provides hydrodynamic equations which are stable and remain valid also at short length scales, provided that the condition of local equilibrium holds. Thus, the purpose of this work is to offer a short survey over a field of active research, which aims at bridging time and length scales, from the particle-like description inherent in the Boltzmann theory up to the hydrodynamic setting. Our plan is to perform a bottom-up approach, which steps from the statistical foundations of the Boltzmann equation to the spectral properties of hydrodynamic fluctuations. Our natural

inclination is to shape things as they were rooted and naturally emerging each from the other: i.e., by showing, for instance, how typical kinetic equilibration rates affect the normal modes of hydrodynamic fluctuations and, also, how the hydrodynamic setting may be successfully extended to length scales comparable with the mean free path. A great scientist of our times, J. L. Lebowitz, expressed, in [11], his own surprise in considering that, in spite of the hierarchical structure of Nature, characterised by a variety of time and length scales, it is still possible, to some extent, to discuss the various levels of description independently of one another. “*Thus, arrows of explanations between different levels always point from smaller to larger scales, although the origin of higher level phenomena in the more fundamental lower level laws is often very far from transparent*”. Our overall impression, which we would like to share with the reader, is that a unifying approach is finally starting to take shape.

References

1. S. Chapman and T. G. Cowling, *The Mathematical Theory of Nonuniform Gases*, (Cambridge University Press, New York, 1970)
2. C. Cercignani, R. Illner and M. Pulvirenti, *The Mathematical Theory of Dilute Gases* (Springer, Berlin, 1994)
3. C. Cercignani, *Theory and Application of the Boltzmann Equation* (Scottish Academic Press, Edinburgh, 1975)
4. D. Benedetto, F. Castella, R. Esposito and M. Pulvirenti, A short review on the derivation of the nonlinear quantum Boltzmann equations, *Commun. Math. Sci.* **5**, 55 (2007)
5. D. Benedetto, F. Castella, R. Esposito and M. Pulvirenti, From the N-body Schrödinger equation to the quantum Boltzmann equation: a term-by-term convergence result in the weak coupling regime, *Comm. Math. Phys.* **277**, 1 (2008)
6. N. Bellomo Ed., *Lecture Notes on the Mathematical Theory of the Boltzmann Equation* (World Scientific, 1995)
7. D. Jou, J. Casas-Vázquez and G. Lebon, *Extended Irreversible Thermodynamics* (Springer, 2010)
8. M. Colangeli, I. V. Karlin and M. Kröger, From hyperbolic regularization to exact hydrodynamics for linearized Grad’s equations, *Phys. Rev. E* **75**, 051204 (2007)
9. M. Colangeli, I. V. Karlin and M. Kröger, Hyperbolicity of exact hydrodynamics for three-dimensional linearized Grad’s equations, *Phys. Rev. E* **76**, 022201 (2007)
10. A. N. Gorban and I. V. Karlin, *Invariant Manifolds for Physical and Chemical Kinetics*, Lect. Notes Phys. **660** (Springer, Berlin, 2005)
11. J. L. Lebowitz, Statistical mechanics: A selective review of two central issues, *Rev. Mod. Phys.* **71**, S346–S357 (1999)

Acknowledgments

I gratefully thank Prof. N. Bellomo for his support and my advisor, Prof. L. Rondoni, for contributing to shape my knowledge in this field through a continuous exchange of ideas, over the years.

I also acknowledge inspiring discussions with Prof. M. Pulvirenti and with Prof. E. Trizac.

A special mention, then, is for my family and for Alessandra, for their presence.

This work is dedicated to the memory of my beloved grandma, Gemma, who will be with me till the end.

Contents

1	Introduction	1
	References	2
2	From the Phase Space to the Boltzmann Equation	3
	2.1 The Phase Space Description	3
	2.2 The Boltzmann Equation	9
	2.2.1 H-Theorem and the Maxwellian Distribution	15
	2.3 Hydrodynamic Limit and Other Scalings	16
	2.4 Linearized Collision Integrals and Kinetic Models	18
	References	20
3	Methods of Reduced Description	23
	3.1 The Bogoliubov Hypothesis and Macroscopic Equations	23
	3.2 The Hilbert and the Chapman–Enskog Methods	28
	3.3 Grad’s Moment Method	30
	3.4 The Invariant Manifold Theory	31
	References	35
4	Hydrodynamic Spectrum of Simple Fluids	37
	4.1 Correlations in Space and Time	38
	4.2 Linearized Hydrodynamics and Collective Modes	42
	References	47
5	Hydrodynamic Fluctuations from the Boltzmann Equation	49
	5.1 Eigenfrequencies of the Boltzmann Equation	50
	5.2 The Invariant Manifold Technique	53
	5.3 Coordinate Representation and Symmetries	56
	5.4 The BGK Kinetic Model	60

- 5.5 The Maxwell Molecules Gas. 63
 - 5.5.1 Hydrodynamic Modes and Transport Coefficients. 66
 - 5.5.2 Short-Wavelength Hydrodynamics 69
- References 72

- 6 Grad’s 13-Moments System 75**
 - 6.1 Derivation of the 13-Moment System from the Boltzmann Equation 76
 - 6.2 Hydrodynamics from the Linearized One-Dimensional Grad’s System. 77
 - 6.2.1 Invariance Equations 81
 - 6.2.2 Exact Hydrodynamic Solutions. 82
 - 6.2.3 Hyperbolic Transformation for Exact Hydrodynamics . . . 85
 - 6.3 Exact Hydrodynamics from Three-Dimensional Linearized Grad’s Equations 88
- References 94

- 7 Conclusions 95**
 - References 96