

Current Topics in Microbiology and Immunology

Volume 328

Series Editors

Richard W. Compans

Emory University School of Medicine, Department of Microbiology and Immunology, 3001 Rollins Research Center, Atlanta, GA 30322, USA

Max D. Cooper

Department of Pathology and Laboratory Medicine, Georgia Research Alliance, Emory University, 1462 Clifton Road, Atlanta, GA 30322, USA

Tasuku Honjo

Department of Medical Chemistry, Kyoto University, Faculty of Medicine, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

Hilary Koprowski

Thomas Jefferson University, Department of Cancer Biology, Biotechnology Foundation Laboratories, 1020 Locust Street, Suite M85 JAH, Philadelphia, PA 19107-6799, USA

Fritz Melchers

Biozentrum, Department of Cell Biology, University of Basel, Klingelbergstr. 50-70, 4056 Basel Switzerland

Michael B.A. Oldstone

Department of Neuropharmacology, Division of Virology, The Scripps Research Institute, 10550 N. Torrey Pines, La Jolla, CA 92037, USA

Sjur Olsnes

Department of Biochemistry, Institute for Cancer Research, The Norwegian Radium Hospital, Montebello 0310 Oslo, Norway

Peter K. Vogt

The Scripps Research Institute, Dept. of Molecular & Exp. Medicine, Division of Oncovirology, 10550 N. Torrey Pines. BCC-239, La Jolla, CA 92037, USA

James L. Van Etten
Editor

Lesser Known Large dsDNA Viruses

 Springer

Editor

James L. Van Etten
Department of Plant Pathology
and Nebraska Center for Virology
University of Nebraska
Lincoln, NE 68583-0722
USA

ISBN 978-3-540-68617-0 e-ISBN 978-3-540-68618-7

DOI 10.1007/978-3-540-68618-7

Current Topics in Microbiology and Immunology ISSN 0070-217x

Library of Congress Catalog Number: 2008930849

© 2009 Springer-Verlag Berlin Heidelberg

This work is subject to copyright. All rights reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September, 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Product liability: The publisher cannot guarantee the accuracy of any information about dosage and application contained in this book. In every individual case the user must check such information by consulting the relevant literature.

Cover design: WMX Design GmbH, Heidelberg, Germany

Printed on acid-free paper

9 8 7 6 5 4 3 2 1

springer.com

Preface

Several large dsDNA-containing viruses such as poxviruses (smallpox) and herpes viruses are well known among the scientific community, as well as the general populace, because they cause human diseases. The large dsDNA insect-infecting baculoviruses are also well known in the scientific community because they are used both as biological control agents and as protein expression systems. However, there are other large dsDNA-containing viruses, including the giant 1.2-Mb mimivirus, which are less well known even though all of them play important roles in everyday life. Seven of these virus families are reviewed in this book.

Examples of their importance include the virus that causes white spot disease of shrimp (WSSV) (a single representative in the family *Whispoviridae*). WSSV is responsible for the loss of millions of dollars in the commercial shrimp farming industry every year. Likewise, some iridoviruses, such as members in the genus *Megalocytivirus*, cause serious diseases in commercial fish farms throughout Asia. Other Iridovirus members in the genus *Ranavirus* are the causative agent for approximately 50% of the documented cases of amphibian mortality reported in the United States between 1996 and 2001.

African swine fever virus (ASFV) (a single representative in the family *Asfarviridae*), which is vectored by argasid ticks and may actually be a tick virus, is usually lethal to domestic swine and outbreaks of the virus have led to large swine kills in Southern Europe, primarily in Spain.

Large dsDNA viruses that infect algae, family *Phycodnaviridae*, are also just beginning to be appreciated by the scientific community because of their influence on the global environment. That is, more than 50% of the CO₂ fixed on the planet is by marine microorganisms, called phytoplankton. The majority of these microorganisms are photosynthetic cyanobacteria. However, a significant number of these microorganisms are eukaryotic algae. Studies in the past 10 years indicate that approximately 20% or more of these photosynthetic microorganisms are infected with a virus at any one time; thus viruses are playing a large role in the turnover of these microorganisms and their role in global CO₂ fixation is only beginning to be appreciated by many marine scientists. Phycodnaviruses also contribute to the disappearance of some massive algal blooms, often referred to as red tides or brown tides.

The huge *Mimivirus*, which was originally reported to only infect protozoans belonging to the *Acanthamoeba* (amoeba) genus, may be involved in some human pneumonia-like diseases, although this role is still subject to verification. Currently, *Mimivirus* is the only member in the family *Mimiviridae*, but other relatives have been discovered recently and their genomes are being sequenced. Also, the massive genomic sequencing projects that are occurring, e.g., DNA from the Sargasso Sea, are revealing many genes that are related to *Mimivirus genes*, indicating that these large viruses are probably more common than expected.

Members of the virus family *Ascoviridae* infect insects. Ascoviruses primarily infect lepidopterous insects and they have a fascinating life style.

Some viruses reviewed in this book, specifically ASFV, *Mimivirus*, the iridoviruses, and the phycodnaviruses, along with the poxviruses, probably have a common evolutionary ancestry. Collectively, these viruses are referred to as nuclear, cytoplasmic, large dsDNA viruses, abbreviated as NCLDV. As mentioned in the chapter on the *Ascoviridae*, the ascoviruses have a strong evolutionary connection to the iridoviruses and so eventually they will probably be included as a member of the NCLDVs.

The NCLDVs are gaining the attention of some scientists interested in evolution for two reasons. First, accumulating evidence indicates that the NCLDVs may be ancient viruses; in fact, there is a suggestion that the predecessor of the NCLDVs may have existed at about the time eukaryotes and prokaryotes diverged and that these viruses might even represent the fourth domain of life. A separate cladistic study suggests that the *Phycodnaviridae* δ DNA polymerases may be near the origin of all eukaryotic δ DNA polymerases. Second, other investigators have suggested that an NCLDV predecessor may have been the origin of the nucleus in eukaryotic cells or that the nucleus is the origin of the NCLDVs.

The evolutionary origin of some of these other large dsDNA viruses is a complete mystery. For example, currently WSSV does not seem to be related to any other family of viruses. Of its predicted 181 open reading frames (ORFs), only approximately 25% match proteins in the public databases; however, another 40 of these proteins are structural proteins associated with the virus particles.

The large bacteriophage (~500 kb genome) Phage G, which infects *Bacillus megaterium*, was always considered to be an abnormality. Although, the phage was first described roughly 40 years ago, it was essentially ignored until recently. However, scientists have recently isolated more of these giant, plaque-forming bacteriophage from nature. Ironically, many of these large dsDNA bacteriophage were missed because the agar concentrations used to plaque the phage were so high that the viruses did not have time to diffuse in the soft agar on the plates.

It should also be noted that many of these large viruses, such as *Mimivirus*, are trapped in filters commonly used to remove bacteria from natural samples. Consequently, they can be easily missed unless special precautions are taken. Without doubt many more large dsDNA-containing viruses remain to be discovered.

Finally, as editor of this volume, I want to thank each of the authors for their contributions. However, only one, Greg Chinchar and his co-authors, actually met

the original deadline for their chapter and so he receives a gold medal. A special thank you goes to Chu-Fang Lo and her co-authors for agreeing at the last minute to write the chapter on the *Whispovirus*, after the original author backed out of the assignment. Finally, I want to thank Anne Clauss and her assistants at Springer for making this book happen.

Lincoln, NE, USA

Jim Van Etten

Contents

The <i>Phycodnaviridae</i>: The Story of How Tiny Giants Rule the World	1
W.H. Wilson, J.L. Van Etten, M.J. Allen	
African Swine Fever Virus	43
E.R. Tulman, G.A. Delhon, B.K. Ku, D.L. Rock	
Mimivirus	89
J.-M. Claverie, C. Abergel, H. Ogata	
Family <i>Iridoviridae</i>: Poor Viral Relations No Longer	123
V.G. Chinchar, A. Hyatt, T. Miyazaki, T. Williams	
Ascoviruses: Superb Manipulators of Apoptosis for Viral Replication and Transmission	171
B.A. Federici, D.K. Bideshi, Y. Tan, T. Spears, Y. Bigot	
Whispovirus	197
J.-H. Leu, F. Yang, X. Zhang, X. Xu, G.-H. Kou, C.-F. Lo	
Jumbo Bacteriophages	229
R.W. Hendrix	
Index	241

Contributors

C. Abergel

Structural & Genomic Information Laboratory, Parc Scientifique de Luminy,
Case 934, 13288, Marseille cedex 09, France

M.J. Allen

Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth PL1 3DH, UK

D.K. Bideshi

Department of Entomology and Interdepartmental Programs in Microbiology,
Genetics, and Molecular Biology, University of California, Riverside,
Riverside, CA 92507, USA

Y. Bigot

Université François Rabelais, U.F.R. des Sciences et Techniques, Laboratoire
d'Etude des Parasites Génétiques, FRE-CNRS 2535, Parc Grandmont,
37200 Tours, France

V.G. Chinchar

Department of Microbiology, University of Mississippi Medical Center,
Jackson, MS, USA
vchinchar@microbio.umsmed.edu

J.-M. Claverie

Structural & Genomic Information Laboratory, Parc Scientifique de Luminy,
Case 934, 13288 Marseille cedex 09, France
Jean-Michel.Claverie@univmed.fr

G.A. Delhon

Department of Pathobiology, College of Veterinary Medicine, University of
Illinois, Urbana, IL 61802, USA

J.L. Van Etten

Department of Plant Pathology and Nebraska Center for Virology,
University of Nebraska, Lincoln, NE 68583-0722, USA

B.A. Federici

Department of Entomology and Interdepartmental Programs in Microbiology,
Genetics, and Molecular Biology, University of California, Riverside, Riverside,
CA 92507, USA

brian.federici@ucr.edu

R.W. Hendrix

Pittsburgh Bacteriophage Institute and Department of Biological Sciences,
University of Pittsburgh, Pittsburgh, PA 15260, USA

rhx@pitt.edu

A. Hyatt

Australian Animal Health Laboratory, Geelong, VIC, Australia

G.-H. Kou

Institute of Zoology, National Taiwan University, Taipei 106,
Taiwan, ROC

B.K. Ku

National Veterinary Research and Quarantine Service,
Anyang Kyonggido 430-016, Korea

J.-H. Leu

Lo Institute of Zoology, National Taiwan University, Taipei 106,
Taiwan, ROC

C.-F. Lo

Institute of Zoology, National Taiwan University, Taipei 106,
Taiwan, ROC

gracelow@ntu.edu.tw

T. Miyazaki

Graduate School of Bioresources, Mie University, Mie, Japan

H. Ogata

Structural & Genomic Information Laboratory, Parc Scientifique de Luminy,
Case 934, 13288, Marseille cedex 09, France

D.L. Rock

Department of Pathobiology, College of Veterinary Medicine,
University of Illinois, Urbana, IL 61802, USA

T. Spears

Department of Entomology and Interdepartmental Programs in Microbiology,
Genetics, and Molecular Biology, University of California, Riverside,
Riverside, CA 92507, USA

Y. Tan

Department of Entomology and Interdepartmental Programs in Microbiology,
Genetics, and Molecular Biology, University of California, Riverside,
Riverside, CA 92507, USA

E.R. Tulman

Department of Pathobiology and Veterinary Science, and Center of Excellence for Vaccine Research, University of Connecticut, Storrs, CT 06269, USA
edan.tulman@uconn.edu

T. Williams

Instituto de Ecología AC, Xalapa, Veracruz, Mexico

W.H. Wilson

Bigelow Laboratory for Ocean Sciences, 180 McKown Point, P.O. Box 475, West Boothbay Harbor, ME 04575-0475, USA
wwilson@bigelow.org

X. Xu

Key Laboratory of Marine Biogenetic Resources, Third Institute of Oceanography, SOA, Xiamen, PR China

F. Yang

Key Laboratory of Marine Biogenetic Resources, Third Institute of Oceanography, SOA, Xiamen, PR China

X. Zhang

Key Laboratory of Marine Biogenetic Resources, Third Institute of Oceanography, SOA, Xiamen, PR China